

## Population dynamics of Fall armyworm *Spodoptera frugiperda* (J.E. Smith) and its Natural Enemies Maize in India

---

**Abstract** The studies on seasonal incidence of Fall armyworm (FAW) *Spodoptera frugiperda* (J.E. Smith) on Maize in India was carried out at Maize research centre Rajendranagar, Hyderabad, during Rabi 2022-23. The correlation studies revealed that the percent infestation of FAW was found to be negatively significant with maximum temperatures (-0.491<sup>\*</sup>) and positively significant with morning relative humidity (0.565<sup>\*</sup>). Correlation of natural enemies (coccinellids, spiders, earwigs) with weather parameters (Maximum temperature, Minimum temperature, relative humidity morning & evening) revealed that coccinellids, spiders and earwigs were found to be positively significant with maximum temperature and spiders found to be negatively significant with evening relative humidity. They were found to be non-significant with remaining parameters.

Keywords: Armyworm, Maize, *Spodoptera frugiperda*, cereal crop

---

### Introduction

Maize is the most important cereal crop, in India maize is the third most important cereal after rice and wheat, both in terms of area and production. The maize crop suffers from both biotic and abiotic stresses. Insect pests are among the main factors leading to lower maize yield. Over 40 species of insect pest have been identified in maize Fall armyworm (FAW) *Spodoptera frugiperda* (J.E. Smith) belonging to lepidoptera is one of the main pests in maize and known to cause reduction of yields. FAW is a polyphagous pest which is native to tropical and subtropical region of America, where it is an important pest of maize but is also known to target over 100 hosts. Moreover, it has been found that it causes significant damage to economically important crops such as rice, sorghum, sugarcane as well as horticultural crops such as cabbage, beetroot, tomato, potato and onion and also to cotton, pasture grasses, peanut, soybean, alfalfa and millets. FAW is the major pest to maize, reducing its yield to 40% in a mono cropped system. FAW had not been recorded outside of America until 2015. However, a serious outbreak of FAW was recorded from the African countries such as Sio Tome, Nigeria, Benin and Togo in 2016 and later in 2017 it was reported in Ghana. In Zambia, FAW infestation was first detected on maize in late 2016 and was reported to affect 22% of the maize grown for the 2017 crop season (IPPC 2017). In 2018, 98% of farmers reported maize to be affected in Zambia with an average maize loss of 35% (Rwomushana et al. 2018). It has now spread to majority of sub-Saharan Africa, as well as to Asia and the Middle East. The introduction of FAW into new areas has caused significant damage to crops such as maize, sorghum, and millet, which are staple foods for

millions of people in Africa. The first report of fall armyworm on maize in India is from the Shivamogga district of Karnataka during May-June, 2018 and then it was found to spread in other states like Maharashtra, Gujarat, Chhattisgarh, Andhra Pradesh, Telangana, Tamil Nadu and Odisha. This has led to significant economic losses for farmers and has threatened food security in affected countries.

Insect population fluctuations are closely tied to the biotic and abiotic factors present in their micro and macro climates (Kamata 2000). Insect populations vary throughout the year due to changes in environmental conditions such as temperature, humidity, and rainfall (Wallner 1987). Studying seasonal population dynamics in relation to these factors offers valuable insights for managing key crops (Aasman 2001). Research on the effects of temperature on insect pests is essential for comprehending their population dynamics and is instrumental in creating targeted pest control strategies (Ahmad 2016). Another critical abiotic factor affecting insect survival is humidity. Changes in humidity levels can impact the developmental stages of insects. High relative humidity (RH) can benefit the survival and growth of immature stages of some insect species (Lu & Wu 2011). The present study was conducted to know the influence of various biotic and abiotic factors affecting the population dynamics of Fall armyworm in field conditions.

## **Material and Methods**

A field study was conducted at Maize Research Centre, Rajendranagar, Hyderabad during *Rabi* 2022-23. Maize crop was sown in 100 m<sup>2</sup> plot at spacing of 60 X 20 cm. The cultivar used is DHM 117 and sowing was done on 22/11/2022. The crop was grown following all the recommended package of practices. No plant protection measures were followed for raising the crop. The data was collected on Percent infestation of fall armyworm on plants. The data on percent infestation was collected by selecting randomly 10 plants. The incidence was observed from the 1 week after the germination and continued till harvesting. The incidence was peak during 50<sup>th</sup> SMW. Observations on natural enemies like coccinellids, spiders and earwigs were also collected and correlated with weather parameters like maximum temperature, minimum temperature, morning relative humidity and evening relative humidity. The data on percent infestation of plants and natural enemies were collected in weekly intervals from randomly selected 10 plants. The correlation between weather parameters and natural enemies, Percent infestation was done by using OPSTAT. Total number of infested plants were calculated by using the following formula.

$$\text{Percent infestation} = \frac{\text{Number of plants infested}}{\text{Total number of plants}} \times 100$$

## **Results and discussion**

### **Percent infestation of plants:**

The data was collected on the percent infestation of FAW and presented in the Table 1. The observations revealed that the percent infestation of plants was observed from 1 week after emergence and continued till harvest. The highest number of infested plants was observed during 50<sup>th</sup> SMW (10 dec-16 dec) 43.48%. Weather parameters during this period were Maximum temperature (27.4°C), Minimum temperature (18.6°C), Relative humidity morning (97.1), Relative humidity evening (63.9). Lowest number of infested plants were observed during 12<sup>th</sup> SMW (1.79). Number of infested plants were in the range of 1.79% to 43.48%. percentage of infested plants were found to be less at the start of the crop but was found to reach peak when the crop is at 50<sup>th</sup> SMW.

### **Natural Enemies:**

The data on number of natural enemies like coccinellids, spiders and earwigs were collected during *Rabi* 2022-23 and presented in the Table 1. The observations revealed that the coccinellids were first observed during 50<sup>th</sup> SMW mean number of coccinellids per 10 plants were 0.1. Coccinellids were recorded throughout the season from 50<sup>th</sup> SMW to 12<sup>th</sup> SMW. Maximum number of coccinellids were observed during 7<sup>th</sup> SMW. The weather parameters during this period were Maximum temperature (33.2°C), Minimum temperature (11.7°C), Relative humidity morning (74.6 %), Relative humidity evening (17.4 %).

The observations on the mean number of spiders per 10 plants revealed that spiders were recorded throughout the season. The spiders appeared first during 50<sup>th</sup> SMW. The highest spider population was observed during 9<sup>th</sup> SMW (3.3) when the maximum temperature (33.8°C), minimum temperature (14.7°C), relative humidity morning (76.3%), relative humidity evening (19.4%). The population of spiders ranged from 0.2 to 3.3 per 10 plants.

The observations on mean number of earwigs per 10 plants revealed that the highest population of earwigs was observed during 7<sup>th</sup> SMW, where the maximum temperature (33.2°C), minimum temperature (11.7°C), relative humidity morning (74.6%), relative humidity evening (17.4%). The population of earwigs were in the range of 0.4 to 3.12 per 10 plants. The population of earwigs found to appear throughout the season.

### **Correlation between weather parameters, natural enemies and percent FAW infestation on Maize**

The data collected on natural enemies, Percent FAW infestation were correlated with weather parameters and are analysed statistically and presented in Table 2. Percent FAW infestation found to be negatively significant with maximum temperatures (-0.491<sup>\*</sup>) and positively significant with morning relative humidity (0.565<sup>\*</sup>) whereas negatively non- significant with minimum temperatures and positively non-significant with evening relative humidity.

It is evident from the results that coccinellids were found to be positively significant with maximum temperature (0.638<sup>\*\*</sup>) and negatively nonsignificant with minimum temperature, morning and evening relative humidity, percent FAW infestation.

Spiders were found to be positively significant with maximum temperature (0.820<sup>\*\*</sup>) and negatively significant with evening relative humidity (-0.555<sup>\*</sup>) whereas negatively non-significant with minimum temperature, morning relative humidity and percent FAW infestation.

Earwigs were found to be positively significant with maximum temperatures (0.688<sup>\*\*</sup>) and positively non-significant with % FAW infestation whereas negatively non-significant with minimum temperatures, morning and evening relative humidity.

Sunitha *et al.* (2021) who reported that plant infestation caused by *Spodoptera frugiperda* had shown negative significant correlation with maximum temperature and positive significant correlation with morning relative humidity, the present findings were in accordance with earlier findings. Pradeep *et al.* (2022) reported that percentage of plant infestation was positively correlated with relative humidity and negatively correlated with maximum temperature. The present findings were in close conformity with Sunitha *et al.* (2021) who reported that population of spiders showed positive significant correlation with maximum temperatures. Warkad *et al.* (2021) reported that earwig's population positive significant correlation with maximum temperature and negative significant correlation with evening relative humidity, the present findings were in accordance with earlier findings.

**Table1:** Seasonal Incidence of Fall armyworm and its natural enemies on Maize during *Rabi* 2022-23

Std week	Max temp (°C)	Min temp (°C)	RH Mrng (%)	RH Evng (%)	Mean no.of coccinellids per 10 plants	Mean no. of spiders per 10 plants	Mean no.of earwigs per 10 plants	%infestation
47	28.50	17.50	77.86	47.43	0	0	0	0
48	29.6	13.9	85.9	39.7	0	0	0	5.05
49	29.1	15.4	80.7	37.4	0	0	0	7.45
50	27.4	18.6	97.1	63.9	0.1	0.2	0.4	43.48
51	29.6	13.8	87.1	37.0	1.1	1.01	2.2	41.67
52	30.7	16.1	91.6	41.5	1.48	2.3	2.12	40.82
1	29.9	17.1	89.0	54.3	2.12	1.02	2.12	39.22
2	30.1	11.2	79.9	35.7	2.26	2.1	2.3	34.48
3	30.7	13.3	84.7	30.0	2.35	1.44	2.42	30.00
4	29.7	14.1	85.6	31.4	2.21	2.1	1.33	30.61
5	29.8	14.6	85.6	35.1	2.32	2.13	1.4	27.27
6	32.4	14.4	87.7	27.1	2.18	2.35	2.8	11.32
7	33.2	11.7	74.6	17.4	3.1	2.99	3.12	10.20
8	33.9	14.5	77.7	22.9	1.19	2.1	1.63	10.00
9	33.8	14.7	76.3	19.4	2.7	3.3	1.99	9.09
10	33.4	15.6	77.0	25.0	2.0	2.5	2.5	8.16
11	32.9	16.9	87.4	49.4	2.5	3.1	3.0	3.57
12	32.0	18.9	87.7	43.9	3.1	2.3	2.19	1.79

**Table-2:** Correlation of fall armyworm and its natural enemies on maize with weather parameters during Rabi 2022-23

Natural enemies	Temperatures		Relative humidity		% FAW infestation
	Maximum	Minimum	Morning	Evening	
Coccinellids	0.638**	-0.240	-0.181	-0.428	-0.027
Spiders	0.820**	-0.250	-0.281	-0.555*	-0.144
Earwigs	0.688**	-0.289	-0.104	-0.394	0.081
%infestation	-0.491*	-0.113	0.565*	0.323	-

### Conclusions

Percent FAW infestation was correlated negatively significant with maximum temperature and positively significant with morning relative humidity, coccinellids, spiders and earwigs are correlated positively with maximum temperature. The population of spiders are correlated negatively with evening relative humidity.

### Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. NO generative AI technologies were used by the authors
- 2.
- 3.

## References

- Aasman, K (2001) Effect of temperature on development and activity of maize stem borer *Chilo partellus*. *Bull Environ Ent* 125-127.
- Ahmad T, Hassan MW, Jamil M, Iqbal J (2016) Population Dynamics of Aphids (Hemiptera: Aphididae) on Wheat Varieties (*Triticum aestivum* L.) as Affected by Abiotic Conditions in Bahawalpur, Pakistan. *Pakistan J Zool* 48(4):1039-1044.
- IPPC (2017) IPPC ofcial pest report, (No. ZMB-02/2). Rome, Italy, FAO. <https://www.ippc.int/>
- Kumar, M., Basavaraju, B.S., Kumar, L.V., Kumar, V.S. and Gowda, P.T., 2020. Assessment of yield loss at different levels of infestation by fall armyworm, *Spodoptera frugiperda* J.E.Smith, (Lepidoptera: Noctuidae) in maize. *J. Entomol. Zool. Studies*. 8: 1018-1022.
- Lu Y, Wu K (2011) Effect of Relative Humidity on Population Growth of *Apolygus lucorum* (Heteroptera: Miridae). *Appl Entomol Zool* 46(3):421-427.
- Pradeep, P., Deshmukh, S.S., Sannathimmappa, H.G., Kalleshwaraswamy, C.M. and Firake, D.M., 2022. Seasonal activity of *Spodoptera frugiperda* (JE Smith) in maize agroecosystem of South India. *CURRENT SCIENCE*. 123(1):81.
- Rwomushana I, Bateman M, Beale T, Beseh P, Cameron K, Chiluba M et al (2018) Fall armyworm: Impacts and implication for Africa. Evidence Note Update, CAB International, Wallingford.
- Sharanbassappa., Kalleshwaraswamy, C. M., Asokan, R., Mahadeva Swamy, H. M., Maruthi, M. S. and Pavithra, H. B. 2018. First report of the fall armyworm *Spodoptera frugiperda* (J E Smith) (Noctuidae: Lepidoptera), an alien invasive pest on maize in India. *Pest Management in Horticultural Ecosystems*. 24 (1): 23-29.
- Suby, S.B., Soujanya, L.P., Yadava, P., Patil, J.K., Subaharan, Prasad, G.S., Babu, K.S., Jat, S.L., Yathish, K.R., Vadassery, J.L., Vinay, K., Kalia, N., Bakthavatsalam, J.C., Shekhar and Rakshit, S. 2020. Invasion of Fall Armyworm (*Spodoptera frugiperda*) in India: Nature, distribution, management and potential impact. *Current science*. 119(1): 44-51.
- Sunitha, S.V.L., Swathi, M., Madhumathi, T., Kumar, P.A. and Chiranjeevi, C.H., 2021. Population Dynamics of Fall Armyworm, *Spodoptera frugiperda* (JE Smith) on Sorghum. *International Journal of Environment and Climate Change*. 11(11): 222-229.
- Warkad, T.P., Bhede, B.V. and Shinde, G.S., 2021. Seasonal variations in fall armyworm *Spodoptera frugiperda* and its natural enemies on maize. *Journal of Entomological Research*. 45(4): 702-706.
- Wallner WE (1987) Factors affecting insect population dynamics: differences between outbreak and non-outbreak species. *Ann rev entomol*, 32(1):317-340.

UNDER PEER REVIEW