

Short Research Article **Assessment of Molecular Diversity in Popular Pigeonpea Genotypes using Genic Microsatellite markers**

ABSTRACT

Fusarium wilt is an important soil borne disease of pigeonpea causing significant yield losses in susceptible cultivars throughout the pigeonpea growing areas. Thirty one diverse pigeonpea genotypes were screened for wilt resistance and analysed using 14 genic SSRs which included the five SSR markers that were reported to differentiate resistant and susceptible genotypes. Polymorphic information content ranged from 0.69 to 0.90 and SSR marker ASSR-352 displayed the highest PIC value of 0.90. Dissimilarity coefficient ranged from 0.4 to 0.93. Highest similarity was observed between genotypes LRG 133-33 and TDRG 59/ICPL 99050. The 14 SSR markers generated 116 alleles and the genotypes were grouped into 4 clusters. Cluster 3 is the largest with 10 genotypes and comprises of two subclusters of wilt resistant genotypes and one subcluster of susceptible genotypes. Subcluster 2 consists of resistant and moderately resistant genotypes except PRG 176 & WRGE 93. The SSR markers used in our study could differentiate the resistant and susceptible genotypes to some extent.

Keywords: SSR, Fusarium wilt, genetic diversity, Pigeonpea, Genic SSRs

1. INTRODUCTION

Pigeonpea [*Cajanus cajan*(L.) Millspaugh] is one of the major pulse crops of India. India is the largest producer and consumer of pigeonpea in the world with 70% of the production (Bohra et al., 2012). It plays a key role in food security, balanced diet and subsistence agriculture through integrated farming systems and symbiotic nitrogen fixation (Reddy et al., 2005). Analyzing genetic relationships and studying variability is important for revealing diversity available in the species, to conduct successful breeding programs and also for germplasm conservation. Among the molecular markers many studies have reported the potential of SSR markers for studying population structure and genetic diversity because of

high polymorphism, detection of multi-allelic variation, co-dominance, reproducibility and ease of detection.

Fusarium wilt and sterility mosaic disease are the two major biotic constraints affecting pigeonpea production. Pigeonpea wilt caused by a fungus *Fusarium udum* Butler is soil borne resulting in yield losses upto the extent of 100% if the disease incidence occurs during flowering stage of crop growth (Reddy et al., 1990). Being soil borne disease, the use of fungicides is not the effective way to control the disease and development of resistant varieties is the most viable option. Four different races of *Fusarium* pathogen have been reported with pathogenic race present in Bangalore being most virulent. Wilt symptoms usually appear during flowering and podding stage of the crop. Fusarium wilt resistance in pigeonpea is controlled by different gene actions depending on the genetic backgrounds, ranging from single to multiple genes (complementary to duplicate gene action) (Saxena, 2008). Wild relatives and land races play an important role in the genetic improvement of cultivated crops especially tolerance to biotic and abiotic stresses. *Cajanus scarabaeoides* (L.) Thouars is one of the important wild species, which possess several desirable traits (Nagaraja et al., 2016).

Resistant sources for wilt disease are available, but screening requires the development and maintenance of wilt sick plot. This is not only laborious but also time consuming as it takes one entire crop season. The studies conducted by various workers suggested the presence of significant marker traits association for *Fusarium* wilt of pigeonpea (Saxena et al. 2020; Singh et al. 2016; Saxena et al., 2021). However closely linked markers associated with wilt disease or workable markers are not available to carry out marker assisted breeding. Molecular markers to differentiate the resistant and susceptible genotypes can aid the breeders in the selection of parental lines. Thirty-six elite pigeonpea genotypes adapted to different agro climatic regions have been grouped into two main clusters based on wilt reaction and six genic SSR markers were reported to be significantly associated with *Fusarium* wilt resistance (Singh et al., 2013). The inheritance of resistance to *Fusarium* wilt disease in long duration pigeonpea genotypes is governed by one dominant gene in BDN-2004-1 and BDN-2001-9, two duplicate dominant genes in BWR-133 and two dominant complimentary genes in resistance source IPA-234. The group validated the association of SSR markers ASSR-1, ASSR-23 and ASSR-148 with wilt for use in pre-screening of diverse parental lines (Singh et al., 2016). The present study was carried out to assess genetic diversity in a set of elite pigeonpea genotypes and evaluate the efficiency of recently reported SSR markers for *Fusarium* wilt screening in breeding programs.

2. MATERIAL AND METHODS

2.1 Plant material

A set of thirty one pigeonpea genotypes including released varieties, germplasm and advanced breeding lines were selected for this study. The genotypes included both wilt resistant and susceptible genotypes which were grown in pots and leaves from 20 days old seedlings were collected and frozen in liquid nitrogen (Table 1).

Table 1. List of Pigeonpea genotypes used for diversity analysis along with Fusarium wilt reaction

S.No	Genotype	FW reaction
1.	PRILB130	R
2.	ICPL20096	R
3.	PRILC 138	R
4.	ICP 7035	R
5.	ICP11015	R
6.	ICP8860	S
7.	ICPL9045	S
8.	PRILC 54	S
9.	PRILC 9	S
10.	LRG 30	S
11.	TDRG 58	R
12.	BSMR 736	R
13.	ICP7119	S
14.	LRG 133-33	S
15.	TDRG 59/ICPL 99050	R
16.	CRG 2012-25	R
17.	RVSA 16-1	S
18.	CRG 150	S
19.	AGL 1640	R
20.	ICPL 161	S
21.	WRGE 122	S
22.	ICP11230	S
23.	ICPL 2376	S
24.	WRGE 93	S
25.	TS 3R	R
26.	WRGE 121	MR
27.	PRG 176	S
28.	TDRG-4	R
29.	TDRG 107	R

30.	WRGE 120	MR
31.	ICPL 87119	R

R: Resistant; MR: Moderately Resistant; S: Suceptible

2.2 Genomic DNA Isolation and PCR using SSR markers

200mg of leaf tissue was ground into fine powder and genomic DNA was extracted following CTAB procedure (Abdelnooret al. 1995). The quality and quantity of DNA was checked through agarose gel electrophoresis (0.8%). A total of 14 ASSR primers were used for PCR amplification (Table 2). The PCR mixture comprised of 25ng of genomic DNA, 2x EmeraldAmp GT PCR Master Mix, 2.5pm of forward and reverse primers. Amplification was carried out in thermal cycler (Master Cycler, Eppendorf) by initial denaturation at 94°C for 5 min followed by 35 cycles comprising of denaturation at 94°C for 1min, optimum annealing temperature for 30 sec, primer extension at 72°C for 30 sec and final extension of 7 min. The PCR products were resolved using 3% metaphor agarose and documented in gel documentation system.

Table 2. List of polymorphic SSR markers along with primer information

S.No	Primer Code	Forward primer sequence	Reverse primer sequence	Motif
1	ASSR 277	GGAAATACAAATAGCGTAGTGG	CAGCCTAGAACTTAACACGAG	(TCCTGT) ₅
2	ASSR 304	ACTCAACGGTGCTACTCATC	AACCGATGCTTTCTTCTACTAC	(GTT) ₇
3	ASSR 66	CAAGTAAACGATTCCTCTTCTC	GTTCCCTATAGTTGATGTTTGG	(CT) ₁₂
4	ASSR 148	AACCGATGCTTTCTTCTACTAC	ACTCAACGGTGCTACTCATC	(CAA) ₇
5	ASSR 352	CACAAACCTGTAACCTTGATCC	GTGACGAGGATATACAAATTCC	(TTTAA) ₆
6	ASSR 97	GAGGATGAAGAGGAAGAGTATG	TAGCCATCATCTTGTTATGCTG	(ATGGAC) ₈
7	ASSR366	CTCTGCAACTCGCTCATTTTC	ACGTGATGGAGAAGATCCAAC	(CGT) ₈
8	ASSR 1	GTCCGTTGAAAAACAAAGAG	CGTTTTAGGTTTCTTCTCTGC	(GA) ₁₀
9	ASSR 229	ATAGTGGGACAGTAGAAAATCC	CAACTCATCTCTTGGTTCTCC	(TAAGGG) ₅
10	ASSR 23	CTTTCCCTTCTCTCTCAACAC	AAGCAGAAGCAGAAGCAGAG	(CCTTCT) ₅
11	ASSR 363	GGGAGAAGTATAAGGAGAAATG	TCACCCTTTGATAATGTTCC	(GCATCA) ₅
12	ASSR 70	TGTTAACTGTAGTGGGAGTGAG	AGAGACTACCGTAAACGAACAC	(GGTAGA) ₆
13	ASSR 317	AATCAGTGACAGTGAGGAAAAG	ACGTAACCTCCTCCTCGTTTAC	(GAGCAT) ₉

14	ASSR 228	TAGAGGAGGTGAAAAAGGATAG	TTTACTTTACCCGTACCCTTAC	(CTAAGG) ₅
----	----------	------------------------	------------------------	-----------------------

2.3 Statistical analysis

For genetic diversity studies the PCR products from SSR were analyzed by scoring qualitatively for presence (1) or absence (0). A genetic dissimilarity between the cultivars was measured by the dissimilarity coefficient. Polymorphism Information Content (PIC) was calculated according to Anderson *et al.*, (1993) using the following equation:

$$PIC_{j=1} = \sum_{i=1}^n P_i^2$$

Where, i = the i^{th} allele of the j^{th} marker, n = the number of alleles at the j^{th} marker and P = allele frequency. Hierarchical cluster analysis was performed using software DARWinver 6.0.2 (Perrier *et al.* 2003). For calculating genetic relationship among individuals, allelic data was used to generate pair wise dissimilarities using simple matching coefficients. The dissimilarity coefficients were used to construct neighbor-joining tree.

3. RESULTS AND DISCUSSION

3.1 Polymorphic information content (PIC) of SSR markers

A set of thirty one pigeonpea genotypes comprising of germplasm, advanced breeding lines and released cultivars from different states and ICRISAT were included in this study. The genotypes included wilt resistant, moderately resistant and susceptible cultivars (Annual reports, PJTSAU). Fourteen genic-SSR markers were used to analyse genetic diversity among the pigeonpea genotypes (Dutta *et al.*, 2011; Singh *et al.*, 2013). These included the six markers (ASSR-1, ASSR-23, ASSR-148, ASSR-229, ASSR-363 and ASSR-366) that were reported to have significant association with *Fusarium* wilt resistance (Singh *et al.*, 2013). The 14 SSR markers revealed 116 alleles with an average of 8.2 alleles. PIC ranged from 0.69 to 0.90 indicating the usefulness of the markers used in the study. The average PIC value for all the 14 markers was 0.82, while SSR marker ASSR-352 displayed the highest PIC value of 0.90 (Table 3). The size of the amplified products ranged from 110-272bp. Earlier studies identified a total of 83 putative alleles from the 16 polymorphic microsatellite markers with PIC ranging from 0.11 to 0.80 (Sousa *et al.*, 2011). While some studies reported very low level of polymorphism was observed in pigeonpea varieties with SSR markers (Hulluret *et al.*, 2018). Molecular characterization of five distinct pigeonpea genotypes as carried out using antioxidant (AO) enzyme [APX and SOD] and pathogenesis-related (PR) protein [CHS and β -1, 3-glucanase] families (Biswas *et al.*, 2020).

Table 3. Allelic information and Polymorphic information content of SSR markers

S.No	Primer Code	No. of alleles	Allele size range (bp)	PIC
1	ASSR 277	8	134-150	0.83
2	ASSR 304	5	123-132	0.73
3	ASSR 66	9	191-216	0.85
4	ASSR 148	7	125-138	0.83
5	ASSR 352	9	138-162	0.9
6	ASSR 97	8	144-175	0.88
7	ASSR 366	11	230-272	0.89
8	ASSR 1	8	110-121	0.89
9	ASSR 229	10	154-178	0.86
10	ASSR 23	12	149-175	0.86
11	ASSR 363	9	164-188	0.88
12	ASSR 70	7	167-185	0.70
13	ASSR 317	8	160-178	0.78
14	ASSR 228	5	138-149	0.69

3.2 Estimation of Genetic Similarities

Several studies have reported low level of molecular diversity in pigeonpea gene pool using various marker systems such as AFLP (Panguluri *et al.*, 2006), DArT (Yang *et al.*, 2006) and SSRs (Odeny *et al.*, 2007). In this study genetic similarity matrices of all twenty five genotypes were generated using genic SSR markers (Darwin version 6.0) (Perrier *et al.*, 2003). The genotypes were classified as susceptible/resistant/moderately resistant based on wilt screening at ICRISAT/ARS, Tandur locations (Annual reports of AICRP on pigeonpea). Allelic diversity data was used to generate dendrogram explaining the genetic relationships among the genotypes. The genetic dissimilarity coefficient between individuals based on SSR data ranged from 0.40 to 0.95. Minimum dissimilarity coefficient of 0.4 was observed

between genotypes LRG 133-33 and TDRG-59 of cluster IV. Similarly wilt resistant genotypes RVSA 16-1 and CRG 150 of cluster I also displayed minimum dissimilarity coefficient of 0.4. Maximum dissimilarity coefficient of 0.95 was observed between the wilt resistant genotypes TDRG 107 and PRILB-130 which belong to clusters II and III. Total soluble proteins can be employed as biochemical markers to assess the genetic variability in pigeonpea. Analysis of total soluble seed protein profiles classified the pigeonpea wild species into two major clusters with 6 out of 7 wilt resistant accessions in one cluster (Nagaraja et al., 2016). Genetic diversity in pigeonpea A and R lines was estimated using 123 SSRs and 54.47% polymorphism was reported (Sheikh et al. 2015). Genetic divergence in ninety six accessions of pigeonpea including 15 male sterile, 13 maintainer and 68 germplasm were analysed using 33 polymorphic primers. Genotypes were grouped into 8 clusters with highest dissimilarity coefficient (32%) of Pusa 991 and ULA 11 (Sharma et al., 2020).

3.3 Clustering of pigeonpea genotypes

The fourteen polymorphic primers grouped the 31 genotypes into 4 major clusters with cluster III having maximum number of genotypes (Figure 1).




Figure1. Weighted neighbor-joining dendrogram of pigeonpea genotypes depicting four major clusters. Color key: Susceptible entries (Red)

Cluster I consisted of two sub clusters: cluster IA with wilt susceptible genotypes, cluster IB comprised of all wilt resistant and susceptible genotypes. Cluster II consisted of all wilt resistant genotypes except PRG 176 & WRGE 93. Pigeonpea entries developed at RARS, Warangal (WRGE 93, WRGE 121, and WRGE 120) were grouped together in cluster II indicating higher genetic similarities. Cluster III comprised of ten genotypes which were further classified into three distinct sub clusters. The SSR markers could clearly differentiate the wilt susceptible and resistant genotypes. The susceptible genotypes were grouped into IIIA while IIIB and IIIC comprised of wilt resistant genotypes. Cluster IV comprised of all wilt susceptible genotypes except TDRG 59 & CRG 2012-25 which is a susceptible genotype (Table 4).

Table 4. Clustering of pigeonpea genotypes based on SSR markers

Clusters	Genotypes
Cluster 1	ICPL 161, ICP11230, ICPL 2376, WRGE 122 (all susceptible)
	RVSA 16-1, CRG 150, AGL 1640, BSMR 786
Cluster 2	WRGE 93, WRGE 121, WRGE 120, PRG-176, TS3R, TDRG 4, TDRG 107, ICPL 87119 (resistant & moderately resistant except PRG 176 & WRGE 93)
Cluster 3	ICP8860, ICPL 9045, PRIL C 54, PRILC 9 (all susceptible)
	PRILB130, ICPL 20096, PRIL C 138 (all resistant)
	ICP 7035, ICP11015, TDRG 58 (all resistant)
Cluster 4	ICP 7119, LRG 133-33, TDRG 59/ICPL 99050, CORG 2012-25, LRG 30

The SSR markers used in this study could differentiate the wilt susceptible and resistant genotypes to a major extent indicating their usefulness in the selection of parental lines for breeding program. Principle component analysis (PCA) also, classified the pigeonpea genotypes into 4 distinct clusters. The grouping of the genotypes remained almost the same

except for genotypes LRG 30 and LRG-133-33 of cluster IV which were shifted to cluster III. Molecular marker correlated with fusarium wilt will aid the breeders in selection of parental lines for resistance breeding in African and Asian Pigeonpeas (Mayomba, 2015). Saxena et al., (2021) have reported SNPs/Indels for Fusarium wilt and sterility mosaic diseases resistance in pigeonpea. Five SSR markers from earlier studies (Singh et al., 2013) - ASSR 363, ASSR 366, ASSR 1, ASSR 23 and ASSR 148 could clearly differentiate a set of 19 wilt resistant and susceptible genotypes with marker ASSR 363 having highest PIC (Bisht et al., 2022).

4. CONCLUSION

Genetic diversity in 31 elite pigeonpea Fusarium wilt resistant and susceptible genotypes was analyzed using 14 genic-SSR markers. ASSR-352 is the most useful marker in discerning genetic diversity. Fusarium wilt resistant genotypes of divergent clusters are useful in selection of parental lines hybridization program and also for mapping of wilt resistance in pigeonpea.

REFERENCES

- Bisht C, Verma SK, Gaur, AK, Chauhan C, Deep H, Karn A, Sharma. Characterization of Elite Genotypes for *Fusarium* Wilt Resistance in Pigeonpea [*Cajanus cajan*(L.) Millspaugh] Legume Research. 2022; 45 (1): 116-121.
- Biswas K, Tarafdar A, Kumar R, Singhvi N, Ghosh P, Sharma M, Pabbi S, Shukla P. Molecular Analysis of Disease-Responsive Genes Revealing the Resistance Potential Against Fusarium Wilt (*Fusarium udum*Butler) Dependent on Genotype Variability in the Leguminous Crop Pigeonpea. Frontiers in Genetics. 2020; 11: 862.
- Bohra A, Saxena RK, Gnanesh BN, Saxena KB, Byregowda M, Rathore A. *et al.* An intra-specific consensus genetic map of pigeonpea [*Cajanus cajan*(L.) Millspaugh] derived from six mapping populations. Theor. Appl. Genet. 2012;125:1325–1338.
- Burns MJ, Edwards KJ, Newbury HJ, Ford-Lloyd BV, Baggott CD. Development of simple sequence repeat (SSR) markers for the assessment of gene flow and genetic diversity in pigeonpea (*Cajanus cajan*). Mol Ecol Notes. 2001; 1:283–5.
- Dutta S, Kumawat G, Singh BP. *et al.* Development of genic-SSR markers by deep transcriptome sequencing in pigeonpea [*Cajanus cajan* (L.) Millspaugh]. BMC Plant Biol. 2011; 17 (2011).
- Hullur N, Radha BN, Basavaraja B, Channakeshava BC, Byregowda M. Genetic diversity assessment in pigeonpea cultivars using microsatellite (SSR) markers. International Journal of Current Microbiology and Applied Sciences. 2018; 7(3): 3268-3274.

Mayomba M. Characterization and incorporation of fusarium wilt disease resistance in African and Asian Pigeonpeas [*Cajanus cajan*(L.) Millsp.] Germplasms. Doctoral dissertation, Sokoine University of Agriculture, Tanzania (2015) pp. 166.

Nagaraja NR, Arun Kumar MB, Manjunath Prasad CT, Rajkumar, Sinha P, Jaagrati J. Genetic Diversity Studies and Screening for Fusarium Wilt (*Fusarium udum*Butler) Resistance in Wild Pigeonpea Accessions, *Cajanus scarabaeoides*(L.) Thouars. Indian J. Plant Genet. Resour. 2016; 29(2): 121–129.

Odeny DA, Jayashree B, Ferguson M, Hoisington D, Crouch J, Gebhardt C. Development, characterization and utilization of microsatellite markers in pigeonpea. Plant Breed 2007; 126:130–6.

Panguluri SK, Janaiah J, Govil JN, Kumar PA, Sharma, PC. AFLP fingerprinting in pigeonpea [*Cajanus cajan*(L.) Millsp.] and its wild relatives. Genet. Res. Crop. Evol. 2006; 53: 523-531.

Reddy LJ, Upadhyaya HD, Gowda CLL, Singh, S. Development of core collection in pigeonpea [*Cajanus cajan*(L.) Millspaugh] using geographic and qualitative morphological descriptors. Genet. Resour. Crop Evol. 2005; 52:1049–1056.

Sharma P, Singh I, Sirari A, Singh S, Khosla G. Genetic divergence studies through microsatellite markers in pigeonpea [*Cajanus cajan*(L) Millsp.]. Legume Research. 2020;43 (3): 312-319.

Sheikh W, Acharya S, Patel JB, Kalaskar SR, Shinde AS, Patel KA. Genetic fingerprinting of A and R lines of pigeonpea [*Cajanus cajan*(L.) Millsp.] using RAPD and SSR markers. Indian J. of Biotechnology. 2015; 14: 328-333.

Saxena RK, Kale S, Mir RR, Mallikarjuna N, Yadav P, Das RR, Molla J, Sonnappa M, Ghanta A, Narasimhan Y, Rathore A, Sameer Kumar CV, Varshney R.K. Genotyping-by sequencing and multilocation evaluation of two interspecific backcross populations identify QTLs for yield-related traits in pigeonpea. Theoretical and Applied Genetics. 2020; 133: 737-749.

Singh D, Sinha B, Rai VP, Singh MN, Singh DK, Kumar R, Singh A.K. Genetics of *Fusarium* wilt resistance in pigeonpea (*Cajanus cajan*) and efficacy of associated SSR markers. Plant Pathology Journal 2016; 32(2): 95-101.

Singh AK, Rai VP, Chand R, Singh RP, Singh MN. Genetic diversity studies and identification of SSR markers associated with *Fusarium* wilt (*Fusarium udum*) resistance in cultivated pigeonpea (*Cajanus cajan*). Journal of Genetics. 2013; 92: 273-280.

Saxena RK, Hake A, Bohra A, Khan AW, Hingane A, Sultana R, Singh IP, Naik SJS, Varshney RK. A diagnostic marker kit for Fusarium wilt and sterility mosaic diseases resistance in pigeonpea. Theoretical and Applied Genetics. 2021;134(1): 367-379.

Sousa CA, Godoy R, Sforca AD, Campos T, Zucchi M, Jank L, Souza A. Genetic diversity analysis among pigeonpea genotypes adapted to South American regions based on microsatellite markers. Scientia Agricola (Piracicaba, Braz.) 2011; 68(4): 431-439.

Yang S, Pang W, Harper J, Carling J, Wenzl P, Huttner E, Zong X, Kilian A. Low level of genetic diversity in cultivated pigeonpea compared to its wild relatives is revealed by diversity arrays technology (DArT). *Theor. Appl. Genet.* 2006; 113: 585-595.

UNDER PEER REVIEW