

Effect of Different Growth Regulators on the Rooting of Hardwood Cuttings of *Bougainvillea glabra* var. "Mohan"

Abstract

The *Bougainvillea* plant is a common climber and shrub, and one of the most difficult tasks for its growers is cutting roots. It's nice that this study employed three different kinds of auxins to root the cuttings. *Bougainvillea*, a popular decorative plant, is generally propagated by cuttings and tissue culture. However, large-scale cultivation of *Bougainvillea* is sometimes problematic due to roots concerns. An experiment about the effect of different growth regulators on the rooting of hardwood cuttings of *Bougainvillea glabra* var. "Mohan" was conducted during the year 2020-21 at the Faculty of Horticulture, Bidhan Chandra Krishi Vidyalaya, Mohanpur, Nadia, West Bengal, India. The study employed a completely randomized design with ten treatments (T₁: IAA @500ppm, T₂: IAA @1000ppm, T₃: IAA @1500ppm, T₄: IBA @500ppm, T₅: IBA @1000ppm, T₆: IBA @1500ppm, T₇: NAA @500ppm, T₈: NAA @1000ppm, T₉: NAA: 1500ppm & T₁₀: control), 210 hardwood cuttings of *Bougainvillea* var. "Mohan" were made, and fungicide was applied before planting. Black polythene was placed beneath the bed for quicker roots. IAA, IBA, and NAA solutions were created using distilled water. Cuttings were treated with growth regulator and planted in beds, each replicated three times. The findings of this investigation revealed that among the various concentrations of IAA-Indole acetic acid, IBA-Indole-3-butyric acid, and NAA-Naphthalene acetic acid, hardwood cuttings treated with IAA at 1500 ppm exhibited the earliest bud initiation (7 days). The best result for the first root initiation (7 days) was observed with IAA at 1000 ppm. IBA at 1500 ppm resulted in the highest number of leaves per cutting (25), while IBA at 1000 ppm produced the highest number of buds per cutting (4). Furthermore, the highest number of roots (9.2) and the greatest average root length (8.7 cm) were achieved with IBA at 1500 ppm. From the foregoing discussion and summary, it can be concluded that efficient management of IBA in higher concentration i.e. 1500 ppm gave the best result in rooting of Hardwood cuttings of *Bougainvillea* var. "Mohan" for better rooting.

Keywords: *Bougainvillea*, Mohan, Hardwood cuttings, Rooting and PGRs

Introduction

Bougainvillea belongs to the dicot family Nyctaginaceae and is native to tropical and subtropical South America, making it well-adapted to similar climatic conditions. The *Bougainvillea* plant is a popular climber and shrub, and one of the most challenging chores for its cultivators is root removal. It's wonderful that this study used three distinct types of auxins to root the cuttings. *Bougainvillea*, a popular ornamental plant, is often grown by cuttings and tissue culture. However, large-scale production of *Bougainvillea* might be troublesome owing to root issues.

There are about 10 species of *Bougainvillea*, but only three are horticulturally important: *B. spectabilis*, *B. peruviana*, and *B. glabra*. This plant can tolerate both semi-shaded areas and full sun, though it thrives best in full sunlight. A minimum of 5 hours of direct sunlight per day is required for good blooming, and more hours of sunlight are even

better. Bougainvillea does not require a lot of water and is prized by gardeners for its wide range of habitats, prolonged flowering seasons, and variety of flower colours. It also has numerous uses in landscaping. (Bhardwaj and Kumar, 2020).

Bougainvillea blooms in the winter, starting in October when the weather becomes cool and pleasant, with a second flush of flowers in the summer. However, some varieties, like Snow Queen and Snow-White, only bloom once during the winter. The plant can be used as a shrub or climber and can be trained to grow in pergolas. It can grow over 30 feet high, with large, simple, mostly ovate leaves that may have rippling edges and hairs underneath. The flowers are usually surrounded by three brightly coloured bracts, while the actual flowers are small, tubular, and often inconspicuous, located in the centre of the bracts. Bract colours include white, light mauve, magenta, pink, deep mauve, orange, yellow, and red. Bougainvillea has a fragile root system that does not form a good root ball and has many adaptations to its climate and environment. (Bhardwaj and Kumar, 2020)

Betacyanin's and flavanols are extracted from the bracts of *B. glabra* (Heuer *et al.*, 1994), and pinitol, an anti-diabetic compound, has been isolated from the leaves of *B. spectabilis* (Narayanan *et al.*, 1987). Bougainvillea is a valuable ornamental plant with culinary uses and is also utilized in traditional medicine for treating common ailments. It is traditionally used against several diseases, including diarrhoea, hypotension, intestinal disorders, stomach-ache, nausea, inflammation-related ailments, and pain management (Saleem *et al.*, 2021). The leaves have anti-inflammatory potential (Giri *et al.*, 1988), and traditional practitioners in Mandsaur, India, use the leaves to treat various gastrointestinal disorders like diarrhoea and acidity. Additionally, the anti-diarrheal activity of Bougainvillea *glabra* is related to its antimicrobial properties.

Adventitious root formation is a crucial step in the vegetative propagation of woody or horticultural species, and problems associated with rooting cuttings can result in significant economic losses. The application of exogenous auxin/rooting hormones can help overcome these challenges. Commercially available exogenous auxins that aid in the formation of adventitious roots include Indole-3-acetic acid (IAA), Indole-3-butyric acid (IBA), and Naphthalene acetic acid (NAA). Naturally occurring auxins like IAA and IBA favor apical dominance, help control xylem differentiation, and aid in cell division. NAA is a synthetic plant hormone in the auxin family and is an ingredient in many commercial plant rooting products. IAA and IBA are also used commercially in the field of tissue culture. It is used as a rooting agent for the vegetative propagation of plants from stem and leaf cuttings and in plant tissue culture (Memon *et al.*, 2013). The study looks at how different Auxin concentrations (IAA, IBA, and NAA) affect the rooting of Bougainvillea var. Mohan hardwood cuttings, which are used to treat a variety of ailments and traditional treatments such as diarrhoea, hypotension, and inflammation.

Materials and methods

The experiment was conducted at a site in Jalpaiguri, West Bengal, India, using hardwood cuttings of *B. glabra* var. Mohan collected from the premises (26°32'06.9" N 88°43'37" E), of the Faculty of Horticulture, Bidhan Chandra Krishi Vidyalaya, Mohanpur, Nadia, West Bengal, India. The hardwood cuttings, each 15-20 cm in length and with the thickness of a pencil, were prepared in January. A quick dip treatment (10 seconds) at the basal portion with different concentrations of IBA (Indole-3-butyric acid) and NAA (Naphthalene acetic acid) was applied. The stock solutions were prepared by dissolving 1 mg/l of NAA or IBA in distilled water, using ethyl alcohol to aid dissolution.

The experiment followed a Randomized Block Design (RBD) with three replications and ten treatments as follows:

- T1: IAA 500 ppm
- T2: IAA 1000 ppm
- T3: IAA 1500 ppm
- T4: IBA 500 ppm
- T5: IBA 1000 ppm
- T6: IBA 1500 ppm
- T7: NAA 500 ppm
- T8: NAA 1000 ppm
- T9: NAA 1500 ppm
- T10: Control (No chemical)

A total of 210 hardwood cuttings of Bougainvillea var. "Mohan," each 15-20 cm long and of pencil thickness, were prepared. The beds were drenched with fungicide before planting the cuttings, and black polythene was used below the bed for bottom heating to encourage faster rooting. Local Coarse Sand is used as rooting media and the climate was sub-tropical, Rainfall in July August. IAA solutions of 500 ppm, 1000 ppm, and 1500 ppm were prepared by dissolving 0.5 g/l, 1 g/l, and 1.5 g/l, respectively, in distilled water. IBA and NAA solutions were prepared in the same manner.

The observations recorded included:

- Days taken for first bud initiation in cuttings
- Days taken for first rooting of cuttings
- Number of leaves per cutting
- Number of buds per cutting
- Number of roots per cutting
- Length of roots in cuttings

Final data for the parameters (number of leaves, number of buds per cutting, number of roots per cutting, and length of roots) were recorded after 8 weeks. Data were also collected in the 1st, 3rd, 4th, 6th, and 8th weeks after planting the cuttings.

Statistical analysis

The study used a Randomised Block Design to analyze data from field experiments or lab studies, assessing treatment-related variation using Gomez and Gomez's (2010) 'F' values, and determining $S.Em \pm$ using Excel statistical analysis.

4.RESULTS AND DISCUSSIONS

The experiment concluded that treatment T₃ (IAA 1500 ppm) showed the earliest bud initiation, taking the fewest days (7) for the first bud to appear. Treatments T₉ (NAA 1500 ppm) and T₇ (NAA 500 ppm) produced similar results, with the first buds appearing in 8 and 9 days, respectively. In contrast, the control treatment T₁₀, which did not use any hormones, resulted in late bud initiation (15 days). This suggests that the exogenous application of auxin helps break down starch into simple sugars, which are essential for new cell production and

increased respiratory activity during the regeneration of tissue at the time of new primordial initiation. These results are consistent with the findings of Kumar *et al.* (2014) in carnation.

For the days taken for first rooting of cuttings, treatment T₂ (IAA 1000 ppm) gave the best results, with root initiation starting very quickly (7 days). However, this treatment did not produce a large number of roots later on. Treatments T₄ (IBA 500 ppm), T₆ (IBA 1500 ppm), T₃ (IAA 1500 ppm), and T₅ (IBA 1000 ppm) also performed well, with root initiation occurring in 8 to 9 days. This indicates that the appropriate concentration of plant growth regulators can enhance cell division, cell elongation, and early differentiation of callus tissue toward root formation, resulting in early growth in cuttings. These findings align with the observations of Netam *et al.* (2018) and Bhardwaj and Kumar (2020) in bougainvillea.

The highest number of leaves was observed in treatment T₆ (IBA 1500 ppm) with 25 leaves. Treatments T₅ (IBA 1000 ppm) and T₄ (IBA 500 ppm) also performed well, with 22 and 19 leaves, respectively. However, treatments T₁ (IAA 500 ppm) and T₇ (NAA 500 ppm) produced fewer leaves (13 and 12, respectively), and the control treatment T₁₀ had the fewest leaves (10). The highest number of buds per cutting was found in treatment T₅ (IBA 1000 ppm) with 4 buds, followed closely by T₄ (IBA 500 ppm) and T₆ (IBA 1500 ppm). The control treatment performed poorly in this regard. This can be attributed to IBA's ability to produce healthier and longer roots, which absorb more nutrients and water, resulting in more leaves. Additionally, the vigorous growth and early root initiation induced by growth regulators allow for better nutrient absorption and increased leaf production. These results are consistent with the findings of Sahariya *et al.* (2013) and Kale and Bhujbal (1972) in bougainvillea.

Treatment T₆ (IBA 1500 ppm) also produced the highest number of roots and had the best average root length. Treatments T₄ (IBA 500 ppm) and T₅ (IBA 1000 ppm) also performed well. In contrast, the control treatment had poor rooting results. This suggests that IBA increases cell wall plasticity, cell division, callus development, and root growth. Cuttings treated with the appropriate concentration of auxin showed early and better root initiation, leading to a higher number of roots per cutting. Increased root length in IBA-treated cuttings may be due to enhanced carbohydrate hydrolysis, new protein synthesis, cell enlargement, and cell division induced by auxins. Auxins also initiate the synthesis of structural enzyme proteins involved in adventitious root formation, increasing root length through acidification. These results align with the findings of Singh *et al.* (2017) in bougainvillea and Bhatt and Chauhan (2012) in marigold.

Conclusion

The experiment aimed to study the effect of growth regulators on the sprouting and rooting of Bougainvillea hardwood cuttings, and the findings were significant. It was observed that IAA at 1500 ppm resulted in the earliest bud initiation, while NAA at both 1500 ppm and 500 ppm produced similar results for bud initiation. For root initiation, IAA at 1000 ppm gave the best results. The highest number of leaves were seen in cuttings treated with IBA at 1500 ppm, and the most buds per cutting were observed with IBA at 1000 ppm. In terms of root development, the number of roots and the average root length were highest in cuttings treated with IBA at 1500 ppm. These results indicate that IBA at a higher concentration (1500 ppm) yielded the best overall outcomes for rooting Bougainvillea hardwood cuttings var. "Mohan," demonstrating a significant positive impact on both shoot and root growth.

Table 1. Effect of different growth regulators on the rooting parameters of *B. glabra* var. Mohan

Treatment	Days taken to first bud	Days taken to first rooting of cutting	Number of leaves per cuttings:	Number of buds per cuttings
T ₁	12	8	13	3
T ₂	13	7	18	2.7
T ₃	7	9	15	2.5
T ₄	10	8	19	3.3
T ₅	11	9	22	4
T ₆	11	8	25	3.3
T ₇	9	11	12	3
T ₈	12	13	14	3.2
T ₉	8	12	17	3
T ₁₀	15	15	10	2
SEm	0.80	0.87	1.55	0.18
CD @ 5%	2.31	2.60	4.48	0.51

Table 2. Effect of different growth regulators on the number of roots and their length

Treatment	Number of roots per cuttings	Avg. root length (cm)
T ₁	7	3.5
T ₂	8.2	5.2
T ₃	8.9	3
T ₄	9	5
T ₅	10	8.5
T ₆	15	8.7
T ₇	8.6	3
T ₈	9.2	3.5
T ₉	8	4
T ₁₀	5	3.2
SEm	0.85	0.72
CD @ 5%	2.46	2.09

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

- Bhardwaj, D. R and Kumar M. Comparative evaluation of hardwood and semi-hardwood cutting with different rooting hormone in (*Bougainvillea buttiana*) cv. Mahara. *Int. J Chem Std.* 2020;8(5): 606-610.
- Bhatt, S. T., & Chouhan, N. M. (2012). Effect of auxin on rooting of African Marigold (*Tegeteserecta* L.). *Advanced Res. J. Crop Improvement*, 11(3), 69-70.
- Giri, S. N., Biswas, A. K., Saha, B. P., Pal, S. P., & Pal, M. (1988). Studies on the anti-inflammatory action of *Bougainvillea glabra* leaves. *Indian J. Pharmaceut. Sci.*, 50(1), 42.
- Gomez, A.K., and Gomez, A.A. (2010). Statistical procedures for agricultural research. 2nd edn. Wiley India Pvt. Ltd., New Delhi, pp. 134-138.
- Heuer, S., Richter, S., Metzger, J. W., Wray, V., Nimtzt, M., & Strack, D. (1994). Betacyanins from bracts of *Bougainvillea glabra*. *Phytochemistry*, 37(3), 761-767.
- Kale, P. N., & Bhujbal, B. G. (1972). Use of growth regulators in rooting of cuttings of bougainvillea var. Mary Palmer. *Indian J. Horti.*, 29(3and4), 307-309.
- Kumar R, Ahmed N, Sharma OC, Lal S. Influence of auxins on rooting efficacy in Carnation (*Dianthus caryophyllus*L.) cuttings. *J Hortl. Sci.* 2014;9(2):157- 160.
- Memon N, Ali N, Baloch M, Chachar Q. Influence of naphthalene acetic acid (NAA) on sprouting and rooting potential of stem cuttings of Bougainvillea. *Sci. Int.* (Lahore). 2013;25(2):299-304
- Narayanan, C. R., Joshi, D. D., Mujumdar, A. M., & Dhekne, V. V. (1987). Pinitol—a new anti-diabetic compound from the leaves of *Bougainvillea spectabilis*. *Current Science*, 56(3), 139-141.
- Netam, N., Shukla, N., Sharma, G., & Sahu, J. K. (2018). Effect of Different IBA concentration on survivability and rooting of Jasmine (*Jasminum sambac* (L.) Aiton) stem cuttings. *J.Pharmacog.Phytochem.*, 7(1S), 614-617.

- Sahariya, K. S., Singh, J. N., & Singh, A. S. (2013). Studies on the effect of IBA on rooting of bougainvillea (var. Thimma) cuttings in open field and polyhouse conditions.
- Saleem, H., Usman, A., Mahomoodally, M. F., & Ahemad, N. (2021). Bougainvillea glabra (choisy): A comprehensive review on botany, traditional uses, phytochemistry, pharmacology and toxicity. *Journal of Ethnopharmacology*, 266, 113356.
- Singh, B., Sindhu, S. S., Yadav, H., & Saxena, N. K. (2017). Influence of growth hormones on hardwood cutting of Bougainvillea cv. Dr HB Singh. *Chem. Sci. Rev. Lett*, 6(23), 1903-1907.

UNDER PEER REVIEW