

Original Research Article

Evaluation of different Agro-based waste extract media for *Lentinula edodes*

Comment [MK1]: SUGGESTED TITLE: *In vitro* cultivation of Shiitake mushrooms (*Lentinula edodes*) on the compost of some selected agro-based waste extracts.

Abstract

Shiitake mushrooms (*Lentinula edodes*), known for growing on decaying deciduous trees, are particularly notable for their health benefits. Recent advancements in cultivation techniques involve growing *Lentinula edodes* on sterile substrates to boost yield and shorten the growth cycle. In Telangana state, the climate and agricultural waste provide optimal conditions for mushroom cultivation, which can boost the local economy and nutrition. This study aimed to determine the *in vitro* performance of Shiitake mushrooms (*Lentinula edodes*) using S1 to S17 combinations of 17 types of agro-based waste extract media spawned on synthesized PDA and incubated at temperatures of 20°C and 24°C, to evaluate the best substrate extract media for shiitake mushroom production. At 20°C, the maximum mycelium growth of *L. edodes* was recorded on S17 (PDA, 75.33mm) on 7 DAI, while minimum mycelium growth was recorded on sawdust mixed with paddy straw (S8, 35.67mm). Similarly at 24°C, each of the substrates S17 (PDA) and S10, S11, and S12 (eucalyptus bark supplemented with wheat grains and sorghum grains) recorded the maximum mycelium growth (90mm), and S8 (sawdust mixed with paddy straw) recorded minimum mycelium growth (30.33mm) on 7 DAI. The results from the Principal Component Analysis of recorded data values of the mycelial growth was shown in the form of Eigen values, which shows that the highest Eigen value was recorded for mycelial growth of which was recorded in *L. edodes* grown on 7 DAI and 14 DAI at 24°C, indicating significant influence on substrates.

Comment [MK2]: Delete. Not necessary here

Keywords: *Lentinula edodes*, mushroom compost, substrates, mycelium growth, media, temperature, *in vitro* cultivation, agro-based waste.

Comment [MK3]: Write as eigenvalues.

Comment [MK4]: Delete

Comment [MK5]: What are these? What do they stand for?

Comment [MK6]: Please, recast to give more meaning to your conclusion. You can use the word "compost"

Formatted: Font: Italic

Introduction

Shiitake mushroom (*Lentinula edodes*) is one of the six popular edible mushrooms native to the East Asian region. The practice of cultivating shiitake mushrooms can be traced back to ancient times in the Far East, particularly in areas such as China, and Japan. The generic name *edodes* comes from Latin, meaning "edible". *L. edodes* grows in groups on decaying deciduous trees, particularly on the *Castanopsis cuspidata* species. They are white wood rotting fungi capable of decomposing the cellulose and lignin structural components (Annepuet *et al.*, 2019 and Kobayashi *et al.*, 2020). Shiitake mushrooms are prized for being fat-free and having low cholesterol and sodium content, making them a healthy dietary choice. They are also rich in proteins, lipids, carbohydrates,

fibers, ergosterols, antioxidants, and vitamins like provitamin D, which are not commonly found in other food sources (Paswalet *et al.*, 2024). Due to their exceptional nutritional value and potential medical benefits, this species is often referred as "the queen of plants." (Chen *et al.*, 2017). Season-based cultivation of these mushrooms can be taken up with favourable climatic conditions. Additionally, the abundance of agricultural wastes from local farms can be converted into prized mushroom production. Despite efforts to standardize its cultivation, this mushroom has not yet been commercially produced on large scale ever in India, with no reports of its cultivation in Telangana. Mushroom cultivation helps recycling agricultural wastes and addresses nutritional deficiencies and health issues. This study was taken up to evaluate the suitable media and growth performance of shiitake mushrooms at different temperatures prevailing in Telangana state on some selected agricultural wastes.

Comment [MK7]: Ref not listed

Comment [MK8]: Suggestion: Reports on its cultivation in Telangana is also not available.

Comment [MK9]: Be precise about the temperature

Comment [MK10]: Mention the details of the climatic conditions and the month of this study.

Comment [MK11]: Why not study site?

Experimental site

The experiment trial was conducted in the Mushroom Cultivation Scheme, Department of Plant Pathology, College of Agriculture, Rajendranagar, PJTSAU, Telangana State, India. This study was supported by the Central Instrumentation Cell, which provided essential resources and facilities for the research.

Materials and Methods:

Pure culture

The shiitake mushroom (*Lentinula edodes*) culture was procured from the IIHR, Bangalore. The culture was grown on sterilized Petri plates containing Potato Dextrose Agar (PDA) media in a BOD incubator at a temperature of 25±2°C. The fully grown 7 days old culture was then transferred onto fresh PDA slants, and after full growth, they are allowed to fully grow and then stored in a refrigerator at 10-12°C for future use.

Comment [MK12]: Suggested sub-title: Preparation of pure tissue culture of *Lentinula edodes*

Pouring, incubation, and data collection

In each 500 ml conical flask, 17 different types of the prepared media was filled up to two-thirds of its volume. The flask was then tightly sealed-plugged with non-absorbent cotton plugs wool, and covered with aluminium foil, which was secured with rubber bands. Further, the flasks were autoclaved for 20 minutes, sterilized at 121.6°C and 15 psi pressure, autoclaved for 20 minutes. After sterilization, the media was poured into 90mm sterilized petri plates, with 20 ml per plate for every treatment, three replications were done within a laminar air flow cabinet. The centres of the Petri plates were inoculated with 7mm discs of 7-day-old mushroom culture at the centre of the Petri plate within the laminar air flow chamber and incubated at two different temperatures (20°C and 24°C) in a BOD incubator. The growths of the mushroom mycelium was recorded after 7 DAI along with the mycelium grown in the control plate.

Comment [MK13]: Suggested sub-heading: Preparation of the agro-based waste extract

Comment [MK14]: You can present this in form of experimental design just before this section

Comment [MK15]: Each of the 17 different types of the prepared media was filled up to two-third of 500 ml conical flask.

Comment [MK16]: In three replicates, 20 mL of each of the prepared media was aseptically poured into 90 mm sterilized petri plates inside a laminar air flow cabinet.

Comment [MK17]: Do you mean 7 days?

Comment [MK18]: Mention the components of your control plates.

Preparation of Potato Dextrose Agar media

To prepare the Potato Dextrose Agar (PDA) medium, 200 g of peeled potatoes were boiled in water, and the extract was diluted to a total volume of 1 litre by adjusting with water. Then, 20 g of dextrose and 20 g of agar were dissolved in the solution and dispensed into 250 ml conical flasks, which were

Comment [MK19]: What type of potato? Give the cultivar name if possible.

plugged with non-absorbent cotton and sterilized at 121.6°C and 15 psi pressure for 20 minute in an autoclave.

Comment [MK20]: Get a referenced procedure for this preparation.

Preparation of different agro based waste extract media

Various The extracts of the agro-based waste listed in Table 1 in extract media were prepared by boiling each substrate composition separately as listed (Table 1) in 1 litre of water for 25-30 minutes. The resulting substrate extract was then strained through muslin cloth, and the volume was adjusted to 1 litre. Subsequently, 20 g of dextrose and 20 g of agar was dissolved made into the solution. The prepared substrate media was poured into 250 ml conical flasks, plugged with non-absorbent cotton, and sterilized at 121.6°C and 15 psi pressure for 20 minutes.

Comment [MK21]: It will be fine if the procedure used for this can also be referenced.

Comment [MK22]: How was this done and what was it meant for? Please, provide the detail.

Effect of temperature on growth of *Lentinula edodes*

For the present study, The prepared 17 different types of media were prepared and poured in duplicate at 20 ml per sterilized plate into respectively held at two different temperatures of, 20°C and 24°C on the sterilized Petri plates, with 20 ml per plate. This study aimed to evaluate the effect of mycelial growth on the *Lentinula edodes*. The Petri plates were allowed to solidify in aseptic conditions and each inoculated with 7mm diameter mycelium discs in at the center of the Petri plate containing media. Each treatment inoculation was done with in three replicates along with uninoculated plates as control, plate at two temperatures. The mycelial growths of *Lentinula edodes* was were observed measured in mm and recorded on the 7 and 14 DAI.

Comment [MK23]: Delete

Comment [MK24]: ?????

Table 1. List of different substrate compositions of agro waste extracts media used for in vitro evaluation cultivation of *Lentinula edodes* under in vitro conditions

S.No.	Substrate composition	Ratio and composition
S1	Sorghum Grains +Sawdust	(1:4) 40g +160g
S2	Wheat Grains+ Sawdust	(1:4) 40g+160 g
S3	Wheat Bran + Sawdust	(1:4) 40g+160 g
S4	Rice Bran + Paddy Husk and Sawdust	(1:1:3)40g+40g +120g
S5	Paddy Husk + Sawdust	(1:4) 40g+160 g
S6	Ashoka Woodchips + Paddy Husk + Wheat Bran	(3:1:1) 120g+40g+40g
S7	Ashoka Wood Chips +Paddy Husk + Rice Bran	(3:1:1) 120g+40g+40g
S8	Sawdust + Paddy Straw	(4:1) 160g+40 g
S9	Eucalyptus Bark + Sawdust	(4:1) 160g+40 g
S10	Eucalyptus Bark + Paddy Husk	(4:1) 160g+40 g
S11	Eucalyptus Bark + Rice Bran	(4:1) 160g+40 g
S12	Eucalyptus Bark + Wheat Grains + Sorghum Grains	(1:1:1) 66g+66g+66g
S13	Rice Bran +Sawdust	(1:4) 40g+160 g

S14	Only Sawdust	200g
S15	Only Sorghum Grains	200g
S16	Only Ashoka Wood Chips	200g
S17	PDA	200g

Statistical analysis

In this research, an experiment was carried out using a Factorial completely randomized design (FCRD) with three replications of each treatment by following the standard statistical procedure suggested by Gomez and Gomez (1984). The data obtained from the experiment were analysed using the analysis of variance (ANOVA) technique employed using SPSS software.

Comment [MK25]: 1. Please, re-cast this.
2.

Results and discussion

Lentinula edodes mycelial growth on different agro-based waste extract media at 20° C

Under in vitro conditions, various agro-based waste extracts were evaluated as media for the growth of *Lentinula edodes*. The mycelial growth ranged from 39.59mm to 64.40mm on different days after inoculation (DAI) recorded significant results. Maximum growth was recorded at 14 DAI (64.40mm) and lowest at 7 DAI (39.59mm). The impact of different substrates on mycelial growth was found to be significant. On S17 (82.67mm) media maximum growth was recorded while minimum on S8 (28.50mm) medium. Significant interaction effects of DAI and substrates evaluated were observed on the mycelial growth of shiitake mushroom. The maximum interaction effect was recorded on S17 (90mm), and S12 (90mm) media on 14 DAI, the minimum was recorded on S8 (22mm), and S14 (23.67mm) media at 7 DAI was on par with the minimum growth (Table 2 and Fig 1. Fig 2). Similar findings were reported earlier by Gbolagade *et al.* (2006) that PDA was the most suitable medium for culturing *Lentinula edodes* mycelium. Verma (2021) and his co-workers reported that the maximum mycelial growth was recorded on the PDA media. Iqbal (2009) and his associates reported similar findings, maximum mycelial growth of *L. edodes* was on the PDA media. Similarly, Shanmugarajet *al.* (2024) found that PDA media was best for the maximum growth of *Lentinula edodes*. Similar observations were also reported by Andrade *et al.* (2008). The mycelial growth of *Lentinula edodes* was assessed using various sawdust extracts from eucalyptus species. The results showed that the fastest growth was recorded when eucalyptus sawdust extract was used as growing media.

Comment [MK26]: What are the days?

Comment [MK27]: DAI means days after inoculation; please, indicate as such in the abstract.

Comment [MK28]: Construction of grammar and punctuations need to be thoroughly checked.

Comment [MK29]: Please, discuss your results as you relate to similar research works.

Table 2. Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 20°C temperature

S.No.	TREATMENTS	Mycelial growth(mm) *At 20°C		
		7 DAI	14 DAI	MEAN
S1	Sorghum Grains +Sawdust	34.00	67.75	50.88
S2	Wheat Grains+ Sawdust	26.33	58.00	42.17
S3	Wheat Bran + Sawdust	30.33	58.75	44.54
S4	Rice Bran + Paddy Husk and Sawdust	26.00	56.75	41.38
S5	Paddy Husk + Sawdust	25.33	55.75	40.54
S6	Ashoka Woodchips + Paddy Husk + Wheat Bran	49.33	66.50	57.92
S7	Ashoka Wood Chips +Paddy Husk + Rice Bran	31.33	56.50	43.92
S8	Sawdust + Paddy Straw	22.00	35.00	28.50
S9	Eucalyptus Bark + Sawdust	43.67	65.25	54.46
S10	Eucalyptus Bark + Paddy Husk	56.00	82.00	69.00
S11	Eucalyptus Bark + Rice Bran	56.33	83.25	69.79
S12	Eucalyptus Bark + Wheat Grains + Sorghum Grains	64.00	90.00	77.00
S13	Rice Bran +Sawdust	24.00	51.25	37.63
S14	Only Sawdust	23.67	52.00	37.84
S15	Only Sorghum Grains	57.67	80.00	68.84
S16	Only Ashoka Wood Chips	27.67	46.00	36.84
S17	PDA	75.33	90.00	82.67
	MEAN	39.59	64.40	
	CV	2.24		
		CD @5%	SE(m)	
	DAI (A)	0.466	0.165	
	Substrate (B)	1.359	0.482	
	Interaction (A X B)	1.922	0.681	

Note: * Average of 3 replications




Fig.1 Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 20°C temperature

Comment [MK30]: The horizontal axis in the chart should be titled.



Fig.2. Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 20°C temperature at 14 DAI

Note: SDPS-Sawdust and Paddy Straw, EBPH-EucalyptusBark and Paddy Husk, EBRB-Eucalyptus Bark and Rice Bran, EBWGSG- Eucalyptus Bark and Wheat Grains and Sorghum Grains, PDA-Potato Dextrose Agar

Comment [MK31]: Keys

Comment [MK32]: Indicate S8 to S17 inside Fig. 2

***Lentinula edodes* mycelial growth at 24° C temperature**

Growth of *Lentinula edodes* mycelium was significant among two DAI evaluated and ranged between 55.86mm to 80.45mm. Maximum growth was recorded on 14 DAI (80.88mm) and minimum on 7 DAI (55.86mm). The impact of different substrates on mycelial growth was significant, varying from 90mm to 36mm. The maximum growth was recorded on S17 (90mm) media while the lowest was recorded on S8 (36mm) medium. Additionally, the interaction effect of two DAI and among 17 various substrates, mycelial growth of *L. edodes* was significant at a temperature of 24°C. The maximum interaction effect was recorded 90mm at 7 DAI and 14 DAI on S17 and S12 media. On S11, and S10 media also recorded maximum growth (90mm) at 14 DAI. While the minimum growth was recorded on S8 (30.33mm) medium at 7 DAI and 14 DAI (Table 3 and Fig 3, Fig 4.). The experimental results

Comment [MK33]: At what probability?

Comment [MK34]: Lower to higher i.e. 36 – 90mm

revealed a significant effect of temperature on the mycelial growth of shiitake mushroom under *invitro* conditions. These findings are in accordance with the results of Verma, I.K. and Singh, P. (2021) who reported that PDA served as the most conducive medium for the growth of *Lentinula edodes* mycelium and provided evidence supporting the suitability of PDA as an effective media. The study revealed that the optimal temperature for the *L. edodes* growth was 24°C, where maximum mycelial growth was recorded within 7 DAI when compared to other agro-based waste extract media. Similarly, Kumar and his co-workers (2019) reported that the maximum growth of mycelium was recorded at 24°C.

Comment [MK35]: Mention the evidence

Comment [MK36]: Discuss your findings.

Comment [MK37]: *et al.*




Fig 3. Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 24°C temperature

Comment [MK38]: The horizontal axis in the chart should be titled.




Fig 4. Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 24°C temperature at 14 DAI

Note: SDPS-Sawdust and Paddy Straw, AWPHWB-Ashoka Wood chips and Paddy Husk, Wheat Bran, EBPH-Eucalyptus Bark and Paddy husk, EBRB-Eucalyptus Bark and Rice Bran, EBWGSG-Eucalyptus Bark and Wheat Grains and Sorghum Grains, PDA- Potato Dextrose Agar

Comment [MK39]: Indicate S8 to S17 inside Fig. 4

Table 3. Mycelial growth of *Lentinula edodes* on different agro-based waste extract media at 24°C temperature

S.No	TREATMENTS	Mycelial growth(mm)* At 24°C		
		7 DAI	14 DAI	MEAN
S1	Sorghum Grains +Sawdust	46.33	89.60	68.167
S2	Wheat Grains+ Sawdust	43.67	84.90	64.333
S3	Wheat Bran + Sawdust	42.94	83.67	62.333
S4	Rice Bran + Paddy Husk and Sawdust	40.60	81.00	60.867
S5	Paddy Husk + Sawdust	42.67	83.33	62.167
S6	Ashoka Woodchips + Paddy Husk + Wheat Bran	64.00	89.00	76.50
S7	Ashoka Wood Chips +Paddy Husk + Rice Bran	51.00	78.33	64.167
S8	Sawdust + Paddy Straw	30.33	42.97	36.00
S9	Eucalyptus Bark + Sawdust	58.67	79.83	69.00
S10	Eucalyptus Bark + Paddy Husk	68.00	90.00	79.50
S11	Eucalyptus Bark + Rice Bran	75.00	90.00	83.00
S12	Eucalyptus Bark + Wheat Grains + Sorghum Grains	90.00	90.00	90.00
S13	Rice Bran +Sawdust	43.00	88.33	65.167
S14	Only Sawdust	39.00	82.37	60.66
S15	Only Sorghum Grains	73.00	87.00	80.00
S16	Only Ashoka Wood Chips	40.37	62.00	51.33
S17	PDA	90.00	90.00	90.00
	MEAN	55.86	80.88	
	CV	1.58		
		CD @5%	SE(m)	
	DAI (A)	0.427	0.151	
	Substrate (B)	1.244	0.441	
	Interaction (A X B)	1.76	0.624	

Note: * Average of 3 replications

Table 4. Principal Component Analysis of *Lentinula edodes* on different substrates at various temperatures.

Component		Eigenvalues	Proportion	Cumulative (%)
Mycelial growth of <i>L.edodes</i> at 24°C	7DAI	5.148	0.644	0.644
	14 DAI	1.436	0.179	0.823
Mycelial growth of <i>L.edodes</i> at 20°C	7DAI	0.691	0.086	0.909
	14 DAI	0.352	0.044	0.953
Colony characters of <i>L.edodes</i> mycelium at 24°C	Colony colour	0.213	0.027	0.980
	Thickness of mycelium	0.092	0.011	0.991
Colony characters <i>L.edodes</i> mycelium at 20°C	Colony colour	0.062	0.008	0.999
	Thickness of mycelium	0.062	0.001	1.000




Fig 5. Screen plot for growth of *Lentinula edodes* by using Principal Component Analysis

According to the **Principal Component Analysis**, Eigen value ≥ 1 i.e., Mycelial growth of *L. edodes* 7 DAI and 14 DAI at 24°C temperature components was considered for the existence of

Comment [MK40]: 1.If Eigen values have unit of measurement, please, indicate.
2. Add more meaning to the component number inside the chart. component number of what?

Comment [MK41]: Please, include the procedure for this.

influences of mycelial growth of *L. edodes* among 17 substrates extract media as illustrated in (Table 4 and Fig 5). The results from the Principal Component Analysis of recorded data of mycelial growth was shown in the form of Eigen values, which shows that the highest Eigen value was recorded for mycelial growth of *L. edodes* at 7 DAI and 14 DAI indicates significant influence of substrates.

Conclusion

This laboratory study aimed to assess the growth performance of the Shiitake mushroom (*Lentinula edodes*) using various agro-based waste extracts as growth media components. Among the different agro-based wastes, extracts of eucalyptus bark supplemented with wheat grains, sorghum grains, rice bran, and paddy husk and PDA recorded the maximum mycelial growth at temperature of 24°C. These findings suggest that *Lentinula edodes* has the potential to be cultivated using locally available residues for its spawn production. Tissue culture technique can be employed in further research is needed for commercial production of shiitake mushroom spawn.

References

- Kumar, V., Mishra, S.K and Kaur, M. 2019. Effect of different media, temperature and pH on radial mycelial growth of *Lentinula edodes* strain Le-17-04. *Journal of Pharmacognosy and Phytochemistry*. 8(1):345-348.
- Chang, S.T and S.P. Wasser. 2017. The cultivation and Environmental impact of mushrooms. Printed from the Oxford Research Encyclopaedia, Environmental Science (c) Oxford University Press, USA:43.
- Kobayashi, T., Oguro, M., Akiba, M., Taki, H., Kitajima, H and Ishihara, H. 2020. Mushroom yield of cultivated shiitake (*Lentinula edodes*) and fungal communities in logs. *Journal of Forest Research*. 25(4): 269-275.
- Annepu, Sudheer Kumar & Sharma, Ved Parkash & Kumar, Satish & Barh, Anupam. (2019). Cultivation Techniques of Shiitake (A Medicinal Mushroom with Culinary Delight).
- Gbolagade, J.S., Fasidi, I.O., Ajayi, E.J. and Sobowale, A.A. 2006. Effect of physico-chemical factors and semi-synthetic media on vegetative growth of *Lentinus subnudus* (Berk.), an edible mushroom from Nigeria. *Food chemistry*. 99(4): 742-747.
- Shanmugaraj, C., Saranraj, K. and Biswas, M.K., Optimizing growth conditions for shiitake mushroom cultivation in Birbhum, West Bengal: A study on media, temperature, and pH variations.
- Paswal, S., Kakraliya, S.S. and Fogawat, V. 2024. Effect of Different Substrates on Nutritional Composition of Shiitake Mushroom (*Lentinula edodes*). *Archives of Current Research International*. 24(4):156-161.
- Andrade, M.C.N.D., Silva, J.H.D., Minhoni, M.T.D.A. and Zied, D.C. 2008. Mycelial growth of two *Lentinula edodes* strains in culture media prepared with sawdust extracts from seven eucalyptus species and three eucalyptus clones. *Acta Scientiarum. Agronomy*, 30: 333-337.
- Iqbal, S.M., Arif, Z., Zahid, N.Y. and Abbasi, N.A. 2009. Effect of different culture medium and pH on the mycelial growth of shiitake mushroom. *MYCOPATH*, 13(1).

Comment [MK42]: Inconsistent referencing styles

Comment [MK43]: Not cited

Comment [MK44]: Old ref.

Comment [MK45]: Incomplete referencing

Comment [MK46]: Old ref.

Comment [MK47]: Old ref

Verma, I.K. and Singh, P. 2021. In vitro evaluation for optimum growth conditions of shiitake mushroom (*Lentinula edodes*). *Mushroom Research*. 29(2).

Gomez, K.A. and Gomez, A.A., 1984. *Statistical procedures for agricultural research*. John Wiley & sons.

Comment [MK48]: Old ref

UNDER PEER REVIEW