

Effect of integrated nutrient management on growth Characteristics of the soil properties of rice crops

Comment [a1]: Soil properties not added. Only initial soil properties added. Add after harvest soil properties or delete the word from title.

Comment [a2]: Effect of INM on growth parameters of rice grown on alluvial calcareous soil

Abstract

The field experiment was conducted at Main Experimental Station of Narendra Deva University of Agriculture and Technology, Kumarganj, Faizabad (U.P.) during the *Kharif*, 2014 to study the effect of integrated nutrient management on soil properties, growth Characteristics of rice (*Oryza sativa*). The experiment comprised of ten treatments viz. T₁ (control), T₂ (100% RDF) T₃ (125% RDF), T₄ (100% RDF + FYM @ 5 t ha⁻¹), T₅ (100% RDF + Brown manuring), T₆ (100% RDF + BGA @ 10 kg ha⁻¹), T₇ (100% RDF + PSB), T₈ (Farmers practice (N₁₀₀:P₄₀:K₀), T₉ (100% RDF + FYM @ 5 t ha⁻¹ + PSB + *Azotobacter* + BGA) and T₁₀ (100% RDF + FYM @ 5 t ha⁻¹ + PSB + *Azotobacter* + Brown manuring) in Randomized Block Design with three replications with rice variety NDR-97 was taken. The experimental soil was silty loam in texture having the maximum values of pH (7.88), EC dSm^{-1} (0.32 dSm^{-1}), organic carbon (0.34 %).

Keys words: RDF, FYM, PSB , *Azotobacter*, BGA

Introduction

Rice (*Oryza sativa* L.) is the most important staple food crop in the world. It is a rich source of energy and contains a reasonable amount of protein (6-10%), carbohydrate (70-80%), minerals (1.2-2.0%), and vitamins such as riboflavin, thiamine, niacin, and vitamin E. Over 90% of the world's rice is grown and consumed in Asia, often referred to as the "rice bowl of the world," where 16% of the earth's population and two-thirds of the world's poor reside.

Formatted: Font: (Default) Times New Roman

Formatted: Justified

Rice is the primary food source for people living in the humid tropics and subtropics, and it is grown worldwide.

~~Rice is the principal food for people living in the humid tropics and subtropics, and it is cultivated in tropical and subtropical regions globally.~~ The global requirement for rice is projected to be about 880 million tonnes by 2050. According to the FAO (2014), worldwide production of rice was 719.74 million metric tonnes with an acreage of 160.6 million hectares. In India, rice is cultivated over approximately 43.9 million hectares, with an annual production of about 88.02 million tonnes. India contributes around 21% of global rice production from its 28% share of the world's rice area. Rice is a vital food for the majority of the world's population. In Uttar Pradesh, the area under rice cultivation is about 5.964 million hectares, with a production of 2.50 million tonnes and a productivity of 2358 kg per hectare (FAO 2014). The decline or

Comment [a3]: Old statistics ..update with new one.

stagnation in rice yield has been attributed to nutrient depletion and reduced use of organic matter (John et al., 2001). In the current scenario, where food and nutritional security are under threat, it is imperative to enhance the soil-plant-atmosphere system as a whole rather than focusing solely on crop nutrition. The combined use of organic and inorganic nutrient sources, along with biofertilizers, has proven to be more effective than using each component separately (Palanippan and Annadurai, 2007). India's population is expected to reach 1.4 billion by the year 2025, further emphasizing the need for sustainable agricultural practices.

Comment [a4]: Add latest reviews

Material and Methods:

Geographical situation of experimental site:

The experiment was conducted at the Main Experimental Station of Narendra Nagar (Kumarganj), Faizabad (U.P.) during the *Kharif* season of 2014. ~~This chapter describes the climatic and edaphic conditions, materials used, and methods employed during the investigation.~~ The ~~m~~Main ~~e~~Experimental ~~s~~Station ~~was of Narendra Deva University of Agriculture and Technology,~~ located in Narendra Nagar (Kumarganj), Faizabad (U.P.), is situated on the Faizabad-Raibarely road, approximately 42 km from Faizabad. The experimental site falls under the subtropical zone of the Indo-Gangetic plains, characterized by alluvial calcareous soil. It is located between 24.440-26.560 degrees North latitude and 82.120-83.980 degrees East longitude, with an elevation of about 113 meters above mean sea level. The experimental field was well-leveled and had good irrigation and drainage facilities. The mean annual rainfall in this region is 1021.8 mm, with about 90% of it occurring during the monsoon season.

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Soil characteristics:

The soil of the experimental site is part of the Indo-Gangetic plain, classified as the Eastern Plain Zone (EPZ) in the alluvial tract of Uttar Pradesh. The properties of the soil are detailed in Table 1. Soil analysis data indicate that the soil is low in available nitrogen, ~~and has~~ medium levels of phosphorus and potassium.

Formatted: Normal (Web)

Table 1 Initial Analysis of Experimental Soil

S. No.	Properties	Values	Method Employed
A. Physical properties			
1.	Bulk density	1.36	Core method (undisturbed sample) Black, 1965
2.	Texture		
i)	Sand (%)	24.15	Hydrometer method (Bouyoucos, 1936)
ii)	Silt (%)	56.68	
iii)	Clay (%)	19.17	
iv)	Textural class	Silt loam	Triangular method
B. Chemical properties			
1.	pH (1:2.5)	7.90	Glass electrode pH meter (Jackson, 1973)
2.	EC (dSm ⁻¹ at 25°C)	0.38	Conductivity meter (Jackson, 1973)
3.	Organic carbon (%)	0.30	Walkley & Black's rapid titration method (1934)
4.	Available N (kg ha ⁻¹)	130.96	Alkaline permanganate method (Subbiah and Asija, 1956)
5.	Available P (kg ha ⁻¹)	16.90	Olsen's method (Olsen et al., 1954)
6.	Available K (kg ha ⁻¹)	220.75	Flame photometer (Jackson, 1973)
7.	Available Zn (ppm)	0.49	Extraction with DTPA solution (Lindsay & Norvell, 1978)

Comment [a5]: Add in reference

Formatted: Superscript

Comment [a6]: Add in reference

Formatted: Superscript

Comment [a7]: Add in reference

Formatted: Superscript

Comment [a8]: Add in reference

Formatted: Superscript

Formatted Table

Comment [a9]: Add in reference

Treatments details: T1: Control, T2: 100% RDF, T3: 125% RDF, T4: 100% RDF + FYM @5 t ha⁻¹, T5: 100% RDF + Brown manuring, T6: 100% RDF + BGA @10 kg ha⁻¹, T7: 100% RDF + PSB, T8: Farmers practice (N100:P40), T9: 100% RDF + FYM @5 t ha⁻¹ + PSB + Azotobacter + BGA, T10: 100% RDF + FYM @5 t ha⁻¹ + PSB + Azotobacter + Brown manuring.

Comment [a10]: Per ha should be super script all over the manuscript

Layout Details:

The experiment was designed in Randomized Block Design (RBD), with ten treatments (i.e., T1: Control, T2: 100% RDF, T3: 125% RDF, T4: 100% RDF + FYM @5 t ha⁻¹, T5: 100% RDF + Brown manuring, T6: 100% RDF + BGA @10 kg ha⁻¹, T7: 100% RDF + PSB, T8: Farmers practice (N100:P40), T9: 100% RDF + FYM @5 t ha⁻¹ + PSB + Azotobacter + BGA, T10: 100% RDF + FYM @5 t ha⁻¹ + PSB + Azotobacter +

Brownmanuring and three replications. Each plot has a net plot size of $3.2 \times 4 = 12.8 \text{ m}^2$ and a gross plot size of $4 \times 5 = 20 \text{ m}^2$. The spacing is $20 \text{ cm} \times 10 \text{ cm}$. The variety used is NDR-97.

Design: Randomized Block Design (RBD), Number of replications: 3, Number of treatments: 10, Total number of plots: 30, Net plot size: $3.2 \times 4 = 12.8 \text{ m}^2$, Gross plot size: $4 \times 5 = 20 \text{ m}^2$, Spacing: $20 \text{ cm} \times 10 \text{ cm}$, Variety: NDR-97

Comment [a11]: Per ha should be super script

Result and Discussion

The data on plant height as influenced by integrated nutrient management are presented in Table 2. Significant differences in plant height were observed due to the application of various organic, inorganic, and biofertilizer treatments. Treatments included different levels of RDF: 100% RDF alone, 125% RDF alone, 100% RDF combined with FYM @ 5 t ha⁻¹, Brown manuring, BGA @ 10 kg ha⁻¹, PSB, and combinations of 100% RDF with FYM @ 5 t ha⁻¹ + PSB + Azotobacter + Brown manuring or BGA @ 10 kg ha⁻¹. The highest plant height was recorded in the treatment with 100% RDF + FYM @ 5 t ha⁻¹ + PSB + Azotobacter + Brown manuring. The increase in plant height can be attributed to better nourishment, which enhances beneficial effects such as an accelerated rate of photosynthesis, assimilation, cell division, and vegetative growth. These results align with the findings of Lin and Lin (1985), Priyanka *et al.* (2013), and Sujatha *et al.* (2014).

Number of tillers m⁻²:

The number of tillers per square meter is a crucial growth attribute influenced by organic, inorganic, and biofertilizers. The maximum number of tillers was recorded with treatment T10, which included 100% RDF + FYM @ 5 t ha⁻¹ + PSB + Azotobacter + Brown manuring substituted by organic application. The increased number of tillers per square meter can be explained by due to the balanced nutrient supply that encouraged vegetative growth through the combined use of inorganic, organic, and biofertilizer sources. Organic manures and biofertilizers improved soil fertility, enhancing nutrient availability and, consequently, the number of tillers per square meter. This improved nutrient availability ultimately leads to higher crop yields. These results are consistent with the findings of Azad and Leiria (2001), Satsangi *et al.* (2002), Jagdish and Yadav (2008), Lin and Lin (1985), and Dutt and Chauhan (2010).

Comment [a12]: All the parameters no. of tillers dry matter production and height are T10 and T 9 treatments are o par

Dry Matter Production

The increase in dry matter production was observed at different growth stages due to the application of organic, inorganic, and biofertilizer treatments. Various levels of RDF were applied: either 100% RDF alone, 125% RDF alone, or 100% RDF in combination with FYM @

5 t ha⁻¹, Brown manuring, BGA @ 10 kg ha⁻¹, PSB, or combinations of 100% RDF with FYM @ 5 t ha⁻¹ + PSB + Azotobacter + Brown manuring or BGA @ 10 kg ha⁻¹. The maximum dry matter production was recorded in the treatment with 100% RDF + FYM @ 5 t ha⁻¹ + PSB + Azotobacter + Brown manuring, followed by the treatment with 100% RDF + FYM @ 5 t ha⁻¹ + PSB + Azotobacter + BGA.

The increase in dry matter production with these treatments can be explained by the balanced nutrient supply, which enhances better translocation of photosynthates from source to sink. The significant improvement in dry matter production is attributed to the various levels of RDF and their combined application. Treatments with Brown manuring showed relatively higher dry matter production compared to those with BGA. While the use of inorganic fertilizers alone also contributed to dry matter production, the extent was not as significant as when combined with organic and biofertilizers.

The increased dry matter production at different stages under various treatments was due to the increased availability of nutrients in adequate amounts during different growth stages of rice. These results are corroborated by the findings of Begum et al. (2009), Hasan ~~et al.~~ (2012), Priyanka et al. (2013), and Mhalevilie et al. (2013), who also reported significant increases in dry matter production of rice with RDF and its combination with BGA, PSB, or Brown manuring. The variation in dry matter production among different treatments was due to differences in nutrient availability in the soil at respective growth stages.

Comment [a13]: these two treatments were one par no significant difference. Change the result.

Table-2. Effect of integrated nutrient management on plant height (cm) and Number of tillersm⁻² at various growth stages of rice crop

Treatments	Plant height (cm)				Number of tillers m ⁻²			
	30 DAT	60 DAT	90 DAT	At harvest	30 DAT	60 DAT	90 DAT	At harvest
T ₁ - Control	30.21	55.07	68.96	70.85	128.44	290.22	299.48	298.48
T ₂ - 100% RDF	33.52	68.97	79.74	81.28	136.39	338.01	345.68	346.12
T ₃ - 125% RDF	33.96	70.35	83.02	82.72	140.88	349.13	359.78	360.60
T ₄ - 100% RDF + FYM @5 t ha ⁻¹	33.74	69.57	82.98	84.10	139.20	344.94	362.78	360.00
T ₅ - 100% RDF+ Brown manuring	33.50	69.12	82.22	83.52	138.78	343.90	356.56	357.12
T ₆ - 100% RDF + BGA@10 kg ha ⁻¹	33.57	68.25	81.10	82.65	138.10	342.21	358.78	359.90
T ₇ - 100% RDF +PSB	33.46	67.92	80.02	81.99	136.69	338.68	346.01	345.68
T ₈ - Farmers practice(N ₁₀₀ :P ₄₀ :K ₀)	32.06	62.10	77.20	79.00	132.55	329.11	333.17	333.17
T ₉ - 100% RDF + FYM @5 t ha ⁻¹ +PSB+Azotobacter+BGA	35.16	73.20	84.98	85.11	153.00	366.11	375.78	376.12
T ₁₀ - 100%RDF+FYM @5 t ha ⁻¹ +PSB+Azotobacter+Brown manuring	35.68	74.65	85.08	86.95	154.70	368.06	376.12	376.28
SEm+	2.06	2.25	2.56	2.96	1.64	3.29	3.96	4.51
CD (P= 0.05)	NS	6.66	7.76	8.08	4.90	9.86	12.10	13.76

Table-3. Effect of integrated nutrient management on dry matter production (g m^{-2}) at various growth stages of rice crop

Treatments	Dry matter production (g m^{-2})			
	30 DAT	60 DAT	90 DAT	At harvest
T ₁ - Control	48.97	77.53	119.78	135.84
T ₂ - 100% RDF	68.69	114.29	238.70	270.71
T ₃ - 125% RDF	74.94	129.70	260.44	295.37
T ₄ - 100% RDF + FYM @5 t ha ⁻¹	73.05	124.16	258.56	294.09
T ₅ - 100% RDF+ Brown manuring	70.62	122.52	256.49	292.90
T ₆ - 100% RDF + BGA@10 kg ha ⁻¹	72.96	123.42	257.20	292.78
T ₇ - 100% RDF +PSB	69.56	115.76	242.35	272.77
T ₈ - Farmers practice(N ₁₀₀ :P ₄₀ :K ₀)	58.12	93.78	141.19	196.98
T ₉ - 100% RDF + FYM @5 t ha ⁻¹ +PSB+Azotobacter+BGA	86.52	138.96	274.66	314.76
T ₁₀ - 100%RDF+FYM @5 t ha ⁻¹ +PSB+Azotobacter+Brown manuring	89.00	140.17	275.12	316.05
SEm±	2.10	3.04	4.17	5.89
CD (P= 0.05)	3.60	9.12	12.17	17.64

Conclusion:

The results obtained in this experiment, it can be conclusively stated that the application of FYM at 5 t/ha, along with PSB, Azotobacter, and brown manuring, proved to be the most effective approach for enhancing growth characteristics. Additionally, this combination significantly contributed to maintaining soil health, thereby promoting sustainable rice production.

Reference

- Azad, B. S., &Lehria, S. K. (2001). Yield maximization of rice through integrated nutrient management under irrigated conditions. *Annals of Agriculture research*.**22** (4): 471-475.
- Begum, Z. N. T., Mandal, R., & Islam, M. S. (2009). Effect of blue-green algae and Urea-N on growth and yield performance of traditional variety of rice. *Journal of Phytological Research*. **22**(2):211-214.
- Dutta, M. and Chauhan, B.S. (2010). Effect of nutrient management practice on the performance of upland rice in a newly developed terraced land. *Indian Agriculture*. **54** (2): 13-21.
- FAO (2014) <http://www.fao.org/worldfoodsituation/csdb/en/>

Comment [a14]: BGA treatment T 9 also on par with T10

- Hasan, M. A. (2012). Investigation on the Nitrogen fixing cyanobacteria (BGA) in rice fields of North-West region of Bangladesh. *Journal of Environmental Science and Natural Resources*. **5** (2): 185-192.
- Jagdish, K. and Yadav, M.P. (2008). Effect of conjunctive use of organic inorganic and bio-fertilizer on growth and yield attributes, yield and nutrient uptake in hybrid rice (*Oryza sativa* L.). *Research on crop*. **9** (3): 511-113.
- John, P.S., George, M. and Jacob, R., Zool. (2001). Nutrient mining in agro climatic zones of Kerala, *Fertilizer news*. **46**:45-52 and 55-57.
- Lin, J.L., and Lin, T.L. (1985). Tiller number and leaf area index rice community as influenced by planting density and nitrogen fertilizer. *J. Agril. Assoc. of China*. **12** (9): 14-34.
- Mhalevilie, T., Sarkar, N.C., and Atou, K. (2013). Effect of biofertilizer on the productivity of terraced upland rice (*Oryza sativa* L.) *International journal of bio-resource and stress management*. **4** (3): 400-403.
- Palaniappan, S.P. and Annadurai, K. (2007). Organic farming: theory and practices.
- Priyanka G., Sharma G.D., Rana R. and Lal B. (2013). Effect of integrated nutrient management and spacing on growth parameters, nutrient content and productivity of rice under system of rice intensification. *International journal of research in bio sciences*. **2** (3): 53-59.
- Satsangi, G. P., Bhavnagar, V. P. and Srivastava, J. N. (2002). Effect of BGA/Urea as nitrogen application on growth. *Journal of Living World*. **9** (1):5-13.
- Sujatha, K., Mosha, G., Subbaiah P. and Prasuna.P.(2014). Residual soil fertility and productivity of rice (*Oryza sativa* L.) as influenced by different organic sources of nitrogen. *International journal of plant, animal and environmental sciences*. **5** (2): 266-289.