

Original Research Article

STUDIES ON GENETIC VARIABILITY AND MORPHOLOGICAL VARIATIONS OF GREENGRAM [*Vigna radiata* (L.) WILCZEK] GERMPLASM

ABSTRACT

Greengram [*Vigna radiata* (L.) Wilczek] belongs to the family of Leguminosae (Fabaceae) and it is an important annual legume crop widely cultivated in semi-arid tropics. In the present investigation, a total of 20 lines of greengram including one check variety were evaluated during Zaid, 2023 in Randomized Block Design with three replications for genetic variability, heritability, and genetic advance by using 13 quantitative traits and to study morphological variation for these 13 quantitative characters by metroglyph and index score method. Analysis of variance for various quantitative characters revealed that the mean sum of squares due to genotypes showed high significant differences for all characters under study at 1% level and 5% level of significance. High GCV, PCV, heritability, and genetic advance were recorded by number of pods per plant. Based on metroglyph analysis and scatter diagram, 20 genotypes were grouped into 5 complexes, and the maximum number of genotypes (15) were found in group I. The two most variable characters, number of pods per plant and plant height (cm) were selected for the X and Y axes, respectively. The germplasm lines VIRAT (31), CO-7 (31), LGG-407 (30), AMULYA (29) and IPM-2-3 (29) recorded high index scores and fell into different clusters do well for morphological variations for a greater number of traits. Thus, the use of these genotypes in future breeding programs for crossing to obtain maximum variability of good combinations is suggested.

Keywords: Genetic diversity, Heritability, Index score, Metroglyph, Morphological variation.

1. INTRODUCTION

“Pulses, often regarded as “Poor man’s meat” are considered the earliest domestic plants and are the richest source of protein, starch, minerals, and vitamins and are a major source of human food second only to cereals (Poaceae). Greengram [*Vigna radiata* (L.) Wilczek], commonly called mung bean, is also known as “Golden gram”, is one of the most ancient and extensively grown leguminous crops of India ($2n=22$ and genome size of 494 to 579 Mb). It is the third most important pulse crop after chickpea and pigeonpea, cultivated throughout India for its multipurpose uses as a

vegetable, pulse, fodder, and green manure crop. Its seed is more palatable, nutritive, digestible, and nonflatulent than other pulses grown in the world”(Kartik *et al.*, 2020).“Greengram is primarily a native of India and Central Asia may be a secondary center. There are around 7.3 million hectares of mungbean cultivated worldwide, with an average production of 721 kg/ha and 30% of the 5.3 million tonnes of production produced globally is split between India and Myanmar. China, Indonesia, Thailand, Kenya, Tanzania, Nepal, Sri Lanka, Korea, and Pakistan are other significant producers”(Nair *et al.*, 2020). In India, the mungbean alone accounts for 10% of production (17 lakh tonnes) with productivity of 500 kg/ha and 16% of area (36 lakh hectares) of all pulses. The dominant contributors to mungbean cultivation in terms of area and production are Rajasthan, Madhya Pradesh, Maharashtra, Karnataka, Odisha, Bihar, Tamil Nadu, Gujarat, Andhra Pradesh, and Telangana as stated in the **Annual Report (2022-23) by AICRPR on Kharif pulses.**

“In particular, due to poor management and low soil fertility, greengram produces low seed yield and poor growth. However, the yield potential of this crop continues to be low and for this reason, it is facing severe competition from cereal crops, more particularly in recent years with the availability of high-yielding cereal varieties. The presently grown cultivars of mungbean are mostly old land races which are the products of direct and indirect natural and human selections. Due to this unique position in our agricultural system, the crop yield is very low at present. Recently, with the awareness of this reason, intensive work has been initiated to conserve and create diversity and thereby improve the crop” (Nair *et al.*, 2020).

The agriculture practices and successive breeding have narrowed down the genetic base of cultivated greengram. The lack of genetic variability is an important factor for the limited progress achieved in increasing the productivity of grain legumes including greengram. Nevertheless, genetic improvement of morphological traits, yield, and yield components are the most difficult to achieve due to the complex nature of the inheritance and the numerous environmental factors that influence yield and its genetic basis of variation in greengram. Further, to improve the productivity, information about the nature and magnitude of genetic divergence would help selection of diverse parents, which upon hybridization might lead to effective gene recombination. The present study was undertaken to select the divergent parent for future hybridization programme.

“The success of any crop Improvement program largely depends on the selection of diverse parents as breeding material. Genetic diversity has been considered as an important factor which is also an essential prerequisite for hybridization programs or obtaining high-yielding progenies. The inclusion of diverse parents in the hybridization program will serve the purpose of combining desirable genes to obtain superior recombinants”(Johnson *et al.*, 1955). Estimation of genetic variability used in conjunction with heritability and genetic advance gives an idea of the possible improvement obtained through selection. High heritable estimates together with genetic advance are more valid for selection than heritability estimates alone. In view of that, it is therefore appropriate to assess the proportion of genetic and non-genetic variability observed in the traits studied. Exploitation of genetic variability in available germplasm is taken as a key point for making further genetic improvements in economically important traits as well as yield.

“This diversity not only results in inducing genetic variation but also provides scope for new recombination of genes within the existing gene pool. Hence, studies on genetic diversity are of considerable importance to classify the available genotypes into discrete classes so that the parents belonging to diverse groups can be selected. In addition to aiding in the selection of divergent parents for hybridization, metroglyph analysis is a favorable tool for use in plant breeding as it measures the degree of divergence at the genotypic level and determines the relative contribution of each component character to the total divergence”(Anderson, 1957). The presence of genetic variability is of utmost importance for any breeding program and due to this reason, plant breeders have emphasized the evaluation of germplasm for the improvement of crop yield as well as for utilization in further breeding programs. Knowledge of genetic divergence coupled with genetic parameters and genetic gain obtained by selection is an important prerequisite for a systematic breeding program to improve the yield potential of genotypes.

1.1 OBJECTIVES

1. Determine the extent of genetic variation for morphological traits in Greengram.
2. Identify the divergence pattern of greengram germplasm to develop selection-criteria for future hybridization programs.
3. Classify genotypes by metroglyph and index score analysis for morphological characters in Greengram.

2. MATERIALS AND METHODS

The present investigation was carried out at the Research Farm, Department of Genetics and Plant Breeding, SHUATS, Prayagraj, during the *Zaid* season 2023. The experimental material comprises 20 genotypes along with one check variety. A randomized complete block design was used for the experiment in three replications with row-to-row spacing of 30 cm and plant-to-plant spacing of 10 cm and a plot size of 1m x 1m. For this green gram crop, recommended agronomical and plant protection practices were followed.

To select the best yield-giving genotype in the agro-climatic conditions of Prayagraj region, observation was recorded by selecting five plants randomly from each genotype in each replication for various quantitative traits like plant height, days to 50% flowering, days to 50% pod setting, days to maturity, plant height (cm), number of primary branches, number of clusters per plant, number of pods per plant, pod length (cm), number of seeds per pod, seed index (g), biological yield (g), harvest index (%) and seed yield per plant (g). Thereafter, the phenotypic quantitative traits were compared with high-yielding check varieties for varietal selection. To determine the outcomes for the mentioned morphological characters in different crop species, biometrical and statistical analyses using metroglyph and index score method advocated by Anderson (1957) has been used on the recorded data.

Analysis of Variance (ANOVA) was done by subjecting the data to the statistical method based on Randomized Block Design (RBD) described by **Panse and Sukhatme (1978)**. "Estimation of genetic variability parameters viz., range, phenotypic coefficient of variation, genotypic coefficient of variation, heritability (h^2), genetic advance and genetic advance as % of mean were calculated separately for each character. For metroglyph analysis, the first two characters having high variability on the basis of genotypic coefficient of variation are selected. The mean values of X character for each genotype are plotted on the graph against the mean values of Y character. A small circle by which the position of genotypes is represented on the graph is called glyph. Then the genotypes are divided into three groups and these groups are given an index score of 1, 2 and 3 respectively. Thus the whole graph is divided into 9 parts. Genotypes in each part of the graph are representing one variability group. Variation for each character is depicted by the length of ray which depends on the index score value of a genotype. Each character occupies a definite ray position. The variation is analyzed for various traits within the group and between the groups. Thus the maximum and minimum score that an individual genotype can get

is $3n$ and n respectively where n is the total number of characters studied”(Singh and Narayanan, 2000).

Table 1: List of Greengram genotypes used in present study

Sl. no.	Name of genotypes	Sl. no.	Name of genotypes
1	VBN-3	11	CO-7
2	SML-1668	12	MGG-295
3	SM-20-108	13	IPM-2-14
4	MH-421	14	SHAKTI
5	SM-20-103	15	LGG-450
6	TM-96-2	16	IPM-2-3
7	LGG-460	17	AMULYA
8	SHIKHA	18	PUSA BAISAKHI
9	CO-8	19	LGG-407
10	PUSA-105	20	VIRAT (CHECK)

3. RESULT & DISCUSSION

3.1 ANALYSIS OF VARIANCE

Significant differences among the genotypes for all the traits have been observed in the analysis of variance. Analysis of variance for various quantitative characters revealed that the mean sum of squares due to genotypes showed high significant differences for all characters under study at 1% level and 5% level of significance. This reveals that the quantitative characters studied seem to have considerable genetic variability. Thus, there is an ample scope for improvement of different quantitative traits through selection. Based on the mean performance, seed yield per plant was observed high for CO-7 (15.57), AMULYA (14.453), and IPM-2-14 (14.15).

3.2 PHENOTYPIC AND GENOTYPIC COEFFICIENT OF VARIATION

The variability is classified as low if the coefficient of variation is ($< 10\%$), moderate ($10-20\%$) and high ($> 20\%$)(**Robinson et al. 1949**). High estimates of genotypic coefficient of variation (GCV) were recorded for number of pods per plant (27.38). The moderate GCV was recorded for seed yield per plant (17.04), number of seeds per pod (14.11), biological yield per plant (14.02), seed index (13.24), plant height (13.10), number of clusters per plant (12.70), pod length (12.30) and number of primary branches per plant (10.50). Low GCV was recorded for days to maturity (4.96), days to 50% pod setting (6.78), days to 50% flowering (8.02), and harvest index (8.90).

High estimates of phenotypic coefficient of variation (PCV) were recorded for number of pods per plant (31.32) and seed yield per plant (20.43). The moderate PCV was recorded for biological yield per plant (17.20), number of seeds per pod (16.07), number of clusters per plant (15.87), plant height (14.76), pod length (14.07), seed index (13.37), number of primary branches per plant (12.44) and harvest index (10.07). Low PCV was recorded for days to maturity (6.46), days to 50% pod setting (7.21), and days to 50% flowering (9.74).

The estimates of phenotypic coefficient of variation were found higher than their corresponding genotypic coefficient of variation indicating that apparent variation is not only due to genotype but also due to the influence of the environment. Therefore, caution has to be exercised in making the selection based on phenotype alone can be effective for the improvement of these traits. Relatively higher differences between genotypic and phenotypic coefficient of variation were observed for number of pods per plant, seed yield per plant, biological yield per plant, and number of clusters per plant. The large difference between the GCV and PCV indicates a high environmental influence on the expression of particular traits.

Table 2: Analysis of Variance for 13 quantitative characters in Greengram

Sl. No.	Source	Mean Sum of Squares (MSS)		
		Replication	Treatment	Error
	Degrees of freedom	2	19	38
1	Days to 50% flowering	0.3170	29.996**	4.089
2	Days to 50% pod setting	8.717**	39.628**	1.646
3	Plant height (cm)	0.0350	43.101**	3.55
4	Number of branches per plant	0.010	1.314**	0.156
5	Days to maturity	8.150	45.168**	8.483
6	Number of clusters per plant	0.4720	1.968**	0.31
7	Number of pods per plant	5.2670	219.846**	20.517
8	Number of seeds per pod	0.730	7.311**	0.657
9	Pod length (cm)	0.1290	2.833**	0.264
10	Seed index (g)	0.0090	0.711**	0.004
11	Biological yield per plant (g)	13.9130	48.436**	6.983
12	Harvest Index (%)	1.7860	47.114**	4.02
13	Seed yield per plant (g)	3.2270	12.727**	1.619

* Significant at 5% level of significance

** Significant at 1% level of significance

Table 3: Mean and Index Scores for 13 characters of 20 greengramgenotypes during Zaid, 2023

Sl.	Genotypes	DF 50%	DP 50%	PH	NPBP	DM	NCP	NPP	NSPD	PDL	SI	BYP	HI	SYP	TIS
1	VBN-3	36(2.00)	48.67(1.00)	27.2(2.00)	5.85(2.00)	69(2.00)	5.67(2.00)	34.02(2.00)	11.73(2.00)	6.93(2.00)	3.66(2.00)	26.5(2.00)	41.5(2.00)	11.01(2.00)	25
2	CO-8	32.67(1.00)	46.67(1.00)	28.13(2.00)	6.86(3.00)	65.67(1.00)	5.07(2.00)	20.97(1.00)	10.93(2.00)	8.71(3.00)	4.9(3.00)	23.92(2.00)	49.43(3.00)	11.82(2.00)	26
3	LGG-407	37.67(2.00)	53.67(2.00)	28(2.00)	6.68(3.00)	76.67(3.00)	6.73(3.00)	50.29(3.00)	11.07(2.00)	7.17(2.00)	3.64(2.00)	30.15(2.00)	43.31(2.00)	13.06(2.00)	30
4	PUSA	36.33(2.00)	51.33(2.00)	29(2.00)	5.91(2.00)	72.67(2.00)	6(2.00)	22.83(2.00)	10.33(2.00)	7.61(2.00)	3.36(2.00)	24.77(2.00)	44.64(2.00)	11.03(2.00)	26
5	AMULYA	37.33(2.00)	58.33(3.00)	25.67(2.00)	6.66(3.00)	71.33(2.00)	5.8(2.00)	23.32(2.00)	10.07(2.00)	7.54(2.00)	4.21(3.00)	25.81(2.00)	43.1(2.00)	11.17(2.00)	29
6	IPM-2-3	38(2.00)	59(3.00)	31.4(2.00)	6.27(2.00)	69.67(2.00)	6.13(2.00)	27.42(2.00)	12.86(3.00)	9(3.00)	4.14(2.00)	26.91(2.00)	38.63(2.00)	10.39(2.00)	29
7	LGG-450	30.67(1.00)	57.67(3.00)	20.83(1.00)	4.61(1.00)	62.33(1.00)	4.07(1.00)	13.89(1.00)	8.33(1.00)	6.07(1.00)	3.81(2.00)	17.57(1.00)	45.13(2.00)	7.95(1.00)	17
8	SHAKTI	33.33(1.00)	49.67(2.00)	23.73(1.00)	5.99(2.00)	65.33(1.00)	6.46(2.00)	28.18(2.00)	9.93(2.00)	7.21(2.00)	4.16(3.00)	20.86(1.00)	44.19(2.00)	9.25(2.00)	23
9	IPM-2-14	39.67(2.00)	50.33(2.00)	25.47(2.00)	6.25(2.00)	70(2.00)	5.93(2.00)	24.98(2.00)	11.33(2.00)	7.83(2.00)	2.93(1.00)	23.59(2.00)	44.61(2.00)	10.49(2.00)	25
10	MGG-295	33.33(1.00)	51.67(2.00)	26.67(2.00)	5.41(2.00)	69.33(2.00)	5.93(2.00)	32.7(2.00)	8.66(1.00)	6.76(2.00)	3.4(2.00)	23.09(2.00)	45.91(2.00)	10.6(2.00)	24
11	CO-7	38.67(2.00)	54.33(2.00)	37.4(3.00)	4.96(1.00)	72.33(2.00)	6.27(2.00)	44.86(3.00)	14.86(3.00)	10.1(3.00)	3.22(2.00)	33.88(3.00)	45.93(2.00)	15.57(3.00)	31
12	PUSA-105	36.67(2.00)	48.33(1.00)	25.48(2.00)	5.27(2.00)	75.67(3.00)	5.26(2.00)	32.29(2.00)	11.07(2.00)	6.87(2.00)	3.27(2.00)	29.75(2.00)	41.78(2.00)	12.43(2.00)	26
13	VIRAT	41.33(3.00)	52.67(2.00)	27.67(2.00)	6.4(2.00)	74.67(3.00)	7.4(3.00)	35.14(2.00)	10.8(2.00)	8.22(2.00)	3.27(2.00)	31.94(3.00)	44.38(2.00)	14.15(3.00)	31
14	SHIKHA	41.33(3.00)	56(2.00)	28.6(2.00)	5.78(2.00)	74(2.00)	5.8(2.00)	35.58(2.00)	8.77(1.00)	6.49(1.00)	3.31(2.00)	29.14(2.00)	44.09(2.00)	12.85(2.00)	25
15	LGG-460	40.67(3.00)	52(2.00)	29.87(2.00)	7.16(3.00)	71.33(2.00)	6.4(2.00)	29.71(2.00)	7.86(1.00)	8.31(2.00)	3.61(2.00)	27.51(2.00)	42.85(2.00)	11.79(2.00)	27
16	TM-96-2	33.33(1.00)	47.67(1.00)	25.33(2.00)	5.29(2.00)	65.33(1.00)	6.4(2.00)	27.22(2.00)	10.4(2.00)	7(2.00)	3.58(2.00)	24.98(2.00)	42.31(2.00)	10.59(2.00)	23
17	SM-02-103	32.33(1.00)	49.67(2.00)	26.13(2.00)	5.58(2.00)	66.33(1.00)	6.87(3.00)	38.89(3.00)	10.4(2.00)	6.97(2.00)	3.4(2.00)	32.43(3.00)	44.78(2.00)	14.45(3.00)	28
18	MH-421	36.33(2.00)	55.67(2.00)	29.73(2.00)	6.33(2.00)	70.67(2.00)	5.47(2.00)	27.16(2.00)	10.46(2.00)	6.7(2.00)	3.29(2.00)	28.47(2.00)	37.76(1.00)	10.73(2.00)	25
19	SM-20-108	39.67(2.00)	51(2.00)	23.27(1.00)	5.65(2.00)	72.67(2.00)	4.73(1.00)	25.18(2.00)	10.67(2.00)	7.02(2.00)	3.65(2.00)	24.43(2.00)	34.46(1.00)	8.39(1.00)	22
20	SML-1668	37.33(2.00)	55(2.00)	34.6(3.00)	5.47(2.00)	74(2.00)	4.67(1.00)	20.83(1.00)	10.53(2.00)	7.98(2.00)	4.46(3.00)	24.67(2.00)	32.9(1.00)	8.12(1.00)	24

DF 50% = days to 50% flowering, DP 50%= days to 50% pod setting, PH= plant height (cm), NPBP= number of primary branches per plant, DM= days to maturity, NCP= number of clusters per plant, NPP= number of pods per plant, NSPD= number of seeds per pod, PDL= Pod Length (cm), SI= Seed Index (g),
 BYP= biological Yield per Plant, HI= Harvest Index, SYP= Seed Yield per Plant, TIS= total Index Score

3.3 HERITABILITY

The heritability is classified as low (< 30%), moderate (30- 60%) and high (> 60%) by **Johnson et al., 1955**. In the present study, heritability estimates were high for seed index (98.204), days to 50% pod setting (88.492), plant height (78.786), harvest index (78.134), number of seeds per pod (77.139), pod length (76.458), number of pods per plant (76.406), number of primary branches per plant (71.223), seed yield per plant (69.582), days to 50% flowering (67.868), biological yield per plant (66.429) and number of clusters per plant (64.044) while it was moderate for days to maturity (59.04). No characters were observed with low heritability.

Higher values of broad sense heritability for the traits indicate that those characters are less influenced by environmental effects thus helping in the effective selection of the traits based on the phenotypic expression by adopting a simple selection method and suggesting the scope of genetic improvement.

3.4 GENETIC ADVANCE AS PERCENT OF MEAN

Genetic advance as percent of mean was categorized as low (less than 10%), moderate (10-20%), and high (more than 20%) (**Johnson et al., 1955**). In the present investigation, characters number of pods per plant (49.30), seed yield per plant (29.28), seed index (27.04), number of seeds per pod (25.53), plant height (23.96), biological yield per plant (23.54), pod length (22.16) and number of clusters per plant (20.94) recorded high genetic advance as percent of mean. While it was recorded as moderate for number of primary branches per plant (18.25) followed by harvest index (16.21), days to 50% flowering (13.61), and days to 50% pod setting (13.14) and it was recorded as low for days to maturity (7.86). Respectively suggesting that these characters might be viewed as desirable features for selection improvement, which may be attributed to additive gene action and so could be improved upon by modifying selection without progeny testing.

Table 4: Genetic variability parameters of 13 quantitative characters in greengram

Sl. No.	Genetic Parameters	GCV	PCV	h^2 (bs)	GA	GAM
1	Days to 50% flowering	8.02	9.74	67.868	4.99	13.61
2	Days to 50% pod setting	6.78	7.21	88.492	6.90	13.14
3	Plant height (cm)	13.10	14.76	78.786	6.64	23.96
4	Number of branches per plant	10.50	12.44	71.223	1.08	18.25
5	Days to maturity	4.96	6.46	59.04	5.54	7.86
6	Number of clusters per plant	12.70	15.87	64.044	1.23	20.94
7	Number of pods per plant	27.38	31.32	76.406	14.68	49.30
8	Number of seeds per pod	14.11	16.07	77.139	2.70	25.53
9	Pod length (cm)	12.30	14.07	76.458	1.67	22.16
10	Seed Index (g)	13.24	13.37	98.204	0.99	27.04
11	Biological yield per plant (g)	14.02	17.20	66.429	6.24	23.54
12	Harvest Index (%)	8.90	10.07	78.134	6.90	16.21
13	Seed yield per plant (g)	17.04	20.43	69.582	3.31	29.28

PCV: Phenotypic Coefficient of Variation, GCV: Genotypic Coefficient of Variation, h^2 bs: heritability (broad sense), GA: Genetic Advance, GAM: Genetic Advance as Percent of

Mean




Fig. 1: Histogram depicting GCV, PCV, heritability, and genetic advance for 13 quantitative characters of greengram genotypes

3.5 METROGLYPH ANALYSIS

Anderson's metroglyph analysis is a semi-graphical method for assessing the pattern of morphological variation in germplasm lines. It was observed that maximum variability was in number of pods per plant (13.89-50.29) followed by plant height (20.83-37.4), harvest index (32.9-49.43), biological yield per plant (17.57-33.88) and days to maturity (62.33-76.67). The range of mean values was utilized to assess the index scores 1, 2, and 3 for all the characters studied. The simple circle without rays represents index score 1, while others with values for index scores 2 and 3 have short and long rays on respective circles in different directions, respectively. The total index score varied from 17 (LGG-450) to 31 (CO-7, **VIRAT**) with a mean of 25.8. The highest total index score of 31 was recorded in genotypes CO-7 and **VIRAT** (**CHECK**) followed by a total index score of 30 in LGG-407 and a total index score of 29 in AMULYA and IPM-2-3. Hence, among the 20 germplasm lines, CO-7, **VIRAT** (**CHECK**), LGG-407, AMULYA, and IPM-2-3 were observed as high-yielding and identified for the highest index scores. In all these genotypes, number of pods per plant, plant height, and harvest index were contributory characters for more index scores.

Table 5: Index scores and signs used for characters for Metroglyph analysis of 20 genotypes of Greengram

Sl. No.	Character	Range of Mean	Score 1	Sign	Score 2	Sign	Score 3	Sign
			Value <		Value from - to		Value >	
1	Days to 50% flowering	30.67-41.33	33.47	○	33.47-39.8	♀	39.80	♀
2	Days to 50% pod setting	46.67-59	48.83	○	48.83-56.1	♂	56.10	♂
3	Plant height (cm)	20.83-37.4	23.92	○	23.92-31.5	○	31.50	○
4	Number of branches per plant	4.61-7.16	5.26	○	5.26-6.58	♂	6.58	♂
5	Days to maturity	62.33-76.67	66.57	○	66.57-74.33	♂	74.33	♂
6	Number of clusters per plant	4.07-7.4	5.04	○	5.04-6.66	♂	6.66	♂
7	Number of pods per plant	13.89-50.29	21.21	○	21.21-38.33	○	38.33	○
8	Number of seeds per pod	7.86-14.86	8.99	○	8.99-12.11	♀	12.11	♀
9	Pod length (cm)	6.07-10.1	6.55	○	6.55-8.5	♂	8.50	♂
10	Seed Index (g)	2.93-4.9	3.18	○	3.18-4.15	♀	4.15	♀
11	Biological yield per plant (g)	17.57-33.88	22.50	○	22.5-30.54	♂	30.54	♂
12	Harvest Index (%)	32.9-49.43	38.62	○	38.62-46.55	♀	46.55	♀
13	Seed yield per plant (g)	7.95-15.57	9.23	○	9.23-13.35	♀	13.35	♀

Table 6: Mean and Index Scores for 13 characters of 20 greengram genotypes

Sl.	Genotypes	DF 50%	DP 50%	PH	NPBP	DM	NCP	NPP	NSPD	PDL	SI	BYP	HI	SYP	TIS
1	VBN-3	36(2.00)	48.67(1.00)	27.2(2.00)	5.85(2.00)	69(2.00)	5.67(2.00)	34.02(2.00)	11.73(2.00)	6.93(2.00)	3.66(2.00)	26.5(2.00)	41.5(2.00)	11.01(2.00)	25
2	CO-8	32.67(1.00)	46.67(1.00)	28.13(2.00)	6.86(3.00)	65.67(1.00)	5.07(2.00)	20.97(1.00)	10.93(2.00)	8.71(3.00)	4.9(3.00)	23.92(2.00)	49.43(3.00)	11.82(2.00)	26
3	LGG-407	37.67(2.00)	53.67(2.00)	28(2.00)	6.68(3.00)	76.67(3.00)	6.73(3.00)	50.29(3.00)	11.07(2.00)	7.17(2.00)	3.64(2.00)	30.15(2.00)	43.31(2.00)	13.06(2.00)	30
4	PUSA	36.33(2.00)	51.33(2.00)	29(2.00)	5.91(2.00)	72.67(2.00)	6(2.00)	22.83(2.00)	10.33(2.00)	7.61(2.00)	3.36(2.00)	24.77(2.00)	44.64(2.00)	11.03(2.00)	26
5	AMULYA	37.33(2.00)	58.33(3.00)	25.67(2.00)	6.66(3.00)	71.33(2.00)	5.8(2.00)	23.32(2.00)	10.07(2.00)	7.54(2.00)	4.21(3.00)	25.81(2.00)	43.1(2.00)	11.17(2.00)	29
6	IPM-2-3	38(2.00)	59(3.00)	31.4(2.00)	6.27(2.00)	69.67(2.00)	6.13(2.00)	27.42(2.00)	12.86(3.00)	9(3.00)	4.14(2.00)	26.91(2.00)	38.63(2.00)	10.39(2.00)	29
7	LGG-450	30.67(1.00)	57.67(3.00)	20.83(1.00)	4.61(1.00)	62.33(1.00)	4.07(1.00)	13.89(1.00)	8.33(1.00)	6.07(1.00)	3.81(2.00)	17.57(1.00)	45.13(2.00)	7.95(1.00)	17
8	SHAKTI	33.33(1.00)	49.67(2.00)	23.73(1.00)	5.99(2.00)	65.33(1.00)	6.46(2.00)	28.18(2.00)	9.93(2.00)	7.21(2.00)	4.16(3.00)	20.86(1.00)	44.19(2.00)	9.25(2.00)	23
9	IPM-2-14	39.67(2.00)	50.33(2.00)	25.47(2.00)	6.25(2.00)	70(2.00)	5.93(2.00)	24.98(2.00)	11.33(2.00)	7.83(2.00)	2.93(1.00)	23.59(2.00)	44.61(2.00)	10.49(2.00)	25
10	MGG-295	33.33(1.00)	51.67(2.00)	26.67(2.00)	5.41(2.00)	69.33(2.00)	5.93(2.00)	32.7(2.00)	8.66(1.00)	6.76(2.00)	3.4(2.00)	23.09(2.00)	45.91(2.00)	10.6(2.00)	24
11	CO-7	38.67(2.00)	54.33(2.00)	37.4(3.00)	4.96(1.00)	72.33(2.00)	6.27(2.00)	44.86(3.00)	14.86(3.00)	10.1(3.00)	3.22(2.00)	33.88(3.00)	45.93(2.00)	15.57(3.00)	31
12	PUSA-105	36.67(2.00)	48.33(1.00)	25.48(2.00)	5.27(2.00)	75.67(3.00)	5.26(2.00)	32.29(2.00)	11.07(2.00)	6.87(2.00)	3.27(2.00)	29.75(2.00)	41.78(2.00)	12.43(2.00)	26
13	VIRAT	41.33(3.00)	52.67(2.00)	27.67(2.00)	6.4(2.00)	74.67(3.00)	7.4(3.00)	35.14(2.00)	10.8(2.00)	8.22(2.00)	3.27(2.00)	31.94(3.00)	44.38(2.00)	14.15(3.00)	31
14	SHIKHA	41.33(3.00)	56(2.00)	28.6(2.00)	5.78(2.00)	74(2.00)	5.8(2.00)	35.58(2.00)	8.77(1.00)	6.49(1.00)	3.31(2.00)	29.14(2.00)	44.09(2.00)	12.85(2.00)	25
15	LGG-460	40.67(3.00)	52(2.00)	29.87(2.00)	7.16(3.00)	71.33(2.00)	6.4(2.00)	29.71(2.00)	7.86(1.00)	8.31(2.00)	3.61(2.00)	27.51(2.00)	42.85(2.00)	11.79(2.00)	27
16	TM-96-2	33.33(1.00)	47.67(1.00)	25.33(2.00)	5.29(2.00)	65.33(1.00)	6.4(2.00)	27.22(2.00)	10.4(2.00)	7(2.00)	3.58(2.00)	24.98(2.00)	42.31(2.00)	10.59(2.00)	23
17	SM-02-103	32.33(1.00)	49.67(2.00)	26.13(2.00)	5.58(2.00)	66.33(1.00)	6.87(3.00)	38.89(3.00)	10.4(2.00)	6.97(2.00)	3.4(2.00)	32.43(3.00)	44.78(2.00)	14.45(3.00)	28
18	MH-421	36.33(2.00)	55.67(2.00)	29.73(2.00)	6.33(2.00)	70.67(2.00)	5.47(2.00)	27.16(2.00)	10.46(2.00)	6.7(2.00)	3.29(2.00)	28.47(2.00)	37.76(1.00)	10.73(2.00)	25
19	SM-20-108	39.67(2.00)	51(2.00)	23.27(1.00)	5.65(2.00)	72.67(2.00)	4.73(1.00)	25.18(2.00)	10.67(2.00)	7.02(2.00)	3.65(2.00)	24.43(2.00)	34.46(1.00)	8.39(1.00)	22
20	SML-1668	37.33(2.00)	55(2.00)	34.6(3.00)	5.47(2.00)	74(2.00)	4.67(1.00)	20.83(1.00)	10.53(2.00)	7.98(2.00)	4.46(3.00)	24.67(2.00)	32.9(1.00)	8.12(1.00)	24

DF 50% = days to 50% flowering, DP 50%= days to 50% pod setting, PH= plant height (cm), NPBP= number of primary branches per plant, DM= days to maturity, NCP= number of clusters per plant, NPP= number of pods per plant, NSPD= number of seeds per pod, PDL= Pod Length (cm), SI= Seed Index (g), BYP= biological Yield per Plant, HI= Harvest Index, SYP= Seed Yield per Plant, TIS= total Index Score

The scatter diagram has been prepared by taking number of pods per plant on the x-axis and plant height (cm) on the y-axis and five complexes could be distinguished based on morphological variations. The scatter diagram revealed that five complexes could be distinguished based on morphological variations.

Complex - I: was represented by 10 genotypes which are VBN-3, CO-8, PUSA BAISAKHI, AMULYA, IPM-2-3, SHAKTI, IPM-2-14, MGG-295, PUSA-105, VIRAT, SHIKHA, LGG-460, TM-96-2, MH-421 and SM-20-108 with moderate number of pods per plant with moderate plant height with range value of 21- 28 and average index score of the complex 25.73.

Complex – II was represented by two genotypes which are LGG-407 and SM-20-103 with higher number of pods per plant with moderate plant height with a range value of 28-30 and an average index score of the complex 29.

Complex – III: was represented by one genotype LGG-450 with lower number of pods per plant with lower plant height with range and average index score of the complex 17.

Complex – IV: was represented by one genotype CO-7 with moderate number of pods per plant with higher plant height with range and average index score of the complex 24.

Complex – V: was represented by one genotype SML-1668 with higher number of pods per plant with higher plant height with range and average index score of the complex 31.

Table 7: Genotypes in different complexes in metroglyph analysis

Complex	Name of complex	No. of lines	Name of lines	Range and average score
I	Moderate number of pods per plant with moderate plant height	15	VBN-3, CO-8, PUSA BAIKAKHI, AMULYA, IPM-2-3, SHAKTI, IPM-2-14, MGG-295, PUSA-105, VIRAT, SHIKHA, LGG-460, TM-96-2, MH-421 and SM-20-108	22.00-31.00 (25.73)
II	Higher number of pods per plant with moderate plant height	2	LGG-407 and SM-02-103	28.00-30.00 (29.00)
III	Lower number of pods per plant with lower plant height	1	LGG-450	17
IV	Moderate number of pods per plant with higher plant height	1	CO-7	24
V	Higher number of pods per plant with higher plant height	1	SML-1668	31

Fig. 2: Scattered diagram of Metroglyph analysis showing 20 genotypes of greengram





Fig. 3: Frequency diagram of index scores of 20 greengram genotypes



4. CONCLUSION

From the present investigation, it is concluded that among the 20 genotypes of greengram based on mean performance, genotype CO-7 (15.573 g) possessed the maximum seed yield per plant over the check variety VIRAT (10.487 g). It is also concluded from the present study that all the 20 genotypes of the greengram showed significant differences among them for all the characters. Genotypes CO-7, AMULYA, IPM-2-14, SM-20-108, SHAKTI, and MGG-295 were identified as the best-performing genotypes for seed yield and its component characters. The highest GCV, PCV is recorded for number of pods per plant followed by seed yield per plant and biological yield. Also, the Phenotypic Coefficient of Variation values are higher than the Genotypic Coefficient of Variation for all the traits under study indicating the influence of environment on studied characters. Hence, the characters with high-range estimates of GCV should be given top priority during selection. The number of pods per plant recorded high variation among all other characters with high PCV and GCV and heritability coupled with genetic advance as percent of the mean. The scattered diagram was prepared for 20 greengram germplasm lines, which formed five complexes, these complexes comprised 10, 2, 1, 1, and 1 germplasm lines respectively.

Further, it can be concluded that the metroglyph technique was useful in identifying groups of genotypes with yield-enhancing traits among the collection of diverse genotypes thus suggesting its potential value in mungbean improvement. The germplasm lines CO-7, VIRAT, LGG-407, AMULYA, and IPM-2-3 recorded high index scores and fell into different clusters and can be used as parents in hybridization to have maximum variability of a good combination of characters.

If a character is to be improved, which is undesirable or otherwise weak in a genotype, the information furnished here will be helpful in identifying the cross(s) that can be attempted to obtain the desired results.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. Name of AI Technology: ChatGPT

2. Version and model: GPT-4

3. Source: OpenAI

REFERENCES

1. **Afroz, M., Thirumala, Rao. V., Sridhar, V. and Hari, Y. (2022).** Assessment of genetic variability in mung bean [*Vigna radiata* (L.) Wilczek.]. *The Pharma Innovation Journal*, 11(11): 1371-1373.
2. **Anderson, E. (1957).** A semigraphical method for the analysis of complex problems. Proc. National Academy of Sciences, Washington, USA, 43: 923-927.
3. **Ashok, Chhabra., Nilu, A. L. R., Singh, V. P. and Chhabra, A. (1985).** Preliminary classification of broad bean germplasm lines. *Haryana Agriculture University Journal of Research*, 19: 431-435.
4. **Atul, Bhargava., Sudhir, Shukla., Rakesh, Kumar. and Deepak, Ohri.(2009).** Metroglyph Analysis of Morphological Variation in *Chenopodium* spp. *World Journal of Agricultural Sciences*, 5 (1): 117-120.
5. **Bajrang, Lal. Jakhar., Sanjay, Kumar. Sanadya., Smrutishree, Sahoo. and M. Sharma. (2020).** Metroglyph analysis of Groundnut germplasms for the assessment of morphological variations and preliminary classification. *Journal of Pharmacognosy and Phytochemistry*, 9(3): 130-134.
6. **Bhadra, S. K. and Akhtar, M. (1991).** Genetic divergence for yield and its morphological Components of mungbean. *Journal of Breeding and Genetics*, 23: 127-136.
7. **Borah, H. K. and Hazarika, M. H. (1991).** Metroglyph and index score analysis of morphological variation in greengram (*Vigna radiata* (L.) Wilczek) genotypes. *Journal of the Assam Science Society*, 33(4): 1-7.
8. **Chandra, G. S., Lavanya, G. R. and Kulkarni, S. D. (2017).** Studies on Genetic diversity in Greengram (*Vigna radiata* (L.) Wilczek) for seed yield characters. *Journal of Pharmacognosy and Phytochemistry*, 6(6): 1765-1767.
9. **Datta, D., Mukherjee, B. K., Barua, N. S. and Das, S. P.(2013).** Metroglyph analysis of maize (*Zea mays* L.) inbreds for preliminary classification and group constellation.

African Journal of Agricultural Research, 8(45): 5659-5663.

10. **Venkata, D. Sravani., Vinay, Tiwari. and Lavanya, G. R. (2022).** Genetic Variation, Correlation and Metroglyph Analysis in Rice (*Oryza sativa* L.) for Grain Yield Characters. *International Journal of Plant and Soil Science*, 748–758.
11. **Degefa, I., Petros, Y. and Andargie, M. 2014.** Genetic Variability, heritability and genetic advance in Mung bean (*Vigna radiata* (L.) Wilczek). *Plant Science Today*, 1(2):94-98.
12. **Dewan, D. B., Islam, M. A. and Khaleque, M. A. (1992).** Metroglyph analysis of morphological variation in Indian mustard (*Brassica juncea* L.). *Agric. Sci.*, 62: 766 - 777.
13. **Ghulam, Abbas., Muhammad, Jawad. Asghar., Tariq, Mahmud. Shah. and Babar, Manzoor. Atta. (2010).** Genetic diversity in mungbean (*Vigna radiata* (L.) Wilczek) germplasm. *Pakistan Journal of Botany*, 42(5): 3485-3495.
14. **Gokulakrishnan, J., Sunil, K. B. and Prakash, M. (2012).** Studies on Genetic Diversity in mungbean (*Vigna radiata* L.). *Legume Research*, 35(1): 50-52.
15. **Haritha, S. (1999).** Studies on Variability, Genetic Divergence and Metroglyph Analysis in Mung Bean (*Vigna radiata* (L.) wilczek).
16. **Jha, U. C., Singh, D. P., Paul, P. J. and Lavanya, G. R. (2011).** Metroglyph analysis for morphological variation in chickpea (*Cicer arietinum* L.). *Madras Agricultural Journal*, 98:121–123.
17. **Charles, Wesly.K., Madala, Nagaraju. and Lavanya, G. R. (2020).** Estimation of genetic variability and divergence in Greengram [*Vigna radiata* (L.)] germplasm. *Journal of Pharmacognosy and Phytochemistry*, 9(2): 1890-1893.
18. **Munukuru, Likhitha. and Gaibriyal, M. Lal. (2023).** Study of Genetic Divergence Using Metroglyph Analysis in Maize (*Zea mays* L.). *International Journal of Plant & Soil Science*, 35 (18): 1771-1780.
19. **Thakur, N. R., Topnope, V. N. and Sai, Phanindra. K. (2018).** Meteroglyph analysis for morphological variation in chickpea(*Cicer arietinum* L.). *The Journal of Research ANGRAU*, 46(2) 52-57
20. **Punitha, D., Ganesamurthy, K. and Rajarathinam, S. (2010).** Metroglyph analysis of morphological variations in Sorghum germplasm collections. *Electronic Journal of Plant Breeding*, 1: 536-541.
21. **Chandirakala, R., Ameena, Premnath. and Manivannan, N. (2015).** Metroglyph

analysis on genetic diversity in germplasm accessions of sunflower (*Helianthus annuus* L.). *Journal of Oilseeds Research*, 32: 170-173.

22. **Shika, Udayasri., Lavanya, G. R. and andGabriyal, M. Lal. (2022).** Estimation of Variability and Genetic Divergence in Greengram [*Vigna radiata* (L.) Wilczek] for Yield Characters. *International Journal of Plant & Soil Science*, 34(23): 49-56.
23. **Singh, V. P. and Chowdhury, R. K. (1974).** Metroglyph and index score analysis of morphological variation in greengram (*Phaseolus aureus* L.). Haryana Agricultural University Journal of Research, 4(4): 296-299.
24. **Srikanth, Thipanni., Eswari, K. B. and Bhrameshwar, Rao. (2016).** Metroglyph analysis of morphological variation in greengram germplasm collections. *Advances in life sciences*, 5 (20).

UNDER PEER REVIEW