

# “Studies on the effect of integrated nutrient management on growth of chrysanthemum (*Dendranthema grandiflorum*) cv. Pusa shwet”

## ABSTRACT

The present investigation was conducted in the Experimental Farm, Department of Agriculture, Mata Gujri College, Fatehgarh Sahib, Punjab during year 2023 with eight treatments comprising of different doses of integrated nutrients viz: T<sub>1</sub> i.e., Control, T<sub>2</sub> i.e., VAM (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>3</sub> i.e., PSB (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>4</sub> i.e., Azospirillum (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>5</sub> i.e., VAM + PSB (0.05+0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>6</sub> i.e., VAM + Azospirillum (0.05 + 0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>7</sub> i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>8</sub> i.e., RDF (40:20:20 g/m<sup>2</sup>), with three replications in randomized block design. The maximum plant height (41.72cm), number of leaves per stem (12.81cm), number of stems per plant (12.29), stem length (15.19cm), diameter of main stem (8.12mm), plant spread (33.13cm<sup>2</sup>), Leaf length (37.20cm), were recorded in T<sub>7</sub> i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) while minimum plant height (28.94cm), number of leaves per stem (8.14), number of stems per plant (6.92), stem length (9.18cm), diameter of main stem (5.59mm), plant spread (17.67cm<sup>2</sup>), Leaf length (28.76cm), was recorded under T<sub>1</sub> i.e., control.

**Keywords:** Chrysanthemum, Azospirillum, Phosphate solubilizing bacteria, Vesicular- arbuscular mycorrhiza

## INTRODUCTION

The term "Gul-e-Daudi" or "Queen of the East" refers to chrysanthemum, which is scientifically known as *Dendranthema grandiflorum*. It belongs to the family Asteraceae. This incredibly beautiful short-day herb can be grown both as an annual and a perennial flowering plant (Ghafoor & Khan, 2002). The chromosomal number of this lovely flower is 2n = 18. The name "chrysanthemum," which is indigenous to Northern Europe and Asia, comes from the Greek words "Chryos" which means gold, and "Anthemion" which means flower.

They take up a lot of nutrients from the surrounding soil, flower crops react favorably to fertilizer application. Therefore, to ensure optimal flower output, greater dosages of chemical fertilizers in a balanced ratio are required. Fertilizers are necessary for flowers to develop properly and to produce a sufficient quantity and quality of blooms. Adding biofertilizers to chemical fertilizers improves the soil's physical, chemical, and biological qualities while also increasing the chemical fertilizers' efficiency and providing some nutrients. Combining the usage of multiple nutrient sources can result in sustainable harvests of high-quality flowers. Until now, little research has been done on the efficacy of integrated nutrition management in flower crops, particularly in chrysanthemum.

Chrysanthemums are grown for their production, and among their many cultural practices, INM is crucial in determining the growth, yield, and quality of the flowers. INM promotes a greater number of branches and leaf area on the plant, which results in a higher flower yield per plant (Patanwar *et al.* 2014). It has been suggested that using INM in different combinations will improve floral quality and yield. Therefore, farmers will benefit from the adoption of this approach.

## MATERIAL AND METHODS

The present investigation was conducted at Research Farm, Mata Gujri College, Fatehgarh Sahib, Punjab during 2023-2024. The usual climate of Fatehgarh Sahib is subtropical. The area involves maximum temperature ranges from 10.2°C to 35.60°C in the summer, while in the winter lowest temperature is between 4.50°C to 20.10°C. Chrysanthemum cuttings were planted on the backside of Mata Gujri College, Fatehgarh Sahib, on August 23, 2023. On September 26, 2023, 35 days after the rooting process started, the cuttings are transplanted in the main field of Kharaura. The experiment was laid out in Randomized Block Design (RBD) with three replications. The cuttings of uniform size (10-15cm) used for treatment of integrated nutrient management and treated with biofertilizers and then planted in the beds with a spacing of 30 cm plant to plant and 35 cm row to row during September 2023. On September 26, 2023, biofertilizers were applied to the plants prior weighing 0.05 g/m<sup>2</sup> each on weighing scale. The Azospirillum (0.05 g/m<sup>2</sup>), PSB (0.05 g/m<sup>2</sup>) and NPK (20:10:10 g/m<sup>2</sup>) were weighed and then mix with vermicompost for application to the growing crop chrysanthemum. There were 8 treatments of integrated nutrients i.e., T<sub>1</sub> Control, T<sub>2</sub> VAM (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>3</sub> PSB (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>4</sub> Azospirillum (0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>5</sub> VAM + PSB (0.05+0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>6</sub> VAM + Azospirillum (0.05 + 0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>7</sub> Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + NPK (20:10:10 g/m<sup>2</sup>), T<sub>8</sub> RDF (40:20:20 g/m<sup>2</sup>) which were applied on Pusa shwet cultivar of chrysanthemum. Land was prepared to a good tilth by ploughing and then leveling. Earthing up was done at 45 days after planting to provide sufficient support, greater soil volume for spread of corms and prevent lodging of plants.

Comment [cm1]: italic

Comment [cm2]: mention treatment details in materials and methods and mention the outcomes of the experiment in abstract area

Comment [cm3]: italics

Comment [cm4]: italics

Comment [cm5]: Integrated Nutrient Management (INM)

Comment [cm6]: Insert the field geographical coordinates and layout picture of the field

Comment [cm7]: Prior

## RESULTS AND DISCUSSION

The maximum plant height was measured in T<sub>7</sub> (41.72 cm), which was the combination of Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically at par with T<sub>2</sub> (39.29 cm), which was composed of VAM (0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>). Different nutrient levels had a substantial impact on the height of the plants. The favourable effects of biofertilizers Azospirillum + PSB combined with the recommended half dosage of (RDF) inorganic fertilizers may be the cause of the increase in plant height in treatment T<sub>7</sub>. Azospirillum may have contributed nitrogen through atmospheric fixation. Additionally, PSB may have assisted in providing the plant with phosphorus, which subsequently enlarged the flower's diameter. The minimum plant height was recorded in T<sub>1</sub> (28.94 cm) which was statistically inferior. Decrease in plant height might be brought on by insufficient supply of nutrition being available to the plant at vital periods for its luxuriant growth. Azospirillum + PSB with 50% RDF significantly increased the plant height of china aster, according to (Chaitra and Patil's 2007) observations. (Deshmukh *et al.* 2008) also reported increased gaillardia plant height when 50% NPK, Azospirillum and PSB were used as seedling inoculants.

There were more number of leaves per stem was noticed in T<sub>7</sub> (12.81) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically superior. Because nitrogen is a crucial component of protein and chlorophyll, which promotes greater development and in the treatment T<sub>7</sub> had the maximum leaf count per plant can be linked to its enhanced availability. T<sub>1</sub> (8.14) i.e., control, had the minimum number of leaves on the stem, which was statistically at par with T<sub>4</sub> (8.52) i.e., Azospirillum (0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>). Decrease in number of leaves per plant could result from the plant receiving insufficient nourishment during critical stages necessary for its vegetative development. Nandre *et al.* (2005) observed similar findings in China aster, and Shashidhara and Gopinath (2005) reported similar findings in calendula.

T<sub>7</sub> (12.29) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) had the maximum number of stems per plant was noticed which was statistically superior. Better micronutrients and macronutrients flow as well as plant growth substances into the plant system in the plots treated with Azospirillum + PSB with 50% RDF that may be the cause of maximum number of stems per plant in T<sub>7</sub>. T<sub>1</sub> (6.92) i.e., control had the minimum number of stems per plant which was statistically at par with T<sub>4</sub> (7.71) i.e., Azospirillum (0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>). The main reason for the reduction in the number of stems per plant could be the lack of sufficient nutrients at critical stage for the plant's vigorous growth. The findings of Kale *et al.* (1987) in salvia and Nethra (1996) in china aster also support the above conclusions.

Comment [cm8]: Add recent findings

The maximum stem length was observed in T<sub>7</sub> (15.19 cm) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically superior. The main reason of the increase in stem length may be the production of growth promoting compounds from the administration of PSB and Azospirillum as well as the provision of macronutrients through the half dose of inorganic fertilizers. T<sub>1</sub> (9.18 cm) i.e., control resulted smallest stem length which was statistically inferior. Decrease in stem length may result from the plant receiving insufficient nutrition during critical stages necessary for its lush development. Similar results was shown in, African marigolds treated with Azospirillum + PSB with 50% RDF showed considerably higher plants (Sunita *et al.* 2007).

Maximum diameter of the main stem was resulted in T<sub>7</sub> (8.12 mm) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically at par with T<sub>3</sub> (7.73 mm) i.e., PSB (0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>). This could be the result of the synergistic action of biofertilizers as well as inorganic fertilizers i.e., Azospirillum + PSB with 50% RDF which improved nutrient availability that improved root and shoot growth and ultimately favourably affected plant growth, branching out more per plant, causing more cell division and accumulating carbohydrates that resulted in a thick stem. The control i.e., T<sub>1</sub> (5.59 mm) had a minimum diameter of main stem which was statistically inferior. Insufficient nutrients available to the plant during critical times for its luxuriant growth may cause of decline in diameter of main stem. Similar results were also found in the cases of chrysanthemum by Chauhan (2005), marigold by Rathi *et al.*, (2005), marigold by Sunitha *et al.*, (2007) and China aster by Chaitra and Patil (2007).

Comment [cm9]: Inappropriate sentence

The largest plant spread was measured in T<sub>7</sub> (33.13 cm<sup>2</sup>) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically superior. In addition to making fixed phosphorous in soil more soluble and plant-available, bioinoculants and chemical fertilizers such as Azospirillum + PSB with 50% RDF may have been helpful in fixing atmospheric nitrogen and secreting growth promoting substances like auxin which may have increased plant metabolic activity, photosynthetic efficiency and improved plant growth and development (Kumar *et al.*, 2009). T<sub>1</sub> (17.67 cm<sup>2</sup>) i.e., control had the minimum plant spread which was statistically inferior. Insufficient nutrients supplied to the plant during critical times for its luxuriant development may cause a decline in plant spread.

Maximum leaf length was resulted in T<sub>7</sub> (37.20cm) i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) which was statistically at par with T<sub>2</sub> (36.46cm) i.e., VAM (0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>). In addition to facilitating the intake of microelements found in soil, mycorrhizal fungi also mobilize and absorb phosphate. It could possibly be because of the Azospirillum + PSB with 50% RDF which has made more N available to the plants. T<sub>1</sub> (6.92cm) i.e., control group had the minimum leaf length which was statistically inferior. Absence of enough nutrients at crucial points for plants luxuriant growth may be the cause of decrease in leaf length. The similar findings are recorded in *Calendula officinalis* by Sohn *et al.*, (2003), in *chrysanthemum* by HariPriya *et al.*, (2004) and in roses by Mostafa (2002).

**Table 1. Shows the mean performance of the vegetative as well as flowering parameters of Chrysanthemum (*Dendranthema grandiflorum*) cv. Pusa shwet as affected by different biofertilizer treatments during the year 2023-2024.**

**Comment [cm10]:** Insert the table number in the discussion and also show the graphical representation of the data

Treatments	Plant height	No. of leaves per stem	No. of stems per plant	Stem length	Diameter of main stem	Plant spread	Leaf length
T <sub>1</sub> Control	28.94	8.14	6.92	9.18	5.59	17.67	28.76
T <sub>2</sub> VAM (0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	39.29	9.45	8.46	11.25	7.41	26.48	36.46
T <sub>3</sub> PSB (0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	37.01	9.25	8.63	11.42	7.73	26.33	32.61
T <sub>4</sub> Azospirillum (0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	38.45	8.52	7.71	11.36	6.49	26.44	33.34
T <sub>5</sub> VAM + PSB (0.05+0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	36.35	9.10	8.46	12.49	6.14	26.50	33.35
T <sub>6</sub> VAM + Azospirillum (0.05 + 0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	37.75	9.15	8.27	12.20	6.30	26.27	33.47
T <sub>7</sub> Azospirillum + PSB (0.05 + 0.05 g/m <sup>2</sup> ) + NPK (20:10:10 g/m <sup>2</sup> )	41.72	12.81	12.29	15.19	8.12	33.13	37.20
T <sub>8</sub> RDF (40:20:20 g/m <sup>2</sup> )	36.60	9.29	8.21	12.81	6.59	26.44	33.15
SEM±	0.90	0.18	0.26	0.38	0.16	0.27	0.26
CD	2.13	0.73	0.77	1.16	0.48	0.81	0.59

### CONCLUSION

From the presented research it can be concluded that T<sub>7</sub> i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) is statistically superior in all the vegetative parameters, including plant height(41.72cm), plant spread(33.13cm<sup>2</sup>), number of leaves per stem(12.81), diameter main stem(8.12mm), number of stems per plant(12.29), leaf length(37.20cm), and stem length(15.19cm).

T<sub>7</sub> i.e., Azospirillum + PSB (0.05 + 0.05 g/m<sup>2</sup>) + N:P:K (20:10:10 g/m<sup>2</sup>) results best both in almost all the parameters. So it is suggested for the cultivation of chrysanthemum cv. Pusa Shwet.

### REFERENCES

**Comment [cm11]:** Insert the latest references

- Chaitra, R and Patil, V S. 2007. Integrated nutrient management studies in china aster (*Callistephus chinensis* Nees) Cv. Kamini. Karnataka J. Agril. Sci., **20** (3) : 689-690.
- Chauhan PA Effect of biofertilizers and chemical nitrogenous fertilizer on growth and flower yield of chrysanthemum (*Chrysanthemum morifolium*). M.Sc. Thesis (Unpublished). AAU, Anand c (2005).
- Deshmukh, P G, Khiratar, S D, Badge, S A and Bhongle, S A. 2008. Effect of bio-inoculants with graded doses of NPK on growth and yield of gaillardia. J. Soils Crops, **18** (1) : 212-216.

- Ghafoor, A and R Khan. 2002. Anthemideae. In: Flora of Pakistan, (Eds.): S.I Ali & M. Qaiser. University of Karachi, Missouri Botanical Garden, Press, **207**: 45-47.
- Haripriya, K, Senthilkumar, S and Sriramchandrasekaran, M V. 2004. Effect of vermicompost and fertilizer on the growth and yield of rose. *Journal of Interacademia*, **8**(2): 207-210.
- Hemavathi, M. 1997. Effect of organic manures and bio-fertilizers on growth and productivity of chrysanthemum (*Chrysanthemum morifolium Ramat.*). Cv. Local yellow. *M. Sc. (Agri.) Thesis*, Univ. Agric. Sci., Bangalore (India).
- Kale, R D, Bano, K, Sreenivasa, M N and Bagyaraj, D J. 1987, Influence of worms cast on the growth and mycorrhizal colonization of two ornamental plants. *South Indian Hort.*, **35**: 433-437.
- Kumar, D, Singh, B P and Singh, V N. 2009. Effect of integrated nutrient management on growth, flowering behaviour and yield of African marigold (*Tagetes erecta* L.) cv. African Giant Double. *J. Hort. Sci.*, **4**(2) : 134-137.
- Kumar, M, Sharma, S K, Singh, S, Dahiya, D S, Mohammed, S and Singh, V P. 2006. Effect of farm yard manure and different biofertilizers on yield and nutrient content of marigold cv. Pusa Narangi. *Haryana J. Hort. Sci.*, **35**(3-4): 256-257.
- Mashaldi, A. 2000. Effect of organic and inorganic fertilizers on growth, yield and post harvest life of marigold (*Tagetes erecta* L.) cv. Double orange. *M. Sc. (Ag.) Thesis*, UAS, Dharwad, Karnataka.
- Mostafa, M M. 2002. Effect of biofertilizer, salinity and negative technique on the growth of some annuals plants. *Alexandria Journal of Agriculture Research*, **47**(2): 151-162.
- Nandre, D R, Jogdande, N D, Dalal, S R, Bansode, A B and Chaudhale, B S. 2005. Effect of Azotobacter on growth and yield on China aster under reduced nitrogen levels. *Advances in Plant Sciences*, **18**(1): 87-89.
- Patanwar, Mridubhashini, G Sharma, Chetna Banjare, Deepika Chandravanshi and Eshu Sahu. 2014. Growth and development of chrysanthemum (*Dendranthema grandiflora* Tzvelev) as influenced by integrated nutrient management. **6**(Special issue): 459-462.
- Pravina, T, Damke, M M, Gonge, V S and Deshmukh, S. 2007. Influence of integrated nutrient management on vegetative growth parameters of marigold (*Tagetes erecta* L.). *Asian J. of Hort.*, **2**(2) : 33-36.
- Rathi SS, Parmar PB, Parmar BR. Influence of biofertilizers on growth and yield of African marigold (*Tagetes erecta* L.). *Gujarat Agriculture University Research Journal*. (2005) **30**(1-2):50-52.
- Sattar, M A and Gaur, A C. 1987. Production of auxins and gibberellins by phosphate dissolving microorganisms. *Zentralbl Microbiol.*, **142** : 393-395.
- Shasidhara, G R and Gopinath, G. 2005. Growth, flowering, yield, quality and economics of calendula (*Calendula officinalis* Linn.) as influenced by nutrients and bio-inoculants. *Journal of Ornamental Horticulture*, **8**(4): 249-253.
- Sohn, K, Kim, K Y, Chung, S J, Kim, W S, Park, S M, Kang, J G, Rim, Y S, Cho, J S, Kim, T H and Lee, J H. 2003. Effect of different timing of AMF inoculation on plant growth and flower quality of chrysanthemum, *Scientia Horticulturae*, **98**: 173-183.
- Sunita, H M, Ravi hunje, Vyakaranahal, B S and Bablad, H B. 2007. Effect of plant spacing and integrated nutrient management on yield and quality of seed and vegetative growth parameters in African marigold (*Tagetes erecta* Linn.). *J. Orna. Hort.*, **10** (4) : 245-249.