

Review Article

Nutritional benefits of Kodo and Little millet and their role in Dairy industry

ABSTRACT

Aim: The aim of the study is to find out benefits of millets and its role in dairy sector.

Millets are grass crops that have gained importance due to their excellent nutritional profile as well as their adaptability to climate change. Unlike conventional crops like rice and wheat, millets can grow in extreme conditions like drought, hot and arid conditions etc. The major and minor millets both have their benefits. Minor millets include kodo millet, barnyard millet (sawa), little millet (kutki), foxtail millet (Kakum), proso millet (chenna) and two pseudo millets which does not belong to grass family i.e., amaranth (rajkira) and buckwheat (kuttu). These millets have high mineral content compared to other conventional crops. Inclusion of these millets in our daily diet can help in combating malnutrition and deficiency diseases. These crops also have medicinal benefits like lowering the risk of cardiovascular diseases, celiac disease etc. In addition, they are good source of antioxidants like phytates, polyphenols, tannins etc. They are also good source of dietary fiber, have low glycemic index and hence can be a staple food for persons suffering from Diabetes. Millets are an excellent choice for value addition in food processing sector particularly dairy sector. India is the largest producer of millets and they are called as "NUTRI-CEREALS" or "Shree Anna". It has been a part of diet of tribal people of Jharkhand, Karnataka, Uttarakhand, Orissa and Madhya Pradesh. In this paper, the benefits of kodo and little-millets and their role in the dairy industry have been presented.

Keywords: Nutri-cereal, Kodo millet, Little millet, Bioactive compounds, value addition

1. INTRODUCTION

Since 2018, due to climate change and the lack of nutritional profile of conventional crops, there has been a lot of interest shown by the Government of India in promoting millets. The same year, millets were given the tag of "NUTRI-CEREALS" and 2018 was declared as the National year of millets. India's government had asked the UN to declare 2023 the International Year of Millets (IYOM) for which 72 nations agreed with India's request, and on March 5, 2021, the United Nations General Assembly (UNGA) proclaimed 2023 as the "International Year of Millets". Millets were also included in the Ministry of Women & Child Development's "POSHAN MISSION" Abhiyan.

Comment [H1]: Revised Abstract:

Aim: This study aims to evaluate the nutritional benefits of Kodo and Little millet compared to other cereals and explore their applications in the dairy sector.

Millets, including Kodo and Little millet, are increasingly recognized for their exceptional nutritional profiles and resilience to climate challenges. Unlike conventional cereals like rice and wheat, millets thrive in extreme conditions such as drought and heat. These grains are rich in minerals, antioxidants, and dietary fiber, making them vital in combating malnutrition and managing health conditions like diabetes and cardiovascular diseases. Their low glycemic index and high nutrient content also make them valuable for developing fortified dairy products. India, the largest producer of millets, has integrated these "Nutri-Cereals" into traditional diets across various states. This paper presents the comparative nutritional benefits of Kodo and Little millet and examines their potential to enhance the dairy industry's offerings.

This version emphasizes the study's purpose and focuses more on the dairy industry's context, making it more aligned with your research topic.

India is the largest producer of millet with 17.68% of the share in total production during the year 2020 followed by Nigeria (-11%) and China (-9%). In 2022, India produced 17.60 Million Metric Tonnes (MMT) of millets contributing to 19% of the total millet production in the world. With 17.75 MMT, India was the biggest consumer of millets in 2022, followed by China (13.70 MMT) and Nigeria (8.80 MMT). Also, India is placed 10th in terms of volume with 1.68 lakh MT (0.45% of global commerce), and seventh in terms of value, with exports of millet totaling USD 65.10 Million (1.66% of global trade). Globally, there were 40.44 Million hectares (ha) of sorghum planted in 2022, compared to 31.28 million hectares (ha) of other millets. Sorghum accounted for more than 66% (60.13 MMT) of total millets output, with the remaining 34% coming from other millets (small millets, finger millets, pearl millets, and so on. (YES BANK AND APEDA, 2022)

Comment [H2]: This reference is not properly cited. The reference should be included while the period follows. This reference should be the first one i.e. [1]

According to **USDA**, small millets account for 11% of the total millet production in India. Madhya Pradesh is the highest producer of small millets (33%) followed by Arunachal Pradesh, Karnataka and Tamil Nadu (9%), Chhattisgarh (7%), Uttarakhand (6%) and Gujarat and Andhra Pradesh (5%). The average production of millets during the last 5 years (2018-19 to 2022-23) is around 12,189 thousand tonnes.

Minor millets include Kodo millet or cow grass (*Paspalum scrobiculatum*) barnyard millet or sawa (*Echinochloa esculenta*), little millet or kutki (*Panicum sumatrense*), foxtail millet or Kakum (*Setaria italica*), proso millet or chenna (*Panicum miliaceum*) and two pseudo millets which does not belong to grass family include amaranth or rajkira (*Amaranthus caudatus* L.) and buckwheat or kuttu (*Fagopyrum esculentum*).

Comment [H3]: Check the spelling

2. KODO MILLET

2.1 Botanical Description

Kodo millet or cow grass belongs to family of *Poaceae* or also known as grass family. It is a long duration crop. The inflorescence of kodo millet consists of 2–6 racemes that are widely dispersed along a short axis or subdigitate. The grain is protected by sticky, persistent, hard husks. Self-pollination is the norm since Kodo millet possesses a cleistogamous flower and because the proportion of open flowers does not surpass 15-20% [1]. While *Paspalum scrobiculatum* var. *commersoniis* the wild variety native to Africa, *Paspalum scrobiculatum* var. *scrobiculatum* is farmed as an important crop in India. It is a monocot plant with approx. 4 feet in height. The leaves are 20-40 cm long. It produces small, ellipsoidal seeds that are 1.5 mm wide and 2 mm long and range in hue from light brown to dark grey given in Figure 1 [2]. The species *Paspalum scrobiculatum* is split into three races: **Regularis, Irregularis, and Variabilis** based on the morphology of the inflorescence. (ICRISAT, 2017). During 2015-16 the area of production of kodo millet was 1.96 lakh hectare, the production yield was 0.84 lakh tonnes and its total yield was 429 kg/ha. Total 40 improved varieties have been released from 1989 to 2022 (Indian Farming, 2023). Various varieties released by the Indian Institute of Millet Research (IIMR); Hyderabad is given in Figure 2.

Comment [H4]: delete

Comment [H5]: Unbold all these names

Comment [H6]: Kodo. Capital 'K' should start the name anywhere it appears.



Figure 1: Kodo millet seeds




Figure 2

Source: IIMR - SALIENT FEATURES OF RELEASED VARIETIES OF KODO MILLET CROPS (1989 TO 2012)

Comment [H7]: I will suggest that you use the pie chart with keys stated by the side. The keys will indicate the colours and their names

Comment [H8]: Source must followed numbered referencing E.g. [4]

2.2 Nutritional Composition of Kodo millet

According to Muragodet *et al.*, 2019, the physico-chemical composition of kodo millet includes its size (1.7 mm), thousand grain weight (2.8 gm), thousand grain volume (1.2 ml), bulk density (1.84), hydration capacity in gm per 1000 seeds is 0.54 while its hydration index is 24.52%. The swelling capacity in gm per 1000 seeds and swelling index is 0.55 and 42.30%, respectively. The proximate composition per 100 gm of kodo millet includes its moisture (11.2%), fat (1.3 g), protein (8.1 g), carbohydrate (64.3 g) and fiber (8.3 mg). The kodo millet also has significant quantities of minerals like Iron (0.5 mg), phosphorus (169 mg) and calcium (32 mg) [3]. Another author also reported similar findings which said that kodo grains have 8.35% protein, 1.45% fat, 65.65% carbohydrate, and 2.95% ash. The

Comment [H9]: This should follow the citation pattern. E.g. According to Muragodet *et al.* [3]

Comment [H10]: Kodo

Comment [H11]: Kodo. Please do for others. It doesn't matter if it start a sentence or not. It's a name.

study also revealed that kodo millets are high in vitamin B₃, vitamin B₆, folic acid, and minerals including calcium, potassium, magnesium, and zinc, as well as having medical advantages for diabetes, celiac disease, cancer, cardiovascular disease, and so on [4]The vitamin and mineral profile is given in Table 1. The energy value of kodo millet is between 340-342 kcal/100 gm[5].Kodo millet has more ash and dietary fiber content as compared to wheat and rice[6,7](Table 2)

UNDER PEER REVIEW

Table 1: Vitamin and mineral profile of millets compared to cereals

| S.no | Name of cereal/ millets | Vitamin B ₁ (mg) | Vitamin B ₂ (mg) | Vitamin B ₃ (mg) | Total folates (µg) | Iron (mg) | Phosphorus (mg) | Calcium (mg) | Sodium (mg) | Magnesium (mg) |
|------|-------------------------|-----------------------------|-----------------------------|-----------------------------|--------------------|-----------|-----------------|--------------|-------------|----------------|
| 1. | Whole wheat | 0.46 | 0.15 | 2.68 | 30.09 | 3.97 | 315 | 30.94 | 2.04 | 125 |
| 2. | Rice | 0.05 | 0.05 | 1.69 | 9.32 | 0.002 | 96 | 7.49 | 2.34 | 19.3 |
| 3. | Little millet | 0.26 | 0.05 | 1.29 | 36.20 | 1.26 | 130 | 16.06 | 4.77 | 91.41 |
| 4. | Kodo millet | 0.29 | 0.2 | 1.49 | 39.49 | 2.34 | 101 | 15.27 | 3.35 | 122 |

Source: Dayakar Rao et al., 2017 (Indian Institute of Millet Research); Dey et al., 2022

Table 2: Nutritional comparison of millets with wheat and rice per 100 gm

| S.no | Name of cereal and millets | Moisture (g) | Fat (g) | Ash (g) | Protein (g) | Carbohydrate (g) | Total dietary fiber (g) | Energy (Kcal) |
|------|----------------------------|--------------|---------|---------|-------------|------------------|-------------------------|---------------|
| 1. | Whole wheat | 10.58 | 1.47 | 1.42 | 10.6 | 64.70 | 11.23 | 1347 |
| 2. | Rice | 9.93 | 0.52 | 0.56 | 7.94 | 78.24 | 2.81 | 1491 |
| 3. | Little millet | 14.23 | 2.55 | 1.72 | 8.92 | 65.55 | 6.3 | 1449 |
| 4. | Kodo millet | 11.6 | 4.2 | 2.95 | 10.6 | 59.2 | 10.2 | 346 |

Source: Dayakar Rao et al., 2017 (Indian Institute of Millet Research), Indian superfood millet APEDA report; Bunkar et al., 2021

Comment [H12]: 'kcal' not Kcal

2.3 Bioactive compounds in Kodo millet



Malnutrition is common nowadays because individuals are not obtaining adequate nutrition from traditional crops, and dining out is popular in cities, which boosts junk food intake among children. This culture is the root cause of many ailments in today's society including cancer. It may be avoided in the future by replacing wheat and rice with millets, which are high in bioactive chemicals.

Kodo is an ancient millet grain that originated in tropical Africa and was domesticated 3000 years ago in India. This Indian cow grass is also referred to as Kodra and Varagu. The three categories of bioactive chemicals include alkaloids, terpenes and terpenoids, and phenolic compounds which includes phytochemicals such as Naringin, catechin, luteolin, apigenin, daidzein, naringenin etc. These compounds have various health benefits [8]. They are also called secondary metabolites. Secondary metabolites in plants are divided into three categories based on their metabolic pathway: (a) nitrogen-containing chemicals like alkaloids, glucosinolates, and cyanogenic glycosides, (b) phenolic compounds like phenylpropanoids and flavonoids, and (c) terpenes. Alkaloids are a type of nitrogen-containing chemicals generated in plants in response to biotic or abiotic stimuli, endowing alkaloids with extraordinary biological activity while shikimate, pentose phosphate, and phenylpropanoid pathways in plants are the sources of phenolic components, which include an aromatic ring with one or more hydroxyl groups. Also, terpenoids are the most prevalent category of secondary metabolites generated by plants and are found in flowers, vegetative tissues, and roots. They exhibit a wide spectrum of biological actions that lead to reduced total cholesterol, triglycerides, or LDL cholesterol, as well as blood pressure [9]. Some bioactive compounds reported by Khare *et al.*, 2020 in Kodo millets are Naringin (11.9 ppm), Catechin (1.10 ppm), Ferulic acid (2.45 ppm), Sinapic Acid (7.9 ppm) and *P*-Coumaric acid (1.38 ppm)[10]. Another author reported that the concentration of campesterol in raw kodo millet was 0.31 % while after germination the concentration increased to 2.6 % [11]. The protective functions of different bioactive compounds are given in Table 3



Comment [H13]: ...reported by Khare et al. [10]. Follow numbered referencing. If you want to follow Vancouver style, just follow it strictly.

Comment [H14]: Put a full stop or period at the end of every sentence. You've done this repeatedly. Please go over the work to see others.



Table 3: Bioactive compounds found in Kodo millet and their protective functions.

| S.no | Name of bioactive compound | Molecular Formula | PubChem CID | Molecular Structure | Nature of Protective functions | Reference |
|------|----------------------------|---|-------------|---|---|------------|
| 1. | Naringin | C ₂₇ H ₃₂ O ₁₄ | 442428 |  | Anti-oxidant and anti-inflammatory, treatment of Parkinson's disease, Alzheimer's disease lowers cholesterol and reduces the risk of CVDs | [10,12,13] |
| 2. | Catechin | C ₁₅ H ₁₄ O ₆ | 9064 |  | Anti-inflammatory, Anti-cancer, Anti-oxidative properties | [10,14] |

Comment [H15]: 'F' Functions

| S.no | Name of bioactive compound | Molecular Formula | PubChem CID | Molecular Structure | Nature of Protective functions | Reference |
|------|----------------------------|--|-------------|--|---|------------|
| 3. | Ferulic acid | C ₁₀ H ₁₀ O ₄ | 445858 |  | Anti-oxidant, anti-microbial, anti-viral, anti-cancerous, vasodilator activity and anti-diabetic activity | [10,15] |
| 4. | Sinapic Acid | C ₁₁ H ₁₂ O ₅ | 637775 |  | Anti-viral property against SARS Cov, Anti-inflammatory, Anti-cancer | [10,16,17] |

Comment [H15]: 'F' Functions

| S.no | Name of bioactive compound | Molecular Formula | PubChem CID | Molecular Structure | Nature of Protective functions | Reference |
|------|----------------------------|--|-------------|---|---|---------------|
| 5. | <i>P</i> -Coumaric acid | C ₉ H ₈ O ₃ | 637542 |  | Anti-inflammatory, Anti-cancer, Anti-oxidant, treatment of Rheumatoid arthritis; Anti-obesity | [10,18,19,20] |
| 6. | Campesterol | C ₂₈ H ₄₈ O | 173183 |  | Lowers cholesterol, Anti-cancerous | [21] |

Source of images: PubChem

Comment [H15]: 'F' Functions

3. LITTLE MILLET

3.1 Botanical Description

Little millet (*Panicum sumatrense*) also known as Samai or kutki is a minor millet which is of *Poaceae* family. It is native of Indian subcontinent. It is self-pollinated with chromosome number 36. The inflorescence of the little millet is a panicle that is 15-45 cm long and 1-5 cm broad. The spikelet is 2.0-3.5 mm in length. At maturity, panicle branches are scabrous and drooping. The lower is sterile, whereas the top is fertile or bisexual with no rachilla extension. Spikelets are elliptical, acute and compressed dorsally [22]. *P. sumatrense* is divided into two subspecies based on inflorescence morphology: *psilopodium* and *sumatrense*. The race **Nana and Robusta** are subspecies of *Sumatrense*. The race Nana has two subraces: **Laxa and Erecta**, and the race Robusta has two subraces: **Laxa and Compacta** (ICRISAT 2017). The grain is 1.8-1.9 mm in length and is smooth on the surface. The major states involved in its cultivation are Orissa, Madhya Pradesh, Andhra Pradesh, Karnataka, Maharashtra and Gujarat. It is resistant to both waterlogging and drought [23]. During 2015-16 the area of production of little millet was 2.34 lakh hectare, the production yield was 1.27 lakh tonnes and its total yield was 544 kg/ha. Total 32 improved varieties have been released from 1989 to 2022 (Indian Farming, 2023). Various varieties released by IIMR of little millet with their average yield is given in Figure 3.




Figure 3

Source: IIMR [SALIENT FEATURES OF RELEASED VARIETIES OF LITTLE MILLET CROPS (1989 TO 2016)]

3.2 Nutritional composition

Comment [H16]: Kutki

Comment [H17]: Unbold the names here

Comment [H18]: (ICRISAT, 2017) number this reference to follow the pattern. E.g. [23].

Comment [H19]: hectares

Comment [H20]: Follow numbered referencing [24]

Comment [H21]: are

Comment [H22]: Use the pie chart with keys stated by the side. The keys will indicate the colours and their names

Comment [H23]: The source should also followed numbered referencing

Kamataret *al.*, 2013 investigated little millet specimens from Karnataka's study districts for physical properties and classified based on seed color. They were divided into four groups: light brown, reddish brown, creamish brown, and blackish brown. The grain's thousand grain weight ranged from 2.40 to 2.80 g, while its bulk density ranged from 0.81 to 1.40 g/ml. The grains' mean hydration capacity varied between 9.13 to 21.50 % whereas their swelling capacity ranged from 11 to 51 %. The grain required 10-15 minutes for cooking. After cooking, the Hydration capacity (160-260 %) and swelling capacity (220-280%) increased significantly [24]. Another author studied the physical characteristics of *Sukshema*, an enhanced cultivar of a local genotype of little millet. The results revealed that the grain's length, width, and thickness were 1.74 mm, 1.53 mm, and 1.09 mm, respectively. The solubility was 23.4 % and water absorption capacity was 0.88 g/g. The total dietary fiber was 8.81 g/100g. Total sugar was around 4.14 g/100g. Table 2 describes the proximate composition of little millet [25].

3.3 Bioactive components

Little millet also has various bioactive components which may have ability to protect against several diseases. One article reported that the Total phenolic content (TPC) content of little millet was around 521 mgGAE/100 g with compounds like Gallic acid, Proto-catechuic acid, Kaempferol, ferulic acid, p-hydroxy-benzoic acid, Gentisic acid, vanillic acid, syringic acid and trans-cinnamic acid [26]. Similarly, another also suggested that little millet can be used as functional food component to control postprandial hyperglycemia. The TPC content of 3 cultivars of little millet was reported between 23-24 μmol ferulic acid equivalents/g and Total soluble Flavonoid content was between 11.33-13.64 μmol catechin equivalents/g. Among the 3 cultivars, compounds identified were Catechin, luteolin, naringenin, quercetin, apigenin, quercetin, and kaempferol. According to the results shown by the study, kaempferol and luteolin were the most abundant flavonoids. The luteolin content was highest in CO4 cultivar (69.24 \pm 2.13 $\mu\text{g/g}$) while kaempferol (60.57 \pm 2.15 $\mu\text{g/g}$) was highest in CO2 cultivar of little millet. The DPPH scavenging activity was also highest in CO2 [27].

Luteolin is a flavonoid that occurs naturally as a glycosylated form in a variety of fruits and vegetables. It has antioxidant, anticancer, anti-inflammatory, and neuroprotective properties. Kaempferol is also a flavanol that belongs to the flavonoid family and is found in a variety of plant genera including Delphinium, Berberis, Camellia, Citrus, and others. Kaempferol is useful in a variety of inflammatory disorders, including cardiovascular, cancer, and neurological diseases. Kaempferol's anti-inflammatory properties are attributable to its reduction of inflammatory cell activity as well as its inhibition of proinflammatory cytokines and chemokine expression [28].

The molecular structure and protective functions of bio-actives molecules found in little millet are presented in Table 4.

Comment [H24]: Follow the numbered referencing. E.g. Kamatar et al. [23]

Comment [H25]: Check this statement and rewrite.

Comment [H26]: 'the ability' adjust the indent to align with others

Comment [H27]: While bolding it?




Comment [H28]: Where did you get this? Please check it.

Comment [H29]: The '2' should be subscript

Comment [H30]: Same thing here

1 **Table 4: Important bioactive components in little millet**

2

| S.no | Name of Compound | Molecular formula | PubChem CID | Molecular structure | Protective functions | Reference |
|------|------------------|--|-------------|---|---------------------------------|-----------|
| 1. | Kaempferol | C ₁₅ H ₁₀ O ₆ | 5280863 |  | Anti-inflammatory | [29] |
| 2. | Luteolin | C ₁₅ H ₁₀ O ₆ | 5280445 |  | Anti-inflammatory, Anti-cancer | [30] |
| 3. | Quercetin | C ₁₅ H ₁₀ O ₇ | 5280343 |  | Anti-inflammatory, Anti-obesity | [31] |

3

4. CHANGES IN THE NUTRITIONAL COMPOSITION OF MILLETS UNDER DIFFERENT PROCESSING CONDITIONS.

Dehusking, soaking, milling, germination, fermentation, malting, heating, and roasting are all part of the processing process. All these practices have significant impact on the nutritional and nutraceutical properties of millets.

Comment [H31]: 'a significant'

Germination is a natural process used to activate grains biologically in order to increase their nutritional and functional qualities. Soaking, sprouting, and drying are part of Malting process. Roasting, parboiling and puffing are examples of heat treatments. These treatments improve millet's eating quality, although their impact on nutritional qualities such as TPC vary according to the heating method [32]. Sprouting of millets can increase its nutritional profile such as essential amino acids, anti-oxidants, minerals etc. During germination, kernels are often soaked in water before being let to germinate in a controlled atmosphere. As a result, sprouting circumstances and water content have a significant influence on its metabolic process. Soaking the kernel at ambient temperature for a few to 24 hours, with a seed weight/volume ratio of 1:1.5 to 1:20, is a common practice performed prior to seed germination. The ideal temperature for germination is 20-30°C [33]. Soaking also reduces anti-nutritional factors like phytates in cereal crops. A comparative study of 5 different millets revealed that 24h soaking and 24h germination was superior for creating nutritionally enhanced millet products. 50% tannin content was reduced in little millet while in kodo millet it was reduced from 1.42 mg/g to 0.52 mg/g after treatment. Also, DPPH activity rose from 34.12% to 56.16% in little millet and from 56.71% to 78.34% in kodo millet after 24 hours of soaking and germination [34]

Comment [H32]: Were
Please work on your grammar and punctuations.

Guha *et al.*, 2015 also reported that processing conditions like germinating, steaming and roasting of little millet also influence the concentration of bioactive components. The comparison between native and processed millets shows that TPC content was highest in Roasted little millet. The decreasing order of TPC value was Roasted (521.0 mg GAE/100 g) > Steamed (485 mg GAE/100g) > germinated (453 mg GAE/100g) > Unprocessed (429 mg GAE/100g). The trend was similar in the case of Total Flavonoid content [35]

Comment [H33]: Follow numbered referencing.
Guha et al. [35]

Similar results were reported by another author where TPC content increased from 27 to 34 mg TAE/100g after 12 h of soaking and 48 h of germination and popping of kodo millet increased TPC content to 63.58 mg TAE/100g [36].

Comment [H34]: Put full stop.

Other studies also indicate that Protein content increased significantly from 6.7% to 7.9%, minerals increased from 232.82 to 251.73mg/100g, dietary fibers increased from 35.30 to 38.34g/100g, and GABA content increased from 9.36 to 47.43mg/100g, while tannins and phytates reduced from 1.603 to 0.234mg/100g and 1.344 to 0.997mol/kg, respectively, in sprouted Kodo millet sample. It was also seen that lipid content decreased from 3.6 to 2.8%. Similarly to prior investigations, the DPPH radical scavenging activity (%) was raised (67.35 to 91.34%) [37].

Roasting also influences the Bioavailability of nutrients. Singh *et al.*, 2018 reported that roasted samples of finger millet showed a reduction in phytochemical elements such as phenols and antioxidant activity. It may be due to exposure to heating. The fat and protein content also decreased while total carbohydrate increased after roasting. The iron content increased from 3.45 to 3.91 mg/100g and calcium was increased from 337-341 mg/100g [38].

Comment [H35]: Singh et al. [38]
Numbered referencing.
Be consistent in your indenting

Comment [H36]: This may be due to....

Fermentation is an ancient technique which is quite popular for increasing probiotic content in food along which influencing the nutrients present in food. One of the literatures reported that little millet fermented with *Saccharomyces boulardii* and *Lactobacillus acidophilus* compared to germinated one has significant impact on its nutrients. The results revealed that Protein content increased from 9.86 to 10.95 %, fat content decreased from 3.26 to 2.61 %, Ash content increased from 2.27 to 2.34 and carbohydrates reduced from 84.54 to 82.01 %. The phytic acid content was reduced as well upon fermentation. So, it can be inferred that germination and fermentation when combined will improve the nutritional quality of the food [39].

Similar study was reported on kodo millet malt beverage inoculated with 4 different strains namely *Pediococcus acidilactici*, *Lactobacillus plantarum*, *Lactobacillus fermentum*. The results were similar to fermented little millet. The protein, antioxidants and crude fat increased [40].

Comment [H37]: 'that'

Comment [H38]: 'and'

Comment [H39]: A similar study....

Comment [H40]: similar to 'those of fermented'

5. MILLETS IN DAIRY INDUSTRY

Value added products of millets has been gaining popularity since the last 5-6 years. In the past decade, weaning mixes, health drinks, imitation dairy products using millets were developed by the researchers. Millet's starch functionality is equivalent to that of other cereals. Millets are appropriate constituents in various food compositions for certain target groups due to their increased proportion of non-starchy polysaccharides, dietary fiber, and low glycemic index. Millet's alkaline composition also makes them beneficial for people suffering from acidosis and gastric ulcers. [41]. Some product categories are described below:

a) Health Drinks

Health drinks are gaining popularity among consumers due to its high nutrient profile. Several studies of milk and millet based health drinks indicate the interest among researchers. Kumar *et al.*, (2013) developed health drink powder by combining malted finger millet (*Eleusine coracana*), different pulse combinations, and skim milk powder. When compared to other health beverages available in the market, it has a fairly high protein (25.01%) and calcium (Ca-1018.7 mg/100 g) content. Sensory analysis revealed that it has a better acceptance among consumers and is also a cost-effective product [42]. Parvathi *et al.*, (2015) also developed a millet-based dairy health drink targeting malnutrition in children. The product was developed by utilizing pearl, kodo, and whole wheat that had been germinated, as well as pulses. The dried germinated millets were roasted until the husks split apart, and they were then ground into a fine flour in a burr mill. Green gram flour, roasted bengal gram flour, and skim milk powder were then added to the flour. The health mix's DPPH scavenging activity was found to be 0.664 0.08 mg/g. In every 100 gm of the standardized mix, there were 17.08% protein, 4.05% fibre, 20.68 mg of calcium, and 7.57 mg of iron [43]. Another study was done on health drink targeting diabetic people using millets like bajra and ragi. Organoleptic and nutritional composition as well as phytochemical analysis showed that multigrain health mix was a superior product and can be used for diabetic people [44]. Similarly, milk-based iron rich health drink was developed to combat malnutrition using sprouted and dried pearl millet flour, finger millet powder, malted soya flour, sugar powder,

Comment [H41]: gaining

Comment [H42]: delete

Comment [H43]: their

Comment [H44]: Rephrased as 'Several studies of milk- and millet-based health drinks have been of interest to researchers.'

Comment [H45]: Kumar et al. [42]

Comment [H46]: delete

Comment [H47]: delete

Comment [H48]: Compared to other....

Comment [H49]: drinks

Comment [H50]: Check spelling

milk powder and popped and milled amaranth seed powder in different combinations. The effect of sprouting on the nutritional content of a bajra-based health drink indicated that sprouting reduced the carbohydrate (67.46g), fat (5.30g), moisture, and fibre contents (1.12g) compared to the control (100% pearl millet flour). The fat content reduced due to action of lipolytic enzyme and protein content increased from 11.2 g to 13.54g[45]. Another author developed a millet-based health drink using germinated kodo millet, barnyard millet, small millet, and finger millet and compared it with other health drinks available on the market. The results revealed that apart from energy, a health drink made from minor millets contains vital nutrients and is economically feasible for the consumers[46].

b) Porridge/Kheer mix

A ready to constitute kheer mix powder using pearl millet, dairy whitener, sugar and cardamom was developed. The millet was autoclaved for 15 minutes at 121°C and incubated at 35°C for 30 minutes before making the kheer mix. RSM (Response surface methodology) was used to optimize the final product which showed that increase in dairy whitener had positive effect on OA (Overall Acceptability) score[47]. Similar study was conducted where nutrient-enriched millet-based composite flour using skimmed milk powder and vegetables was developed for children aged 6 to 59 months. Finger Millets were germinated and roasted and combined with pumpkin seeds, carrots, cowpea leaves, and skimmed milk powder to make a composite flour. The findings indicate that increasing the germination time improved the protein, protein digestibility, and total sugar content of millet flour by 0.03%, 0.79%, and 0.07%, respectively, while decreasing total phytates by 0.005 mg/g[48]. Also Bunkar *et al.*, (2020) developed instant Kodo Millet Based Porridge Mix using Kodo millet, skim milk powder with stevia @ 8%. The optimized product had 72.6± 4.34 g/100gm of anti-oxidant content and 72.6± 4.34 g/100gm of ash content[49].

c) Weaning mix

Weaning is a phase of transition in which the infant's nutrition changes in terms of source and consistency. Weaning is a phase when children are nutritionally vulnerable. To benefit from increased micronutrients, cereals in composite mixes might be substituted with millets in the formulation with milk. Both milk and millets can provide a balanced nutrition to children. Several studies were conducted one of which was on weaning food based on using Barnyard millet, foxtail millet, soyabean and skim milk powder. Millets were malted while soyabean was roasted to create the mix. The product was rich in protein (18.37 %) and Ash content (4%). The calcium and ascorbic acid content was also high. The sensory analysis indicated that product was highly accepted by the consumer with OA (Overall acceptability) score 7.7[50]. Similarly Balasubramanian *et al.*, (2011) also developed a weaning mix using pearl millet, barley, whey protein concentrate vegetable oil and skim milk powder. He used extrusion to create extrudates of raw and malted pearl millet and barley. The overall acceptability of optimized product was between 6.60 to 8.20 [51]. Another study was done on weaning mix for children using little, kodo, foxtail and finger millet in different proportions. All millets were germinated before using. In comparison to the wheat-based market sample, millet-based weaning mixtures were higher in protein (14.3 to 23.17%), ash

Comment [H51]: 'an increase'

Comment [H52]: 'a positive'

Comment [H53]: Rewrite as 'Overall Acceptability (OA)' score

Comment [H54]: delete

Comment [H55]: Also,

Comment [H56]: Bunkar *et al.* [49] Not included on the reference list

Comment [H57]: Millet-based

Comment [H58]: Replaced with 'at'

Comment [H59]: Delete since OA has been previously described

Comment [H60]: 'use'

(2.8 to 7.6%), and crude fiber (2.5 to 6.0%). The calcium content of formulated millet-based goods is similar to that of the market sample without fortification[52].

d) Fermented Milk-Millet Foods

Fermented foods are source of probiotics which is becoming popular in developing countries. Fermentation of grains is an old and low-cost technique of food preservation. One of the product developed was a fermented skim milk and pearl millet product. Skim milk and ungerminated pearl millet flour were used as raw materials, and curry patta(*Murrayakoenigii*), black pepper (*Piper nigrum*), cumin (*Cuminum cyminum*), and salt were used as flavouring agents. The milk was inoculated with 2% *S. thermophilus* and *L. rhamnosus*RSI3. The study revealed that the product has 5.01% protein, 0.82 % ash, 8.95 cfu/g LAB count, 0.73% fat and 82.18 % moisture. The product was similar to Rabadi in consistency[53].

Another author also developed a milk and millet based fermented drink using *S. thermophilus* C106 and *L. rhamnosus* GR-1. 1 litre of 3.25 % homogenized milk was mixed with hulled millet and heated at 85-90°C for 3.-60 minutes. Then the culture was added after cooling it at 40°C. Sensory tests revealed that millet fermented in milk was the most appealing. Viable counts of probiotic bacteria was more than 106 colony-forming units (CFU)/ml[54].Also, Millet milk curd was developedby germinating foxtail millet, little millet, kodo millet, proso millet, and barnyard millet. Germinated millets are used to extract milk and inoculated with NCDC 26. The curd like mass obtained was analysed and it was observed that acidity ranged from 0.74 to 1.2%, while the pH ranged from 3.5 to 4.5 [55]. Similarly a probiotic rich millet milk was developed using *Lactobacillus kefir*. Kefir millet fermented milk had much more protein, polyphenol, viable count, and acidity than the other commercially available comparable drinks [56].

6. CONCLUSION

Minor millets are not only cost effective but is excellent tool for value addition due to their nutraceutical properties. Minor millets are a field which is already getting attention due to their potential in decreasing, malnutritional as well as their bioactive profile. Kodo and little millet are one of the oldest grains and have been source of micro-nutrients in various traditional recipes. Also, with shortage of conventional cereal crops all around the world, millets can be an excellent replacement. Not only they are drought resistant, they have the ability to resist other climatic stress conditions. So, it is important to encourage millet production. Commercialization of millet-based product can attract consumers to make millet as part of their staple diet. Today's consumer is health conscious and demands health and tasty food. So, there is a great opportunity to undertake extensive research for developing higher quality millet goods that are healthy, delicious, have a long shelf life, are appealing in colour and appearance, as well as accessible to customers of all economic levels. Dairy products have wider acceptability among consumers, so it is an excellent tool of value addition. Millets can compensate for nutrients like iron which is deficient in milk.

Please check and correct your grammar and use proper punctuation. You may employ Grammarly to do most of these jobs for you. I have included some corrections in the

Comment [H61]: 'products'

Comment [H62]: 'millet-based'

Comment [H63]: Then, the....

Comment [H64]: 'were'

Comment [H65]: Curd-like

Comment [H66]: Similarly, a probiotic-rich millet...

Comment [H67]: Rewrite as 'Minor millets are not only cost-effective but are excellent tools for value addition due to their nutraceutical properties.'

Comment [H68]: 'that'

Comment [H69]: Rewrite as 'to decrease malnutrition...'

Comment [H70]: 'a shortage'

Comment [H71]: delete

Comment [H72]: Rewrite as 'Not only are they drought-resistant, but they also have the ability to resist other climatic stress conditions.'

Comment [H73]: 'products'

comment section, but it is still necessary to do a thorough job in order to have better communication. After making those corrections to your references, rearrange your in-text citation numbering. An example is shown in the comment section but not numbered accurately.

REFERENCES

- [1] Gupta A, Sood S, Agrawal PC, Bhatt JC. Floral Biology and Pollination System in Small Millets. Eur J Plant Sci Biotechnol. 2011;6:81-6
- [2] Shukla SK, Singh A, Kumar A, Singh A, Paul B, Sharma A, Pandey NK. Biomolecule Reports ISSN: 2456-8759.
- [3] Muragod PP, Muruli NV, Padeppagol S, Kattimani A. Physico-chemical properties and nutritional factors of kodo millet. Int J Pure App Biosci. 2019;7(1):117-23
- [4] Bunkar DS, Goyal SK, Meena KK, Kamalvanshi V. Nutritional, functional role of kodo millet and its processing: a review. International Journal of Current Microbiology and Applied Sciences. 2021;10(01):1972-85. DOI: <https://doi.org/10.20546/ijcmas.2021.1001.229>
- [5] Patel A, Parihar P, Dhumketi K. Nutritional evaluation of kodo millet and puffed kodo. International Journal of Chemical Studies. 2018;6(2):1639-42. <https://www.chemijournal.com/archives/2018/vol6issue2/PartX/6-1-88-649.pdf>
- [6] Dayakar Rao B, Bhaskarachary K, Arlene Christina GD, Sudha Devi G, Vilas AT, Tonapi A. Nutritional and health benefits of millets. ICAR_Indian Institute of Millets Research (IIMR) Rajendranagar, Hyderabad. 2017;2
- [7] Bunkar DS, Goyal SK, Meena KK, Kamalvanshi V. Nutritional, functional role of kodo millet and its processing: a review. International Journal of Current Microbiology and Applied Sciences. 2021;10(01):1972-85. DOI: <https://doi.org/10.20546/ijcmas.2021.1001.229>
- [8] Dey S, Saxena A, Kumar Y, Maity T, Tarafdar A. Understanding the antinutritional factors and bioactive compounds of kodo millet (*Paspalum scrobiculatum*) and little millet (*Panicum sumatrense*). Journal of Food Quality. 2022;2022(1):1578448. <https://doi.org/10.1155/2022/1578448>
- [9] Rabizadeh F, Mirian MS, Doosti R, Kiani-Anbouhi R, Eftekhari E. Phytochemical classification of medicinal plants used in the treatment of kidney disease based on traditional persian medicine. Evidence-Based Complementary and Alternative Medicine. 2022;2022(1):8022599. <https://doi.org/10.1155/2022/8022599>
- [10] Khare P, Maurya R, Bhatia R, Mangal P, Singh J, Podili K, Bishnoi M, Kondepudi KK. Polyphenol rich extracts of finger millet and kodo millet ameliorate high fat diet-induced metabolic alterations. Food & function. 2020;11(11):9833-47 <https://doi.org/10.1039/d0fo01643h>

Comment [H74]: There is no year in your reference

Comment [H75]: delete

Comment [H76]: Similar ref to [7]

Comment [H77]: delete

Comment [H78]: Repeated ref of [5]. Where is Bunkar et al. 2020?

- [11] Sharma S, Saxena DC, Riar CS. Using combined optimization, GC–MS and analytical technique to analyze the germination effect on phenolics, dietary fibers, minerals and GABA contents of Kodo millet (*Paspalum scrobiculatum*). *Food chemistry*. 2017 Oct 15; 233:20-8. <http://dx.doi.org/10.1016/j.foodchem.2017.04.099>
- [12] Emran TB, Islam F, Nath N, Sutradhar H, Das R, Mitra S, Alshahrani MM, Alhasaniah AH, Sharma R. Naringin and naringenin polyphenols in neurological diseases: understandings from a therapeutic viewpoint. *Life*. 2022 Dec 29;13(1):99. <https://doi.org/10.3390/life13010099>
- [13] Wang DM, Yang YJ, Zhang L, Zhang X, Guan FF, Zhang LF. Naringin enhances CaMKII activity and improves long-term memory in a mouse model of Alzheimer's disease. *International journal of molecular sciences*. 2013 Mar 11;14(3):5576-86. <https://doi.org/10.3390/ijms14035576>
- [14] Fan FY, Sang LX, Jiang M. Catechins and their therapeutic benefits to inflammatory bowel disease. *Molecules*. 2017 Mar 19;22(3):484. [doi:10.3390/molecules22030484](https://doi.org/10.3390/molecules22030484)
- [15] Srinivasan M, Sudheer AR, Menon VP. Ferulic acid: therapeutic potential through its antioxidant property. *Journal of clinical biochemistry and nutrition*. 2007;40(2):92-100. [doi: 10.3164/jcbn.40.92](https://doi.org/10.3164/jcbn.40.92)
- [16] Orfali R, Ratab ME, Hassan HM, Alonazi M, Gomaa MR, Mahrous N, GabAllah M, Kandeil A, Perveen S, Abdelmohsen UR, Sayed AM. Sinapic acid suppresses SARS CoV-2 replication by targeting its envelope protein. *Antibiotics*. 2021 Apr 11;10(4):420. [doi: 10.3390/antibiotics10040420](https://doi.org/10.3390/antibiotics10040420)
- [17] Chen C. Sinapic acid and its derivatives as medicine in oxidative stress-induced diseases and aging. *Oxidative medicine and cellular longevity*. 2016;2016(1):3571614. <https://doi.org/10.1155/2016/3571614>
- [18] Zhu H, Liang QH, Xiong XG, Wang Y, Zhang ZH, Sun MJ, Lu X, Wu D. Anti-inflammatory effects of p-coumaric acid, a natural compound of *Oldenlandia diffusa*, on arthritis model rats. *Evidence-Based Complementary and Alternative Medicine*. 2018;2018(1):5198594. <https://doi.org/10.1155/2018/5198594>
- [19] Han X, Guo J, You Y, Zhan J, Huang W. p-Coumaric acid prevents obesity via activating thermogenesis in brown adipose tissue mediated by mTORC1-RPS6. *The FASEB Journal*. 2020 Jun;34(6):7810-24. [DOI: 10.1096/fj.202000333R](https://doi.org/10.1096/fj.202000333R)
- [20] Tehami W, Nani A, Khan NA, Hichami A. New insights into the anticancer effects of p-coumaric acid: focus on colorectal cancer. *Dose-Response*. 2023 Jan 4;21(1):1559325822115070. [DOI: 10.1177/15593258221150704](https://doi.org/10.1177/15593258221150704)
- [21] Choi JM, Lee EO, Lee HJ, Kim KH, Ahn KS, Shim BS, Kim NI, Song MC, Baek NI, Kim SH. Identification of campesterol from *Chrysanthemum coronarium* L. and its antiangiogenic activities. *Phytotherapy Research*. 2007 Oct;21(10):954-9. [DOI: 10.1002/ptr.2189](https://doi.org/10.1002/ptr.2189)
- [22] Nandini C, Bhat S. Modified crossing (SMUASB) method for artificial hybridization in proso millet (*Panicum miliaceum* L.) and Little millet (*Panicum sumatrense*).

Comment [H79]: delete
Remove the month and day unless stipulated by the journal format. Do it for others

Comment [H80]: Follow the pattern in [11], [12], [13] to write the link for your DOI

Comment [H81]: Rewrite as suggested

Comment [H82]: Rewrite

Comment [H83]: rewrite

Comment [H84]: rewrite

Comment [H85]: rewrite

Electronic Journal of Plant Breeding. 2019;10(3):1161-70. DOI: 10.5958/0975-928X.2019.00147.9

Comment [H86]: rewrite

[23] Prabhakar CG, Ganiger BB, Bhat S, Nandini C, Tippeswamy VK, Manjunath HA. Improved production technology for little millet. Project Coordinating Unit ICAR-AICRP on Small Millets GKV, Bengaluru-560065. 2017:16.

[24] Kamatar MY, Hemalatha S, Meghana DR, Talawar S, Naik RK. Evaluation of Little Millet Landraces for Cooking and Nutritional Composition. Current Research in Biological and Pharmaceutical Sciences. 2013;2(1):7-10.

[25] Roopa U, Kasturiba B, Rama Naik RN, Usha Malagi UM, Shanthakumar G, Hemalatha S, Kiran Mirajkar KM. Physico-chemical and functional properties of little millet genotypes. Karnataka Journal of Agricultural Sciences, 2013;26(4), 539-542

[26] Kaur P, Purewal SS, Sandhu KS, Kaur M, Salar RK. Millets: A cereal grain with potent antioxidants and health benefits. Journal of Food Measurement and Characterization. 2019 Mar 15;13:793-806. <https://doi.org/10.1007/s11694-018-9992-0>

[27] Pradeep PM, Sreerama YN. Phenolic antioxidants of foxtail and little millet cultivars and their inhibitory effects on α -amylase and α -glucosidase activities. Food Chemistry. 2018 May 1;247:46-55. doi:10.1016/j.foodchem.2017.11

Comment [H87]: rewrite

[28] Shukla R, Pandey V, Vadnere GP, Lodhi S. Role of flavonoids in management of inflammatory disorders. In Bioactive food as dietary interventions for arthritis and related inflammatory diseases 2019 Jan 1 (pp. 293-322). Academic Press. doi:10.1016/b978-0-12-813820-5.0

Comment [H88]: rewrite

[29] Devi KP, Malar DS, Nabavi SF, Sureda A, Xiao J, Nabavi SM, Daglia M. Kaempferol and inflammation: From chemistry to medicine. Pharmacological research. 2015 Sep 1;99:1-0. doi:10.1016/j.phrs.2015.05.002

Comment [H89]: rewrite

[30] Lin Y, Shi R, Wang X, Shen HM. Luteolin, a flavonoid with potential for cancer prevention and therapy. Current cancer drug targets. 2008 Nov 1;8(7):634-46. doi:10.2174/156800908786241050

Comment [H90]: rewrite

[31] Sato S, Mukai Y. Modulation of chronic inflammation by quercetin: The beneficial effects on obesity. Journal of inflammation research. 2020 Aug 4:421-31. DOI: 10.2147/JIR.S228361

Comment [H91]: rewrite

[32] Wang H, Fu Y, Zhao Q, Hou D, Yang X, Bai S, Diao X, Xue Y, Shen Q. Effect of different processing methods on the millet polyphenols and their anti-diabetic potential. Frontiers in nutrition. 2022 Feb 11;9:780499. doi: 10.3389/fnut.2022.780499

Comment [H92]: rewrite

[33] Hassan S, Hussain MB, Waheed M, Ahmad K, Kassymov S, Shariati MA, Akram M, Mishra AP, Egbuna C. Effect of Germination Processing on Bioactive Compounds of Cereals and Legumes. Functional Foods and Nutraceuticals: Bioactive Components, Formulations and Innovations. 2020:283-306. DOI:10.1007/978-3-030-42319-3_16

Comment [H93]: rewrite

[34] Bhuvaneshwari G, Nirmalakumari A, Kalaiselvi S. Impact of soaking, sprouting on antioxidant and anti-nutritional factors in millet grains. *J Phytol.* 2020;12:62-6. doi: 10.25081/jp.2020.v12.6384

Comment [H94]: rewrite

[35] Guha M, Sreerama YN, Malleshi NG. Influence of processing on nutraceuticals of little millet (*Panicum sumatrense*). In *Processing and impact on active components in food 2015 Jan 1* (pp. 353-360). Academic Press. doi:10.1016/b978-0-12-404699-3

Comment [H95]: rewrite

[36] Devi S, Modgil R. Physico-chemical Characteristics, Functional Properties and Antinutritional Factors of Domestically Processed Sub Himalayan Non-conventional Millet koda (*Eleusine coracana*). *International Journal of Current Microbiology and Applied Sciences* (2020) 9. 2020(11):2808-17. <https://doi.org/10.20546/ijcmas.2020.911.340>

[37] Sharma S, Saxena DC, Riar CS. Using combined optimization, GC-MS and analytical technique to analyze the germination effect on phenolics, dietary fibers, minerals and GABA contents of Kodo millet (*Paspalum scrobiculatum*). *Food chemistry.* 2017 Oct 15;233:20-8. DOI: 10.1016/j.foodchem.2017.04.099

Comment [H96]: rewrite
Repeated ref [11]

[38] Singh N, David J, Thompkinson DK, Seelam BS, Rajput H, Morya S. Effect of roasting on functional and phytochemical constituents of finger millet (*Eleusine coracana* L.). *The Pharma Innovation Journal.* 2018;7(4):414-8.

[39] Pampangouda P, Munishamanna KB, Gurumurthy H. Effect of *Saccharomyces boulardii* and *Lactobacillus acidophilus* fermentation on little millet (*Panicum sumatrense*). *Journal of Applied and Natural Science.* 2015 Jun 1;7(1):260-4. DOI <https://doi.org/10.31018/jans.v7i1.599>

Comment [H97]: Delete

[40] Sharma S, Sharma N. Preparation of probiotic enriched functional beverage of Kodo millet (*Paspalum scrobiculatum*) a nutritionally enriched absolute new product for commercialization. *Journal of Pharmacognosy and Phytochemistry.* 2021;10(1):752-8. <https://www.phytojournal.com/archives/2021/vol10issue1/PartK/9-6-362-529.pdf>

[41] NAAS 2012. Integration of Millets in Fortified Foods. Policy Paper No. 54, National Academy of Agricultural Sciences, New Delhi. 2012;15 p. <https://naas.org.in/Policy%20Papers/policy%2054.pdf>

[42] Kumar A, Goel BK, Karthikeyan S, Asgar S, Gedda AK, Choudhary KK, Uprit S. Protein and calcium rich malted health drink power. *J. Food Process. Technol.* 2013;4(214):10-4172. <http://dx.doi.org/10.4172/2157-7110.1000214>

[43] Parvathi S, Nithya M, Yogeshwari R. Development of a novel health drink from millets. *Int J Home Sci Extn Comm Manage.* 2015;2(2):90-4. DOI : 10.15740/HAS/IJHSECM/2.2/90-94

Comment [H98]: rewrite

[44] Mathangi, & Geethanjali S. Preparation and analysis of value added health mix using germinated millets flour and corn silk (functional food) for diabetic patients. *International journal of advanced research in engineering technology & science.* (2016, december);3(12).

Comment [H99]: Rewrite in title case and italicized

Comment [H100]: Rewrite to align with others

- [45] Vanishree S, Kammar MR, Nidoni U. Development and evaluation of pearl millet based novel health drink. *Advances in Life Sciences*. 2016;5(13):5483-6. <https://doi.org/10.21048/ijnd.2016.53.4.8404>
- [46] Nishad PK, Maitra S, Nilima J. Physicochemical, functional and sensory properties of developed health drink from minor millets. *International Journal of Home Science*. 2017;3(2):503-6. <https://www.homesciencejournal.com/archives/2017/vol3issue2/ParH/3-2-82-113.pdf>
- [47] Bunkar DS, Jha A, Mahajan A. Optimization of the formulation and technology of pearl millet based 'ready-to-reconstitute' kheer mix powder. *Journal of food science and technology*. 2014 Oct;51:2404-14. DOI 10.1007/s13197-012-0800-2
- [48] Tumwine G, Atukwase A, Tumuhimbise GA, Tucungwirwe F, Linnemann A. Production of nutrient-enhanced millet-based composite flour using skimmed milk powder and vegetables. *Food Science & Nutrition*. 2019 Jan;7(1):22-34. <https://doi.org/10.1002/fsn3.777>
- [49] Bunkar DS, Bharti P, Meena KK, Goyal SK, Paswan VK. Studies on the optimization and development of functional instant kodomillet based porridge mix. *Int J Curr Microbiol Appl Sci*. 2020;9(9):1462-80. <https://doi.org/10.20546/ijcmas.2020.909.186>
- [50] Thathola A, Srivastava S. Physicochemical properties and nutritional traits of millet-based weaning food suitable for infants of the Kumaon hills, Northern India. *Asia Pacific journal of clinical nutrition*. 2002 Mar;11(1):28-32.
- [51] Balasubramanian S, Kaur J, Singh D. Optimization of weaning mix based on malted and extruded pearl millet and barley. *Journal of Food Science and Technology*. 2014 Apr;51:682-90. DOI 10.1007/s13197-011-0579-6
- [52] Prasanna MS, Sowjanya VS, Jaya E, Rajender G. Development of millet based instant weaning mix. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(4):1908-13
- [53] Basu S, Tomar SK. Development of novel indigenous pearl millet based fermented skim milk product. *International Journal of Fermented Foods*. 2016;5(1):39-46. <https://doi.org/10.5958/2321-712x.2016.00005.3>
- [54] Di Stefano E, White J, Seney S, Hekmat S, McDowell T, Sumarah M, Reid G. A novel millet-based probiotic fermented food for the developing world. *Nutrients*. 2017 May 22;9(5):529. <https://doi.org/10.3390/nu9050529>
- [55] Sheela P, moorthyUmaMaheswari T, Kanchana S, Kamalasundari S, Hemalatha G. Development and evaluation of fermented millet milk based curd. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(4):714-7

Comment [H101]: rewrite

Comment [H102]: Rewrite in title case and italicized

Comment [H103]: rewrite

[56]Chen S, Wang Y. Response surface optimization of millet milk fermented by Lactobacillus kefir. *Journal of Food Processing and Preservation*. 2022 Jul;46(7):e16675.<https://doi.org/10.1111/jfpp.16675>

Comment [H104]: Italicized

WEB SOURCES

<https://ipad.fas.usda.gov/countrysummary/Default.aspx?id=IN&crop=Millet>

<https://apeda.gov.in/milletportal/Production.html>

Comment [H105]: While separating this? Numbered it and show it in the in-text citation.

Note: All journal names must be in italics, e.g., *Nutrients*, *Journal of Pharmacognosy and Phytochemistry*, and *Journal of Food Processing and Preservation* for [54], [55], and [56], respectively. All in-text citations and references should be renumbered.

Comment [H106]:

UNDER PEER REVIEW