

Effects of *Eleophorbiadrupifera* leaves extract on some Biochemical and Haematological parameters of Albino rats

ABSTRACT

Water and ethanol-water extracts of *Eleophorbiadrupifera* leaves were administered orally in graded doses of 0.5ml (147.06mg/kg body weight), 1.0ml (303.3 mg/kg body weight) water extract and 0.5ml (157.66 mg/kg body weight), 1.0ml (301.72 mg/kg body weight) ethanol-water extract on experimental animals for 3 weeks. The effect of the extracts on some biochemical and haematological parameters were evaluated in albino rats. At the end of 21 days, the growth response was higher in the test groups than in the control ($p < 0.05$). The **Aspartate transaminase** (AST) and **Alanine transaminase** (ALT) levels of the test groups were significantly lower than the control ($p < 0.05$) and total bilirubin showed significantly high values in test groups when compared with the control ($p < 0.05$). The **White blood cell** (WBC) was significantly increased while the **Red blood cell** (RBC) was decreased when compared with the control ($p < 0.05$). There were no significant changes in **Packed cell volume** (PCV) and **haemoglobin** (Hb) of the experimental animals ($p < 0.05$). There was no significant difference between the effects of the water extract and that of the ethanol-water extract on the albino rats ($p < 0.05$). The results suggest that the leaf extract of *Eleophorbiadrupifera* may possess the capacity to stimulate appetite and there may be no adverse effect associated with the use of this extract in phytotherapy.

Keywords: *Eleophorbiadrupifera*, biochemical, haematological, albino rats.

INTRODUCTION

Medicinal plants are plants which have been claimed and confirmed to possess medicinal properties. They can also be referred to as herbal medicines (Okon, 2012). “The healing power of herbs had been recognized and botanic medicine is one of the oldest practiced professions by mankind. Most medicinal plants have not been thoroughly evaluated for their toxicity profiles though medicinal plants constituents mimic more closely the natural constitution of the human somatic system and following the lock and key hypothesis, it is expected that they will fit better into such system” (Samorinia, 2016). “Secondly, in terms of environmental friendliness, medicinal plants and their products are far more advantageous than orthodox medicines since they constitute a lesser form of pollution menace and are renewable” (Gamaniel, 2021, Akpabioet *al.*, 2012; Akpakpanet *al.*, 2012). “However, a blanket assumption should not be

made about the safety of medicinal plants and their products, since seemingly innocuous plants may turn out to be toxic. Though some of the medicinal plants have both medicinal and industrial applications”(Akpakpanet *al.*, 2020, Uwanta, 2024). “*Eleophorbiadrupifera* belongs to the family Euphorbiaceae which is one of the largest families of the angiospermae. It holds about 300 genera and over 600 species. *Eleophorbiadrupifera* looks like a tall Euphorbia with long fleshy leaves. When young, the branches are succulent and angular. Old trees have a clear trunk and densely crowded ascending branches with the leaves in the tufts at the end. Plants of the Euphorbiaceae are frequently used in the indigenous system of medicine. Little had earlier been known about the pharmacological actions of the specie *Eleophorbiadrupifera*, albeit it is listed among the plants that ‘heal’. The leaves, stem, bark, roots and latex are used depending on the type of ailment. Some herbal medicine practitioners posit that the leaves in particular are anti-hypertensive, antidiuretic, skeletal muscle relaxant and antidiabetic” (Comley, 2019). “The leaf is also used as filaricide and against guinea worm infestation. Apart from the use of *Eleophorbiadrupifera* for treatment of various ailments, traditional herbalists claim that it is also effective in aiding or inducing labour or abortion. The latex is used to remove warts though very injurious to the eyes” (Samorini, 2016). Given the traditional use of the plant in the management of various ailments, it becomes necessary to evaluate its toxicological potential in order to caution or encourage its use in traditional medicine.

MATERIALS AND METHODS

The leaves of *Eleophorbiadrupifeara* were collected from the botanical garden of the University of Calabar, Cross River State and were authenticated by a taxonomist in the department of Botany, University of Uyo, Uyo. The leaves of *Eleophorbiadrupifeara* were prepared by washing with distilled water and dried under shade. It was afterwards dried in the oven at 50-60⁰C for 2 days, ground into powder with an electric blender and stored in airtight containers for analytical use.

Extraction of sample

The ground sample (10g) was weighed out into a volumetric flask. Distilled water was added and made up to 100ml mark and the mixture vigorously agitated for 1 hour before it was filtered. The filtrate was stored for use as the water extract. About 10g of the blended leaves sample was again weighed out into a volumetric flask. 80% ethanol was added and made up to 100ml mark. The mixture was vigorously agitated and filtered. The filtrate obtained was poured into a graduated

beaker and evaporated on a hot plate at 65°C. On evaporation to the 20ml mark, the beaker was removed from the hot plate and distilled water was added to the 75ml mark, this was used as the ethanol-water extract.

Experimental Animals

Thirty (30) albino rats weighing 100-141g were obtained from the disease-free stock of the animal house, Department of Biochemistry, University of Calabar and used for the study. The 30 animals were housed in plastic cage with stainless still mesh at the bottom to ensure that faeces and feed droppings were inaccessible to the experimental animals. The environment was under standard conditions of the temperature (28±2°C) and relative humidity (46±5%) with a 12hours light-dark cycle and adequate ventilation. The animals were fed daily with commercial rat mash and water was given from bottles with stainless steel nozzles throughout the study.

Grouping and Treatment of animals

The animals were assigned into five groups of six rats each. Group 1 animals (control) were fed with the commercial rat mash only. Groups 2 and 3 (experimental group) were placed on 0.5ml and 1.0ml water extract respectively, Groups 4 and 5 (experimental group) were respectively treated with 0.5ml and 1.0ml ethanol-water extract. The extracts were daily administered to the rats orally for a period of 21 days.

Collection of Serum sample

Twenty four hours after the last administration, the animals were anaesthsized under chloroform and dissected. Blood was obtained through cardiac puncture into plain sample tubes and allowed to stand for 2 hours and thereafter centrifuged at 2000rpm for 10 minutes to separate serum from the blood cells. The blood serum obtained was used for assay of ALT, AST, total bilirubin, WBC, RBC, PCV and Hb.

Biochemical Assay

Serum and liver activities of ALT and AST were determined using the methods of Reitman and Frankel (2017). "Total bilirubin were estimated according to instructions from laboratory kits obtained from Randox Laboratory Ltd., United Kingdom and Absorbance were read using a UV/visible spectrophotometer –SP-300" (Jendrassik and Grof, 2021).

Haematological Assay

The WBC and RBC were counted by microscopic visual identification using Turk's fluid and Hayem's fluid respectively (Dacie and Lewis, 2015). The PCV was determined by capillary method using haematocrit centrifuge (Jones, 2015). Hb concentration was determined using Drabkin's reagent (Van Kampen and Zijistra, 2014)

Statistical Analysis

Data are presented as mean \pm SD and analyzed for statistical significance among group means by t-test. P-values less than 0.05 were considered statistically significant.

RESULTS and DISCUSSION

Table 1: Haematological and Biochemical Parameters of Rats fed Extracts of *Eleophorbiadrupifera*

Parameter/ group	WBC($\times 10^9$)	RBC($\times 10^9$)	PCV(%)	Hb(g/dI)	AST(U/L)	ALT(U/L)	Total Bilirubin (μ mol/l)
Group 1 (Control)	2.46 \pm 0.28	2.47 \pm 0.06	39.4 \pm 4.34	12.88 \pm 2.24	126.16 \pm 2.98	32.32 \pm 5.02	11.29 \pm 0.99
Group 2	3.15 \pm 0.34	2.33 \pm 0.06	39.2 \pm 9.15	11.82 \pm 2.09	68.5 \pm 16.15	18.00 \pm 2.94	16.19 \pm 1.48
Group 3	3.95 \pm 0.28	2.34 \pm 0.01	42.75 \pm 2.50	13.03 \pm 2.07	69.2 \pm 23.85	19.58 \pm 4.44	18.70 \pm 1.41
Group 4	3.84 \pm 0.23	2.29 \pm 0.05	40.75 \pm 3.77	12.35 \pm 1.08	59.38 \pm 8.26	28.83 \pm 3.57	16.19 \pm 0.61
Group 5	3.36 \pm 0.64	2.17 \pm 0.20	38.5 \pm 7.15	12.18 \pm 1.82	69.93 \pm 20.15	24.17 \pm 2.60	14.06 \pm 2.07

Results are presented as mean \pm SD

As shown in Table 1 above, the AST and ALT activities were found to be significantly lower in the treatment groups than in the control ($p < 0.05$). There was however no significant difference in effect on increasing dose of each extract and between the effects of the water extract and ethanol-water extract respectively ($p < 0.05$). The Total bilirubin of the treatment groups showed significant increase when compared with the control ($p < 0.05$). The treatment groups showed significant increase in WBC and decrease in RBC when compared with the control group

($p < 0.05$). There were no significant differences in PCV and Hb concentrations between the treatment groups and the control ($p < 0.05$).

Medicinal plants contribute immensely in the healthcare of populations in various parts of the world (FAO, 1999) example in Africa (Cunningham, 2003; Mashall, 2016), in China (He and Sheng, 2019), Europe (Gamaniel, 2021) and North America (Brevoort, 2015). “Many of our rural populations depend on a variety of plant materials for their wellbeing” (Akpakpan *et al.*, 2017). “These plants do not only supply essential nutrients but contain active substances which have proven efficacious against many diseases” (Gamaniel, 2021). This study examined the effects of *Eleophorbiadrupifera* on the blood biochemistry and haematology of rats.

It specifically examined the impact of treatment of rats with varying doses of *Eleophorbiadrupifera* on serum levels of AST, ALT and total bilirubin, WBC, RBC, PCV and Hb concentrations, in an attempt to evaluate the hepatotoxic potential of this plant extract. One of the organs usually affected by ingestion of xenobiotics is the liver. Generally, hepatic injury is often associated with alterations in the serum and liver levels of some enzymes notably ALT, AST and ALP (Whitby *et al.*, 2013) and studies with medicinal plant extracts have shown the varying effects of phytochemicals on serum and liver enzyme levels. While some phytochemicals are hepatotoxic, others are hepatoprotective. The significant decrease in both ALT and AST activities of the experimental animals albeit with an ALT/AST ratio of each group below one indicates that there was no liver damage. According to Strove and (Makarova, 2014), ALT/AST ratio greater than one are indicative of liver damage. The decrease in both ALT and AST activities might be due to active participation of the enzymes in tissue and cellular activities (Lehninger, 2002). The increase in total bilirubin though significant was still within normal range as Guyton (2016) posits that normal individuals have less than 20.56 $\mu\text{mol/L}$ of bilirubin circulating in their blood. The increase might have been due to destruction of some red blood cells as indicated by the decrease in RBC of the test animals. RBC though significantly decreased was still within normal range. According to Guyton (2016) normal RBC ranges from 4.7 to $5.2 \times 10^{12}/\text{L}$.

From the statistical comparisons, the effects of the leaf extract are not dose dependent. Some herbal medicine practitioners posit that leaves of *Eleophorbiadrupifera* in particular are anti-hypertensive, antidiuretic, skeletal muscle relaxant and antidiabetic. The leaf is also used as

filaricide and against guinea worm infestation (Comley, 2018). Apart from the use of *Eleophorbiadrupifera* for treatment of various ailments, traditional herbalists claim that it is also effective in aiding or inducing labour or abortion. The latex is used to remove warts though very injurious to the eyes (Samorini, 2016). Eno and Itam (2017) found out that there is a patent hypoglycemic agent(s) present in leaves of *Eleophorbiadrupifera*.

Conclusion

Eleophorbiadrupifera can thus be said to have little or no toxicological effects on biological systems and could be used internally for medicinal purposes without fear of hepatotoxicity.

Disclaimer (Artificial intelligence)

Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

- Akpabio, U. D., Wilson, L. A., Akpakpan, A. E. and I. B. Obot (2012). Phytochemical Screening and proximate composition of *Cassia hirsute* seeds. *Elixir Journal of Applied Chemistry* 8704-8707.
- Akpabio, U. D., Udo, U. E. and Akpakpan, A. E. (2012). Evaluation of Phytochemical, proximate and mineral element composition of stem of costus after (Bush cane). *Asian Journal of Plant Science and Research*, 2(5): 607-612.
- Akpakpan, A. E., Ukpong, E. J., Odiongenyi A. O. and Willie, I. E. (2017) Phytochemical Screening of the Ethanol and Aqueous Extracts of *Dicliptera verticillata* Leaves. *Archives of Current Research International*, 11(2): 1-5.
- Akpakpan, A. E. Uwanta, Emaime J. Akpan, Patrick E. and Owen, Aniekan E (2020). Phytochemical screening and analytical assessment of acid-base indicator properties of red and white kola nuts extracts. *Asian Journal of Advances in Research* 3(1): 1-7
- Brevoort, P. (2015). The booming U.S. botanical market: A new overview. *HerbalGram*, 44:33-43.
- Comely, J. C. M. (2019). New Micro filaricidal leaves from plants. *Trop. Med. Parasitol*, 43:1-9.
- Cunningham, A. B. (2003). African medicinal plants: Setting priorities at the interface between conservation and primary healthcare. UNESCO, Paris, France.
- Dacie, J. V. and Lewis, S.M. (2015). *Practical Haematology* 7th ed. Longman Group Ltd. U.K. p 37-63
- Eno, A.E. and Itam, E. H. (2017). Hypoglycemic agents in leaves of *Eleocharis acicularis*. *Phytotherapy Res.* 10:680-682.
- FAO (1999). State of the world's forests. *FAO*, Rome, Italy
- Gamaniel, K.S. (2021). Toxicity from medicinal plants and their products. *Nig. J. Nat. Prod. and Med.* 4:4-12.
- Guyton, A.C. (2016). *Textbook of Medical Physiology* (7th Ed.) W.B. Saunders Company, U.S.A p 42-53.
- He, S.A. and Sheng, N. (2019). Utilization and conservation of medicinal plants in China In: medicinal plants for forest conservation and healthcare. *FAO*, Rome. Italy
- Jendrassik, L. and Grof, P. (2021). Determination of Bilirubin. *Clin. Path.* 297:81.
- Lahniger, A. L. (2002). *Principles of Biochemistry*. Worth Publishers Inc. U.S.A. p 532-557
- Marshall, N. T. (2016). Searching for a cure: *Conservation of medicinal wildlife resources in East and South Africa*. TRAFFIC International, Cambridge, United Kingdom.
- Okon, A. J. (2012). Effect of *Eleocharis acicularis* on the blood biochemistry and haematology of albino rats. Unpublished BSc. Research, University of Uyo, Uyo.
- Reitman, S. and Frankel, S. (2017). A Colorimetric method for the determination of serum glutamic oxaloacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.* 28:56-63
- Samorini, G. (2016). Visionary Eye drops *Eleusis Periodical* 5:27-32.
- Stroev, E. and Makarova, V. (2014). *Laboratory manual for Biochemistry*. Mir. Publishers, Moscow, p 162-164

- Uwanta, E. J., Udo, I. E. Willie, I. N. and Akpakpan, A. E. (2024). Phytochemical screening, proximate and vitamin composition of Jack fruit (*Artocarpus Heterophyllus*) pulp and seed. *World Journal of Advanced Research and Reviews*, 22(01), 1500–1506
- Whitby, L. G., Percy-Robb, I.W. and Smith, A.F. (2013). Enzymes test in diagnosis. In: *Lecture notes on Clinical Chemistry*, 3rd ed. Blackwell Sci. Publication, London p 138-168.
- Van Kampen, E. J. and Zijistra, W.G. (2014). Spectrophotometry of haemoglobin and haemoglobin derivatives. *Advances in Clin. Chem.* 23:199-202.