

Molecular assessment of dominant genus in heavy metal contaminated soil from selected dumpsites in Port Harcourt, Nigeria

Abstract

Heavy metal contamination distorts and diminishes microbial diversity due to their inhibitory effects on bacteria, resulting in the elimination of susceptible bacteria and the proliferation of resistant strains, which eventually become dominant. Consequently, this study seeks to characterize the dominant bacteria (RCBBR_B37 and RCBBR_B38) isolated from soil at a dump site contaminated with heavy metals. DNA extraction from the isolates was performed using the Zymo Quick DNA Fungal/Bacterial Kit, and the purity of the extracted DNA was assessed using a NanoDrop 2000c spectrophotometer. Gel electrophoresis confirmed the presence of DNA bands, including a control band. Subsequently, PCR amplification using 27F and 1492R complementary primers was carried out on the extracted DNA. The PCR products underwent sequence analysis followed by BLAST search in the National Center for Biotechnology Information (NCBI) database. A phylogenetic tree was constructed based on the BLAST results. Isolate RCBBR_B37 exhibited a 98% similarity index to *Bacillusfungorum*, while RCBBR_B38 showed a 97% similarity index to *Bacillusparamycoides*. The prevailing presence of *Bacillus* in heavy metal contaminated dumpsite is a signature, representing the difference mechanisms of microbial interaction for heavy metals amelioration, including but not limited to bioaccumulation, sequestration, redox transformation, neutralization, metabolization, and detoxification.

Keywords: 27F/1492R primers; *Bacillus* spp.; Detoxification; Heavy metal contamination; Molecular analysis

1. Introduction

Dumpsites indeed represent substantial sources of heavy metal contamination (Abidemi and Theresa, 2015). When waste is deposited in dumpsites, diverse materials containing heavy metals such as lead, cadmium, mercury, arsenic, chromium, and others can seep into the soil and groundwater. Several factors contribute to the presence of heavy metals in dumpsites, including waste from industrial activities, electronic waste, household waste, and construction/demolition waste (Zargooshetal., 2013; Houessiononet al., 2021; Ishchenko and Vasykivskiy, 2020; Chen and Zhou, 2020). The existence of heavy metals in dumpsites presents notable environmental and health hazards. These metals can amass in the soil and water, impacting microbial diversity. Microorganisms exhibit various responses to heavy metal pollution, possessing mechanisms to either endure, accumulate, or detoxify heavy metals (Jamil et al., 2024; Roy et al., 2024). These reactions can carry significant implications for the environment, influencing nutrient cycling and the destiny of heavy metals in ecosystems.

Certain microorganisms possess inherent tolerance to specific heavy metals (Prabhakaran et al., 2016), enabling them to endure elevated metal concentrations in their surroundings without succumbing to toxicity. This tolerance may stem from genetic adaptations (Puente-Sanchez et al., 2018), the presence of particular cellular structures (Tamas et al., 2014), or the existence of enzymes that shield against metal toxicity (Bhaduri and Anwasha, 2012). Some microorganisms can accumulate heavy metals within their cells or on their surfaces (Gadd, 1990) through various mechanisms, including metal binding to cell wall components or sequestration within specialized compartments (Jogawatetal., 2021). By sequestering heavy metals, microorganisms can diminish metal availability in the surrounding environment, potentially mitigating their toxicity. Certain microorganisms can catalyze chemical transformations of heavy metals via redox reactions. For instance, specific bacteria can convert toxic metal forms into less harmful or mobile forms through reduction, oxidation, or methylation processes (Raab and Feldmann, 2003), thereby affecting metal mobility and bioavailability in the environment. Microorganisms possess detoxification pathways that enable them to metabolize or neutralize heavy metals (Alotaibi et al., 2021), often involving the production of metal-binding proteins or peptides and the synthesis of enzymes facilitating the conversion of toxic metals into less harmful forms. Examples include metallothioneins, which bind metals, and metal-reducing enzymes such as metallothionein, glutathione, and phytochelatin (Khandaker et al., 2021; Juwarkaetal., 2010). Moreover, heavy metal pollution can alter the composition and dynamics of microbial communities in affected environments. Some microorganisms may thrive in metal-contaminated habitats due to their

ability to tolerate or utilize metals, while others may be inhibited or displaced (Shukla *et al.*, 2019). Bacteria, in particular, have been extensively studied for their responses to heavy metal contamination.

Bacteria demonstrate diverse reactions to heavy metal contamination, capitalizing on their varied metabolic capacities and genetic flexibility. They can harbor genes responsible for metal resistance mechanisms, often facilitated by synergistic interactions (Xin *et al.*, 2023), which may be situated on bacterial chromosomes, plasmids, or transposons. These genes encode proteins like metal efflux pumps, metal-binding proteins, and metal-detoxifying enzymes (Mathivanan *et al.*, 2021), aiding bacteria in their survival within metal-laden environments. Additional pathways for responding to heavy metal exposure include expelling heavy metals from their cells, sequestering them, undergoing redox reactions, forming biofilms as protective structures, adapting genetically to prolonged exposure, and engaging in synergistic interactions to enhance survival (Pal *et al.*, 2022; Roy *et al.*, 2023; Mohapatra *et al.*, 2020). Commonly implicated bacteria in one or more of these responses include *Cupriavidusmetallidurans*, *acidophilic* bacteria, *Stenotrophomonasmaltophilia*, various *Arthrobacter*, *Pseudomonas*, *Corynebacterium*, and *Bacillus* species (Aljerfetal., 2016). These bacteria, among others, play vital roles in microbial communities thriving in metal-contaminated environments.

This study was designed to characterize the dominant bacterial isolates retrieved from specific dump sites in Port Harcourt using a molecular approach. The findings reveal that *Bacillus* species dominate the dump sites under investigation.

2. Methodology

2.1 Subculturing of dominant bacteria

Bacterial isolates previously obtained from dumpsites contaminated with heavy metals in Port Harcourt were subcultured using the spread plate method. Nutrient Agar was inoculated in duplicate with 0.1 ml of a 10⁻³ dilution using a sterile glass spreader. All plates were properly labeled, inverted, and then incubated at 37°C for 24 hours. Morphological features were utilized to identify the most prevalent bacterial isolate that also spread across the cultured plates examined.

2.2 Molecular Characterization of Isolates

The DNA extraction process utilized the Zymo Quick DNA Fungal/Bacterial Kit, following the kit's protocol with some adjustments. Initially, the bacterial culture was collected and transferred into a sterilized mortar. Next, 750 µl of bashing bead buffer was added to the sample and homogenized. The resulting homogenate was then transferred to a bashing bead tube and further homogenized using a benchtop vortex for 20 minutes. Subsequently, the bashing bead tube containing the homogenized sample underwent centrifugation in a refrigerated centrifuge at ≥10,000 x g for 1 minute. Following centrifugation, 400 µl of supernatant was transferred to a Zymo-Spin III-F Filter in a Collection tube and centrifuged at 8,000 x g for 1 minute, after which the Zymo-Spin III-F Filter was discarded. To the filtrate in the Collection Tube, 1,200 µl of Genomic Lysis Buffer was added and thoroughly mixed. From this mixture, 800 µl was transferred to a Zymo-Spin II-C Column in a Collection Tube and centrifuged at 10,000 x g for 1 minute. The flow-through from the Collection Tube was then discarded, and this step was repeated. Next, 200 µl of DNA Pre-Wash Buffer was added to the Zymo-Spin II-C Column and centrifuged at 10,000 x g for 1 minute. Following this, 500 µl of g-DNA Wash Buffer was added to the Zymo-Spin II-C Column and centrifuged at 10,000 x g for 1 minute. The Zymo-Spin II-C Column was subsequently transferred to a clean 1.5 ml micro-centrifuge tube, and 100 µl (with a minimum of 35 µl for higher output) of DNA Elution Buffer was directly added to the column matrix. Finally, the tube was centrifuged at 10,000 x g for 30 seconds to elute the DNA.

2.3 Determination of DNA Concentration and Purity Using Nanodrop

The DNA concentration and purity were assessed using a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific, USA). Purity was determined by calculating the ratio of ultraviolet (UV) light absorbance at 260 nm to that at 280 nm. The NanoDrop was connected to a computer system, and the sensor was cleaned using cotton wool soaked in 70% ethanol. To blank the system, one microliter (1 µl) of Elution buffer (the solution used to re-suspend the DNA) was dispensed directly into the NanoDrop sensor. Following this, DNA samples (1 µL each) were

individually loaded onto the sensor according to Bunu et al. (2020). Prior to loading a new sample, the sensor was routinely wiped to prevent contamination. Duplicate measurements were taken for each sample using the NanoDrop.

2.4 Gel Electrophoresis

Gel electrophoresis was conducted using a 1.5% agarose gel prepared by dissolving 0.75g of agarose powder in 50 ml of 1X Tris-Boric acid-EDTA (TBE) buffer in a measuring flask, followed by microwaving for 2 minutes until a clear solution was obtained (Bhilocha*et al.*, 2011). Subsequently, 5 μ L of EZ viewing dye (Blue Light) was added to the solution in a conical flask and then poured into the casting tray or gel holder. The comb was positioned within the casting tray, and the mixture was allowed to solidify at room temperature for 20 to 30 minutes. Following solidification, the gel electrophoresis unit was set up by placing the gel holder containing the gel onto the platform inside the gel tank, and 1X TBE buffer was poured into the tank until the gel was fully submerged. A molecular weight marker (1Kb DNA Ladder) was loaded into the first lane, while DNA samples were individually loaded into the wells created by the comb on the gel. Each DNA sample (5 μ L) was mixed with 1 μ L of 6X loading dye before loading onto the gel. Additionally, a control containing all components of the PCR reaction mixture except the template DNA was loaded. The electrophoresis apparatus was operated at 100 volts for 40 minutes. Following the completion of the run, DNA fragments were visualized using an ultraviolet (UV) transilluminator (Gel Documentation microDOCTM, Cleaver Scientific Ltd, UK).

2.5 Polymerase Chain Reaction (PCR)

The target region was amplified using OneTaq® Quick-Load® 2X Master Mix (New England Biolabs, USA, Catalogue No. M0486) in an Eppendorf Mastercycler (Nexus Gradient 230, Germany), following the conditions outlined in Table 1. The primer sets utilized consisted of the forward primer 27F-AGAGTTTGATCMTGGCTCAG and the reverse primer 1492R TACGGYTACCTTGTTACGACTT. Thermal cycling conditions were as follows: initial denaturation at 94°C for 5 minutes; followed by 35 cycles comprising denaturation at 94°C for 30 seconds, annealing at 50°C for 60 seconds, and extension at 72°C for 90 seconds; with a final extension at 72°C for 10 minutes. The thermal cycler was set to maintain the PCR mix at 4°C after completion. PCR products underwent purification using an enzymatic method (ExoSAP) as follows: An ExoSAP master mix was prepared by combining 50 μ l of 20 U/ μ l Exonuclease I (Catalogue No. NEB M0293L) and 200 μ l of 1 U/ μ l Shrimp Alkaline Phosphatase (Catalogue No. NEB M0371) in a 0.6 ml micro-centrifuge tube. The reaction mixture was then prepared by mixing 10 μ l of the amplified PCR product with 2.5 μ l of the ExoSAP mix from step 1. The resulting mix was incubated at 37°C for 15 minutes followed by 80°C for 15 minutes (Frank *et al.*, 2008).

Table 1: Polymerase chain reaction cocktail components

Component	Volume of a 12.5 μ L reaction
Template DNA	2 μ L
10 μ M Forward Primer	0.25 μ L (10nM)
10 μ M Reverse Primer	0.25 μ L (10nM)
One Taq Quick Load 2X Master Mix with Standard Buffer	6.25 μ L
Nuclease free water	3.75 μ L

2.6 Sequence Analysis and blasting

The PCR products underwent sequencing utilizing the Nimagen Brilliant Dye™ Terminator Cycle Sequencing Kit V3.1, BRD3-100/1000, following the manufacturer's instructions as outlined at: [Nimagen Sequencing Kit Instructions](#). Subsequently, the labeled products were purified using the ZR-96 DNA Sequencing Clean-up Kit (Catalogue No. D4053), as detailed in the instructions available at: [Zymo Research DNA Sequencing Clean-up Kit Instructions](#). The purified products were then injected into the Applied Biosystems ABI 3500XL Genetic Analyzer

with a 50cm array, employing POP7 polymer, accessible at: [Applied Biosystems ABI 3500XL Genetic Analyzer with POP7](#), and sequence data were acquired. The sequences obtained from the sequencer underwent base calling using ChromasLite. Subsequent sequence editing was performed using BioEdit software prior to conducting a Basic Local Alignment Search Tool (BLAST) search within the NCBI (National Center for Biotechnology Information) database, available at: [NCBI BLAST](#). Similar sequences retrieved from the search were downloaded and aligned using ClustalW.

2.7 Phylogenetic Tree

The evolutionary lineage was deduced utilizing the Neighbor-Joining technique (Saitou and Nei, 1987), with the optimal tree depicted. The percentage of replicate trees wherein the associated taxa formed clusters in the bootstrap test (500 replicates) is displayed adjacent to the branches (Felsenstein, 1992). The tree is drawn proportionally, with branch lengths reflecting the evolutionary distances used in constructing the phylogenetic tree. These distances were calculated using the Jukes-Cantor method (Jukes and Cantor, 1969) and are expressed in units of base substitutions per site. The evolutionary analysis was carried out using MEGA11 (Tamura et al., 2021).

3. Results

Molecular Characterization of Bacterial Isolates

Table 2 presents the molecular characterization of bacterial isolates, with Fig. 1 showcasing the gDNA bands and Fig. 2 depicting the 16S rRNA PCR bands of the isolates. The results of the blasting are subsequently presented in Table 3 (Fasta sequences) and used in phylogenetic tree reconstruction (Fig. 3), while the outcomes of the similarity assignment check are presented in Table 4.

Table 2. NanoDrop spectrometry characteristics of the isolates

S/N	Sample	Conc. (ng/mL)	A260	A280	Purity (A260/A280)
1	RCBBR_37a	200.5	4.010	2.160	1.86
2	RCBBR_37b	208.5	4.170	2.239	1.86
3	RCBBR_38a	184.0	3.681	1.987	1.85
4	RCBBR_38b	184.2	3.685	1.989	1.85




Fig. 1. gDNA bands from the isolates (Lane 1: RCBBR_37; Lane 2: RCBBR_38)






Fig. 2. 16S rRNA PCR bands from the isolates (1: RCBBR_37; Lane 2: RCBBR_38)

Table 3. The sequencing reads of RCBBR_B37 and RCBBR_B38 isolates

Isolate code	Affiliated sequence
RCBBR_37	<p>>RCBBR_37 GCTTAATGCGTAACTKCAGCACTAAAGGGCGGAAACCCCTAACACTTAGCACTCATCG TTACGGCGTGGACTACCAGGGTATCTAATCCTGTTTGCTCCCCACGCTTTCGCGCCTCA GTGTCAGTTACAGACCAGAAAGTCGCCTTCGCCACTGGTGTTCCTCCATATCTCTACGCA TTCACCCGCTACACATGGAATCCACTTTCCTCTTCTGCACTCAAGTCTCCAGTTTCCA ATGACCCTCCACGGTTGAGCCGTGGGCTTTCACATCAGACTTAARAAACCACCTGCGCGC GCTTACGCCCAATAATTCCGGATAACGCTTGCCACCTACGTATTACCGCGGCTGCTGGC ACGTAGTTAGCCGKGGCTTCTGGTTAGGTACCGTCAAGGTGCCAGCTTATTCAACTAGC ACTTGTTCCTCCCTAACAACARAGTTTTACRACCCGAAAGCCTTCATCACTCACGCGGCG TTGYTCCGTCAGAYTTTCGTCCATTGCGGAARATTCCCTACTGCTGCCTCCCGTAGGAGT CTGGGCCGTGTCTCAGTCCCAGKGGCCGATACCCCTCTCAGGTCCGCTACGCATCGTT GCCTTGGTGAGCCGTTACCTACCAACTAGCTAATGCGACGCGGGYCCATCCATAAGTGA CAGCCGAAGCCGCTTTCATTTTGAACCATGCRGTTCAAATGTTATCCGGTATTAGCC CCGGTTTCCCGGAGTTATCCARTCTTATGGGCAGGTTACCCACGKGTWACTCACCCGTC CGCCGCTAACTTCATARGAGCAAGCTCTTAATCCATTGCTCGACTTGCATGTATTAGG</p>
RCBBR_38	<p>>RCBBR_38 AATGCGTAACTTCAGCACTAAAGGGCGGAAACCCCTAACACTTAGCACTCATCGTTTA CGGCGTGGACTACCAGGGTATCTAATCCTGTTTGCTCCCCACGCTTTCGCGCCTCAGTGT CAGTTACAGACCAGAAAGTCGCCTTCGCCACTGGTGTTCCTCCATATCTCTACGCATTTC ACCGCTACACATGGAATCCACTTTCCTCTTCTGCACTCAAGTCTCCAGTTTCCAATGA CCCTCCACGGTTGAGCCGTGGGCTTTCACATCAGACTTAAGAAACCACCTGCGCGCGCTT TACGCCCAATAATTCCGGATAACGCTTGCCACCTACGTATTACCGCGGCTGCTGGCACGT AGTTAGCCGTGGCTTCTGGTTAGGTACCGTCAAGGTGCCAGCTTATTCAACTAGCACTT GTTCTTCCCTAACAACAGAGTTTTACGACCCGAAAGCCTTCATCACTCACGCGGCGTTGC TCCGTCAGACTTTCGTCCATTGCGGAAGATTCCCTACTGCTGCCTCCCGTAGGAGTCTGG GCCGTGTCTCAGTCCCAGTGTGGCCGATACCCCTCTCAGGTCCGCTACGCATCGTTGCCT TGGTGAGCCGTTACCTACCAACTAGCTAATGCGACGCGGGTCCATCCATAAGTGACAGC CGAAGCCGCTTTCATTTTGAACCATGCRGTTCAAATGTTATCCGGTATTAGCCCCGG TTTCCCGGAGTTATCCAGTCTTATGGGCAGGTTACCCACGTGTTACTCACCCGTCCGCC GCTAACTTCATAAGAGCAAGCTCTTAATCCATTGCTCGACTTGCATGTATTAGGCACGC CGCCAGCGTTCATCCTGAGC</p>



Isolate 1: RCBBR_B37



Isolate 2: RCBBR_B38

Fig. 3. Phylogenetic tree of the identified bacterial isolates using neighbor joining tree

Table 4. Identity of the bacterial isolates - strain, name of organism, closest GenBank, percentage identity and accession

S/N	Strain	Organism	Closest GenBank Match	Similarity (%)	Accession No
1	RCBBR_37	<i>Bacillus fungorum</i>	<i>Bacillus fungorum</i> strain 17-SMS-01	98.09	OR105047
2	RCBBR_38	<i>Bacillus paramycoides</i>	<i>Bacillus paramycoides</i> strain NH24A2(T)	99.88	OR105048

4.0 Discussions

This investigation focused on the molecular characterization of prevalent bacteria within a heavy metal-contaminated dump site. Molecular characterization methods provide distinct advantages over cultural approaches for studying microbial communities. While cultural methods are restricted to identifying only cultivable microbes, molecular techniques enable the thorough examination of both cultivable and non-cultivable microorganisms present in environmental samples. Techniques such as polymerase chain reaction (PCR) and high-throughput sequencing offer precise and sensitive evaluations of microbial diversity, allowing for unbiased detection of various taxa and the identification of rare and novel microbes that may play significant ecological roles (Bella *et al.*, 2013). Moreover, molecular methods facilitate the quantitative analysis of microbial communities, empowering researchers to assess changes in community structure, relative abundance, and gene expression levels over time or in response to environmental changes. When combined with functional assays, molecular techniques also permit the exploration of microbial metabolic capabilities and activities, offering insights into the functional potential of microbial communities. In summary, molecular characterization methods offer robust and comprehensive tools for comprehending microbial diversity, ecology, and function in natural environments, as well as providing more precise characterization of microorganisms.

Culture-dependent molecular characterization has been utilized to identify a significant number of bacteria present in heavy metal-polluted environments. Islam and Sar (2011) employed this dual technique to investigate the bacterial community within uranium ore. Heavy metal-contaminated sites often harbor a diverse array of bacteria capable of surviving and thriving in these challenging conditions (Barman *et al.*, 2020; Saha *et al.*, 2022; Hao *et al.*, 2021). Among the common bacteria found in such environments are *Pseudomonas* spp., known for their versatility and ability to tolerate and detoxify metals through efflux pumps and metal-binding proteins, and *Bacillus* spp., which form resilient endospores and exhibit resistance to heavy metals. Alhasawietal. (2015) identified *P. fluorescens* as capable of tolerating and detoxifying heavy metals in contaminated media. Previously, Aguilar-Barajas *et al.* (2010) reported on the role of *Pseudomonas* spp. in metal homeostasis and detoxification. *Acidithiobacillus* spp. are excellent in bioleaching metals from ores and can tolerate extreme pH and metal concentrations. Akinci *et al.* (2011) confirmed the role of *Acidithiobacillus* spp. in heavy metal leaching in contaminated sediment, both in pure and mixed states. Alviz-Gazitu *et al.* (2019) reported *Cupriavidus metallidurans* as a model for studying extreme resistance to heavy metals such as cadmium, zinc, and cobalt, while *Arthrobacter* spp. and *Stenotrophomonas maltophilia* possess mechanisms for detoxifying and tolerating heavy metal exposure (Pires 2010; Mohapatra *et al.*, 2017). *Rhodococcus* spp. also contribute, often playing roles in bioremediation efforts due to their metabolic versatility and ability to withstand metal toxicity (Nazari *et al.*, 2022). *Bacillus* spp. form resilient endospores and exhibit resistance to heavy metals (Fakhar *et al.* 2022). Collectively, these bacteria play critical roles in the ecology and biogeochemistry of heavy metal-contaminated environments, influencing metal cycling, remediation processes, and ecosystem dynamics.

Bacillus spp. demonstrate significant adaptive responses to heavy metal exposure, rendering them crucial contributors to mitigating metal pollution in contaminated environments. These bacteria utilize various mechanisms to cope with metal toxicity, including metal sequestration through cell surface structures and exopolysaccharides

(Gupta *et al.*, 2014), enzymatic detoxification involving metallothioneins and metalloenzymes (Pandey *et al.*, 2013), and active efflux pump systems that expel metals from the cytoplasm (Fakhar *et al.*, 2022). Additionally, *Bacillus* species are proficient biofilm formers (Syed *et al.*, 2021), which not only shield cells from metal toxicity but also facilitate interactions with metals, such as sequestration and reduction. Maity *et al.* (2023) isolated *Bacillus* strain GH-s29, exhibiting outstanding biofilm formation linked to multi-metal remediation potential. Through genetic adaptation, *Bacillus* spp. can acquire or upregulate genes encoding metal resistance determinants, augmenting their capacity to survive and thrive in metal-contaminated habitats (Moore *et al.*, 2005; Shin *et al.*, 2012; Nanda *et al.*, 2019; Sharma and Shukla, 2021). Their metabolic versatility and resilience position *Bacillus* species as promising candidates for bioremediation applications, wherein they can be employed to enhance the removal or immobilization of heavy metals from contaminated sites, thereby contributing to environmental remediation endeavors

5.0 Conclusion

This research underscores the crucial role of molecular characterization techniques in elucidating microbial dominance within heavy metal-contaminated environments. Molecular methods offer distinct advantages over cultural characterization approaches, allowing for the comprehensive analysis of both cultured microbial taxa present in these challenging habitats. By utilizing techniques such as PCR, high-throughput sequencing, and functional assays, researchers can glean insights into microbial diversity, ecology, and functional potential in natural settings. The study also sheds light on the prevalence and adaptive mechanisms of bacteria, particularly *Bacillus* spp., in response to heavy metal exposure. It provides evidence of *Bacillus* species employing versatile strategies to cope with metal toxicity, including metal sequestration, enzymatic detoxification, and active efflux, thereby making them indispensable contributors to the mitigation of metal pollution in contaminated sites. The resilience and adaptability of *Bacillus* spp. position them as promising candidates for bioremediation applications, offering optimism for the restoration of heavy metal-contaminated environments and the preservation of ecosystem health.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

Abidemi, O. O., & Theresa, O. C. (2015). Environmental fate of heavy metals in soil of Ido-Osun waste dump site, Osogbo, Osun, Nigeria. *American Journal of Environmental Protection*, 3(1), 1-4.

Aguiar-Barajas, E., Ramírez-Díaz, M. I., Riveros-Rosas, H., & Cervantes, C. (2010). Heavy metal resistance in pseudomonads. *Pseudomonas: volume 6: molecular microbiology, infection and biodiversity*, 255-282.

Akinci, G., & Guven, D. E. (2011). Bioleaching of heavy metals contaminated sediment by pure and mixed cultures of *Acidithiobacillus* spp. *Desalination*, 268(1-3), 221-226.

Alhasawi, A., Costanzi, J., Auger, C., Appanna, N. D., & Appanna, V. D. (2015). Metabolic reconfigurations aimed at the detoxification of a multi-metal stress in *Pseudomonas fluorescens*: Implications for the bioremediation of metal pollutants. *Journal of Biotechnology*, 200, 38-43.

Aljerf, L., & AlMasri, N. (2018). A gateway to metal resistance: bacterial response to heavy metal toxicity in the biological environment. *Ann. Adv. Chem*, 2(1), 32-44.

Alotaibi, B. S., Khan, M., & Shamim, S. (2021). Unraveling the underlying heavy metal detoxification mechanisms of *Bacillus* species. *Microorganisms*, 9(8), 1628.

Alviz-Gazitua, P., Fuentes-Alburquenque, S., Rojas, L. A., Turner, R. J., Guiliani, N., & Seeger, M. (2019). The response of *Cupriavidus metallidurans* CH34 to cadmium involves inhibition of the initiation of biofilm formation, decrease in intracellular c-di-GMP levels, and a novel metal regulated phosphodiesterase. *Frontiers in Microbiology*, 10, 434607.

Ayangbenro, A. S., & Babalola, O. O. (2020). Genomic analysis of *Bacillus cereus* NWUAB01 and its heavy metal removal from polluted soil. *Scientific reports*, 10(1), 19660.

Barman, D., Jha, D. K., & Bhattacharjee, K. (2020). Metallotolerant bacteria: insights into bacteria thriving in metal-contaminated areas. *Microbial Versatility in Varied Environments: Microbes in Sensitive Environments*, 135-164.

Bhaduri, A. M., & Fulekar, M. H. (2012). Antioxidant enzyme responses of plants to heavy metal stress. *Reviews in Environmental Science and Bio/Technology*, 11, 55-69.

Bhilocha, S., Amin, R., Pandya, M., Yuan, H., Tank, M., LoBello, J., ... & Cowman, M. K. (2011). Agarose and polyacrylamide gel electrophoresis methods for molecular mass analysis of 5-to 500-kDa hyaluronan. *Analytical biochemistry*, 417(1), 41-49.

Bunu, S. J., Otele, D., Alade, T., & Dodoru, R. T. (2020). Determination of serum DNA purity among patients undergoing antiretroviral therapy using NanoDrop-1000 spectrophotometer and polymerase chain reaction. *Biomedical and Biotechnology Research Journal (BBRJ)*, 4(3), 214-219.

Chen, Y., & Zhou, Y. (2020). The contents and release behavior of heavy metals in construction and demolition waste used in freeway construction. *Environmental Science and Pollution Research*, 27, 1078-1086.

Di Bella, J. M., Bao, Y., Gloor, G. B., Burton, J. P., & Reid, G. (2013). High throughput sequencing methods and analysis for microbiome research. *Journal of microbiological methods*, 95(3), 401-414.

Fakhar, A., Gul, B., Gurmani, A. R., Khan, S. M., Ali, S., Sultan, T., ... & Rizwan, M. (2022). Heavy metal remediation and resistance mechanism of *Aeromonas*, *Bacillus*, and *Pseudomonas*: A review. *Critical Reviews in Environmental Science and Technology*, 52(11), 1868-1914.

Felsenstein, J. (1992). Estimating effective population size from samples of sequences: a bootstrap Monte Carlo integration method. *Genetics Research*, 60(3), 209-220.

- Frank, J. A., Reich, C. I., Sharma, S., Weisbaum, J. S., Wilson, B. A., & Olsen, G. J. (2008). Critical evaluation of two primers commonly used for amplification of bacterial 16S rRNA genes. *Applied and environmental microbiology*, 74(8), 2461-2470.
- Gadd, G. M. (1990). Heavy metal accumulation by bacteria and other microorganisms. *Experientia*, 46(8), 834-840.
- Gupta, S., Goyal, R., & Prakash, N. T. (2014). Biosequestration of lead using Bacillus strains isolated from seleniferous soils and sediments of Punjab. *Environmental Science and Pollution Research*, 21, 10186-10193.
- Hao, X., Zhu, J., Rensing, C., Liu, Y., Gao, S., Chen, W., ... & Liu, Y. R. (2021). Recent advances in exploring the heavy metal (loid) resistant microbiome. *Computational and structural biotechnology journal*, 19, 94-109.
- Houessionon, M. K., Ouendo, E. M. D., Bouland, C., Takyi, S. A., Kedote, N. M., Fayomi, B., ... & Basu, N. (2021). Environmental heavy metal contamination from Electronic Waste (e-waste) recycling activities worldwide: A systematic review from 2005 to 2017. *International journal of environmental research and public health*, 18(7), 3517.
- Ishchenko, V., & Vasylykivskyi, I. (2020). Environmental pollution with heavy metals: case study of the household waste. *Sustainable production: Novel trends in energy, environment and material systems*, 161-175.
- Islam, E., & Sar, P. (2011). Culture-independent and independent molecular analysis of the bacterial community within uranium ore. *Journal of Basic Microbiology*, 51(4), 372-384.
- Jamil, M., Malook, I., Rehman, S. U., Aslam, M. M., Fayyaz, M., Shah, G., ... & Abeed, A. H. (2024). Inoculation of heavy metal resistant bacteria alleviated heavy metal-induced oxidative stress biomarkers in spinach (*Spinacia oleracea* L.). *BMC Plant Biology*, 24(1), 221.
- Jogawat, A., Yadav, B., Chhaya, & Narayan, O. P. (2021). Metal transporters in organelles and their roles in heavy metal transportation and sequestration mechanisms in plants. *Physiologia Plantarum*, 173(1), 259-275.
- Jukes, T. H., & Cantor, C. R. (1969). Evolution of protein molecules. *Mammalian protein metabolism*, 3(24), 21-132.
- Juwarkar, A. A., & Yadav, S. K. (2010). Bioaccumulation and biotransformation of heavy metals. In *Bioremediation technology: Recent advances* (pp. 266-284). Dordrecht: Springer Netherlands.
- Khandaker, M. M., Abdullahi, U. A., Elyni, N., & Alias, N. (2021). Biological processes of heavy metals-contaminated environmental remediation: a review. *Journal of Environmental Treatment Techniques*, 9(3), 601-608.
- Maity, S., Sarkar, D., Poddar, K., Patil, P., & Sarkar, A. (2023). Biofilm-mediated heavy metal removal from aqueous system by multi-metal-resistant bacterial strain Bacillus sp. GH-s29. *Applied Biochemistry and Biotechnology*, 195(8), 4832-4850.
- Mathivanan, K., Chandirika, J. U., Vinothkanna, A., Yin, H., Liu, X., & Meng, D. (2021). Bacterial adaptive strategies to cope with metal toxicity in the contaminated environment—A review. *Ecotoxicology and Environmental Safety*, 226, 112863.
- Mohapatra, R. K., Behera, S. S., Patra, J. K., Thatoi, H., & Parhi, P. K. (2020). Potential application of bacterial biofilm for bioremediation of toxic heavy metals and dye-contaminated environments. In *New and future developments in microbial biotechnology and bioengineering: Microbial biofilms* (pp. 267-281). Elsevier.
- Mohapatra, R. K., Parhi, P. K., Patra, J. K., Panda, C. R., & Thatoi, H. N. (2017). Biotransformation of toxic heavy metals by marine metal resistant bacteria—a novel approach for bioremediation of the polluted saline environment. *Microbial Biotechnology: Volume 1. Applications in Agriculture and Environment*, 343-376.

- Moore, C. M., Gaballa, A., Hui, M., Ye, R. W., & Helmann, J. D. (2005). Genetic and physiological responses of *Bacillus subtilis* to metal ion stress. *Molecular microbiology*, 57(1), 27-40.
- Nanda, M., Kumar, V., & Sharma, D. K. (2019). Multimetal tolerance mechanisms in bacteria: The resistance strategies acquired by bacteria that can be exploited to 'clean-up' heavy metal contaminants from water. *Aquatic toxicology*, 212, 1-10.
- Nazari, M. T., Simon, V., Machado, B. S., Crestani, L., Marchezi, G., Concolato, G., ... & Piccin, J. S. (2022). *Rhodococcus*: A promising genus of actinomycetes for the bioremediation of organic and inorganic contaminants. *Journal of Environmental Management*, 323, 116220.
- Pal, A., Bhattacharjee, S., Saha, J., Sarkar, M., & Mandal, P. (2022). Bacterial survival strategies and responses under heavy metal stress: A comprehensive overview. *Critical reviews in microbiology*, 48(3), 327-355.
- Pandey, S., Barai, P. K., & Maiti, T. K. (2013). Influence of heavy metals on the activity of antioxidant enzymes in the metal resistant strains of *Ochrobactrum* and *Bacillus* sp. *Journal of environmental biology*, 34(6), 1033.
- Pires, C. (2010). Bacteria in heavy metal contaminated soil: diversity, tolerance and use in remediation.
- Prabhakaran, P., Ashraf, M. A., & Aqma, W. S. (2016). Microbial stress response to heavy metals in the environment. *Rsc Advances*, 6(111), 109862-109877.
- Puente-Sánchez, F., Díaz, S., Penacho, V., Aguilera, A., & Olsson, S. (2018). Basis of genetic adaptation to heavy metal stress in the acidophilic green alga *Chlamydomonas acidophila*. *Aquatic toxicology*, 200, 62-72.
- Raab, A., & Feldmann, J. (2003). Microbial transformation of metals and metalloids. *Science Progress*, 86(3), 179-202.
- Roy, B., Maitra, D., Sarkar, S., Podder, R., Das, T., Ghosh, J., & Mitra, A. K. (2023). Biofilm and metallothioneins: A dual approach to bioremediate the heavy metal menace. *Environmental Quality Management*.
- Roy, R., Samanta, S., Pandit, S., Naaz, T., Banerjee, S., Rawat, J. M., ... & Saha, R. P. (2024). An overview of bacteria-mediated heavy metal bioremediation strategies. *Applied Biochemistry and Biotechnology*, 196(3), 1712-1751.
- Saha, R., Banerjee, D. B., Manna, S., & Banerjee, S. (2022). Microbial bioremediation: A promising approach to withstand heavy metal contamination in soil and its future possibilities. In *Synergistic Approaches for Bioremediation of Environmental Pollutants: Recent Advances and Challenges* (pp. 227-262). Academic Press.
- Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular biology and evolution*, 4(4), 406-425.
- Sharma, B., & Shukla, P. (2021). Lead bioaccumulation mediated by *Bacillus cereus* BPS-9 from an industrial waste contaminated site encoding heavy metal resistant genes and their transporters. *Journal of Hazardous Materials*, 401, 123285.
- Shukla, P. K., Misra, P., Maurice, N., & Ramteke, P. W. (2019). Heavy metal toxicity and possible functional aspects of microbial diversity in heavy metal-contaminated sites. *Microbial Genomics in Sustainable Agroecosystems: Volume 2*, 255-317.
- Syed, A., Zeyad, M. T., Shahid, M., Elgorban, A. M., Alkhulaifi, M. M., & Ansari, I. A. (2021). Heavy metals induced modulations in growth, physiology, cellular viability, and biofilm formation of an identified bacterial isolate. *ACS omega*, 6(38), 25076-25088.
- Tamás, M. J., Sharma, S. K., Ibstedt, S., Jacobson, T., & Christen, P. (2014). Heavy metals and metalloids as a cause for protein misfolding and aggregation. *Biomolecules*, 4(1), 252-267.

Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: molecular evolutionary genetics analysis version 11. *Molecular biology and evolution*, 38(7), 3022-3027.

Xin, X., Chen, B., Yang, M., Gao, S., Wang, H., Gu, W., ... & Zhang, B. (2023). A critical review on the interaction of polymer particles and co-existing contaminants: Adsorption mechanism, exposure factors, effects on plankton species. *Journal of Hazardous Materials*, 445, 130463.

Zargoosh, K., Abedini, H., Abdolmaleki, A., & Molavian, M. R. (2013). Effective removal of heavy metal ions from industrial wastes using thiosalicylhydrazide-modified magnetic nanoparticles. *Industrial & Engineering Chemistry Research*, 52(42), 14944-14954.

<https://blast.ncbi.nlm.nih.gov>

<https://files.zymoresearch.com> › *protocols* ›

<https://www.nimagen.com> › IFU_BRD_3-1_v4-1

<https://files.zymoresearch.com> › *protocols* ›

UNDER PEER REVIEW