

## Effect of different fertilizer recommendation approaches on yield, profitability and nutrient uptake by wheat (*Triticum aestivum*) in acid Alfisols of North Western Himalayas

### Abstract

A field experiment was conducted on wheat during *rabi*, 2019-20 at the experimental farm of Soil Science, CSK Himachal Pradesh Krishi Vishwavidyalaya, Palampur. The experiment was laid out in randomized block design with nine treatments which were replicated thrice. The effects of different fertilizer recommendation approaches on yield, profitability and nutrient uptake in wheat were evaluated. Different treatments had a significant effect on the yield and nutrient uptake of wheat crop. STCR treatment for the target yield of 35 q ha<sup>-1</sup> with FYM resulted in the highest wheat grain yield (35.4 q ha<sup>-1</sup>). However, the highest BC ratio was recorded in treatment for target yield 35 q ha<sup>-1</sup> without FYM followed by treatment 25 q ha<sup>-1</sup> without FYM. A significant increase in N, P, K and micronutrients (Fe, Mn, Zn and Cu) uptake by wheat grain and straw was observed in STCR based target yield plots as compared to general recommended dose, farmers' practice, and control. The STCR-based target yield approach maximizes economic yields by supplying nutrients precisely according to crop requirements, thereby enhancing fertilizer use efficiency and conserving expensive fertilizers.

**Keywords:** Wheat, Soil test crop response (STCR), Yield, Uptake, Profitability

### Introduction

Wheat (*Triticum aestivum*) is the primary staple food for around two billion people, accounting for 36 per cent of the world's population. It stands as a cornerstone crop in bolstering global food security, accounting for approximately 20 per cent of the world's dietary energy and protein intake (Gooding and Shewry, 2022). Beside food security it also plays a crucial role in economic stability and agricultural sustainability, making it an essential crop in both developed and developing countries. In India, it is the second most important cereal crop after rice and is a crucial component of the daily human diet.

In the current era, world faces the formidable challenge of producing sufficient food, fodder, fibre, and fuel to meet the diverse needs of its burgeoning human population. According to the United Nations, the global population is expected to reach 8.5 billion by 2030, 9.7 billion by 2050, and grow up to 11.2 billion by the end of the century. The UN Food and Agriculture Organization have estimated that world would need to produce 70 per cent more food in 2050 to fulfil the requirement of its exponentially growing population. As population growth rapidly accelerates, the pressure on food production intensifies, posing a significant threat to global food security. A growing human population demands increased food production. Farmers try to meet that demand by intensive farming, using fertilisers and pesticides. While intensive agriculture has been instrumental in feeding a growing global population and producing food more efficiently, it has also raised concerns regarding sustainability, environmental degradation, animal welfare, and the loss of biodiversity. Some intensive agricultural activities such as the cultivation of monocultures and imbalanced and inadequate use of fertilizers not only results in poor yields but also deteriorates soil fertility and increased multiple nutrient deficiencies. Over the past three decades in India, the intensive cultivation of high-yielding wheat varieties has heavily depleted soil nutrients. This has led to increased reliance on chemical fertilizers to boost yields. Chemical fertilizers

Comment [WU1]: mention nutritive value of Wheat

play a pivotal role in modern agriculture by providing essential nutrients that enhance crop yields. However, over-reliance on chemical fertilizers can have adverse effects on soil health and environment. Heavy use of fertilizers, can lead to soil degradation, water pollution, and biodiversity loss.

As we navigate the complexities of the 21<sup>st</sup> century, sustainable practices which prioritize soil health is a key to ensure that wheat remains a reliable source of nutrition and prosperity for future generations. Use of sustainable agricultural practice such as, precision farming, and integrated nutrient management practices including the balanced use of organic and inorganic sources of fertilizers can preserve soil health, enhance resilience to climate change, and ensure long-term productivity while minimizing environmental impacts.

The Soil Test Crop Response (STCR) approach for fertilizer application, first introduced by Trough (1960), is a scientific method for balanced fertilization, ensuring equilibrium between applied and soil-available nutrients. The method of fertilizer recommendations based on targeted yield of the crops also called “Prescription Based Fertilizer Recommendations”, avoid wide variations in soil rating limits, as it substitutes the exact values for soil available N, P and K (Rao and Srivastava, 2000). Among the different methods of fertilizer recommendations, the prescription based fertilizer application as per STCR approach for target yield is unique, as it involves both the soil test value of a nutrient along with the target yield that can be obtained with ideal agronomic practices (Singh *et al.*, 2005). The Soil Test Crop Response (STCR) approach for fertilizer application, integrates soil and plant analyses scientifically. Under the target yield approach, fertilizer doses are precisely prescribed using adjustment equations. It has emerged as a refined method for optimizing nutrient use from fertilizers and soil resources. Combining targeted yield-based fertilizer application with organic sources like farmyard manure (IPNS-STCR) enhances soil biological activity, physical properties, and chemical properties of soil. This holistic approach promotes ecologically sound nutrient management which not only improves soil health but also increases crop yields synergistically. Therefore, this research work was undertaken to evaluate the effect of different fertilizer recommendation approaches on yield, profitability and nutrient uptake in wheat crop.

## **Material and methods**

### **Experimental details**

The field experiment was conducted on wheat during *rabi*, 2019-20 at the Experimental farm of Department of Soil Science, Chaudhary Sarwan Kumar Himachal Pradesh Krishi Vishwavidyalaya, Palampur. The experimental farm is located at 32° 06' N latitude and 76° 3' E longitude at an elevation of 1290 m above mean sea level in Kangra district of Himachal Pradesh.

### **Climate and weather**

The climate of the experimental site is characterized as wet-temperate. The region has mild summers from March to June and cool winters during December to February with high rainfall during monsoon months. The region receives an average annual rainfall of 2500 mm to 3000 mm per annum with 75% of the showers occurring mainly during monsoon months (June - September). During *rabi*, 2019-2020, the region received a total rainfall of 788 mm and the mean maximum and minimum temperatures were 19.6°C and 17.6°C, respectively.

### Soil characteristics

The soil of the experimental site belonged taxonomically to the order *Alfisol* under the sub-group *Typic Hapludalf*. The soils of the experimental site were silty clay loam in texture, acidic in reaction with pH 5.47, medium in organic carbon 7.80 g kg<sup>-1</sup>, low in available nitrogen 217 kg ha<sup>-1</sup>, high in available phosphorus 28 kg ha<sup>-1</sup> and medium in soil available potassium 234 kg ha<sup>-1</sup> status.

### Experimental details

The experiment was laid out in a randomized block design consisting of nine treatment combinations (Table 1). The treatments were replicated thrice with plot size of 15 m<sup>2</sup>. The details of treatments used were as follows:

**Table 1. Treatment details of the experiment**

Symbol	Description
T <sub>1</sub>	Control
T <sub>2</sub>	Farmers' practice
T <sub>3</sub>	General recommended dose
T <sub>4</sub>	Soil test based fertilizer application
T <sub>5</sub>	Fertilizer dose for target yield 25 q ha <sup>-1</sup>
T <sub>6</sub>	Fertilizer dose for target yield 25 q ha <sup>-1</sup> with FYM @ 5 t ha <sup>-1</sup>
T <sub>7</sub>	Fertilizer dose for target yield 35 q ha <sup>-1</sup>
T <sub>8</sub>	Fertilizer dose for target yield 35 q ha <sup>-1</sup> with FYM @ 5 t ha <sup>-1</sup>
T <sub>9</sub>	<i>Beejamrit + Jeevamrit + Ghanajeevamrit</i>

Wheat (HPW-236) was taken as a test crop. The sowing of crop was done on 14<sup>th</sup> November, 2019 with a seed rate of 100 kg ha<sup>-1</sup>. The row to row spacing was kept 22 cm. All the recommended cultural practices were followed during the entire crop growth. At the time of sowing, half dose of nitrogen and full doses of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O were applied according to the treatments as a basal application. The remaining half dose of nitrogen was applied into two equal splits, one at the time of tillering and another during flowering stage. In T<sub>9</sub>, *Ghanajeevamrit* @ 250 kg ha<sup>-1</sup> was incorporated in the soil before sowing of wheat. The wheat seeds were treated with *Beejamrit* @ 101 kg ha<sup>-1</sup> and then dried in shade before sowing. *Jeevamrit* @ 500 liters ha<sup>-1</sup> was also applied during sowing followed by its spray after 21 days interval with 1:10 dilution in water. Lantana was used for mulching. FYM was applied @ 5 t ha<sup>-1</sup> (dry weight basis) in all the STCR-IPNS treatments.

The following equations given by (Verma et al., 2007) developed for computing the STCR- based fertilizers doses of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O under wheat crop were as follows:

$$FN = 5.27 T - 0.25 SN - 1.06 ON$$

$$FP_2O_5 = 4.13 T - 0.38 SP - 0.98 OP$$

$$FK_2O = 2.87 T - 0.15 SK - 0.55 OK$$

In the above adjustment equations FN, FP<sub>2</sub>O<sub>5</sub>, FK<sub>2</sub>O are doses of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O, respectively in kg ha<sup>-1</sup>. T is the yield target (q ha<sup>-1</sup>), SN, SP and SK are soil available N, P and K contents before sowing of the crop, respectively in kg ha<sup>-1</sup>. Whereas ON, OP and OK are N, P and K supplied by FYM, respectively in kg ha<sup>-1</sup>.

### Laboratory studies

For the chemical analysis of nutrient content in plant, representative grain and straw samples were collected after the harvest of wheat crop. The samples were then air dried and later kept in the hot air oven at 60-70°C for eight hours. The dried samples were then ground and stored in paper bags for subsequent analysis. The analytical methods employed for plant analysis are as follows:

**Table 2. Analytical methods used for FYM and plant analysis**

Parameter	Method	References
Nitrogen	Micro-kjeldahl	Jackson (1973)
Phosphorus	Vanado-molybdo-phosphoric acid	Jackson (1973)
Potassium	Wet digestion	Black (1965)
Zn, Fe, Mn, Cu	Wet digestion	Jackson (1967)

The nutrient uptake was calculated by multiplying (%) concentration of a nutrient with grain and straw yields (q ha<sup>-1</sup>) on dry weight basis using the given formula:

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = [\% \text{ concentration of nutrient} \times \text{yield of crop in q ha}^{-1} \text{ (oven dry weight basis)}].$$

**Comment [WU2]:** it should divided by 100 for N P K

Total uptake = uptake by grains + uptake by straw

### Economic analysis

The economics of the experiment was worked out in terms of net returns and BC ratio. Cost of cultivation (Rs ha<sup>-1</sup>) was obtained by adding the costs of all the inputs and each operation involved during the experimental trial of wheat crop. Gross returns (Rs ha<sup>-1</sup>) were calculated by multiplying yield of produce (q ha<sup>-1</sup>) under different treatments with the price of the produce in the market. Net returns (Rs ha<sup>-1</sup>) were computed by deducting the cost of inputs from gross returns and BC ratio was calculated by dividing gross returns with cost of cultivation.

### Statistical analysis

The experimental data were statistically analyzed for the differential effect of treatments at 5% level of significance for randomized block design as per the standard procedure (Gomez and Gomez, 1984).

## Results and Discussion

### Effect of different fertilizers recommendation approaches on grain and straw yield of wheat

#### Grain and straw yield

Application of fertilizers based on different approaches significantly influenced the grain and straw yield of wheat (Table 1). Wheat grain and straw yield varied from the lowest (14.4 q ha<sup>-1</sup>) and (23.9q ha<sup>-1</sup>) in T<sub>1</sub>(control) to the highest (35.4 q ha<sup>-1</sup>) and (53 q ha<sup>-1</sup>) in T<sub>8</sub>(target yield of 35 q ha<sup>-1</sup> with FYM),

respectively. All the STCR based treatments were significantly superior with respect to the grain yield over the other treatments. Both the STCR based IPNS treatments were at par with their respective non-IPNS treatments with respect to the grain yield and straw yield of wheat. Treatment T<sub>4</sub> (soil test based application of fertilizers) recorded significantly higher grain and straw yield compared to T<sub>2</sub> (farmers' practice), T<sub>9</sub> (treatment where *Beejamrit*, *Jeevamrit*, *Ghanajeevamrit* were applied along with mulch), and T<sub>1</sub> (control). However, it was at par with T<sub>3</sub> (where general recommended dose of fertilizers were applied). Increased yield under STCR approach with and without FYM might be because of balanced use of fertilizers as per soil and crop demand for potential growth and development (Kumar *et al.*, 2023). (Singh *et al.*, 2016) also revealed that the sole or combine application of fertilizers with FYM based on target yield approach had a marked effect on grain and straw yields of wheat. Similar results were reported by (Tiwarie *et al.*, 2013) and (Sellamuthu *et al.*, 2015).

**Comment [WU3]:** mention about IPNS in materials and methods for clear understanding

### Effect of different fertilizers recommendation approaches on profitability of wheat

The highest net returns of Rs 77,695 ha<sup>-1</sup> were manifested under target yield of 35 q ha<sup>-1</sup> without FYM (T<sub>7</sub>) followed by the treatment with same yield target with FYM (T<sub>8</sub>) Rs 69,649 ha<sup>-1</sup> (Table 1). Application of fertilizers based on soil test values (T<sub>4</sub>) recorded net returns of Rs 49,374 ha<sup>-1</sup> which were Rs 6,170 ha<sup>-1</sup>, Rs 26,846 ha<sup>-1</sup> and Rs 21,199 ha<sup>-1</sup> higher in comparison to general recommended dose (T<sub>3</sub>), farmers' practice (T<sub>2</sub>) and *Beejamrit* + *Jeevamrit* + *Ghanajeevamrit* (T<sub>9</sub>). With respect to the benefit drawn per unit rupee invested on inputs, treatment target yield 35 q ha<sup>-1</sup> without FYM (T<sub>7</sub>) was the most remunerative followed by the treatment comprising of target yield 25 q ha<sup>-1</sup> without FYM (T<sub>5</sub>) with the BC ratios of 3.58 and 3.08, respectively. Treatment farmers' practice (T<sub>2</sub>) was found to be least remunerative treatment with the BC ratio of 1.72. The BC ratio was higher in 25 and 35 q ha<sup>-1</sup> targeted treatments without FYM when compared with same targeted treatments with FYM. Application of FYM increased the net returns and BC ratio in IPNS based STCR treatments over non-IPNS based STCR treatments. Almost similar results have also been reported by (Singh *et al.*, 2016) and (Choudhary and Dixit, 2022).

**Table 3. Effect of different fertilizers recommendations approaches on yield (q ha<sup>-1</sup>) and profitability of wheat**

Treatments	Grain yield (q ha <sup>-1</sup> )	Stover yield (q ha <sup>-1</sup> )	Net Returns (Rs ha <sup>-1</sup> )	B:C Ratio
Control	14.4	23.9	28695	2.57
Farmers' practice	16.5	27.2	22528	1.72
General recommended dose	21.1	33.7	43204	2.77
Soil test based	22.9	36.3	49374	3.07
Target yield 25q ha <sup>-1</sup>	25.5	39.8	54703	3.08

Target yield 25q ha <sup>-1</sup> with FYM @ 5t ha <sup>-1</sup>	26.8	41.6	48198	2.31
Target yield 35q ha <sup>-1</sup>	34.8	51.1	77695	3.58
Target yield 35q ha <sup>-1</sup> with FYM @ 5t ha <sup>-1</sup>	35.4	53.0	69649	2.70
Beejamrit + Jeevamrit + Ghanajeevamrit	16.9	28.2	28175	2.04

Sale price of wheat, Grain= Rs 18.5 Rs kg<sup>-1</sup>, Straw = Rs. 850 Rs q<sup>-1</sup>; Jeevamrita=1.5(Rs t<sup>-1</sup>), Beejamrita=2(Rs t<sup>-1</sup>)


Cost of fertilizer (Rs kg<sup>-1</sup>) N=11.9, P<sub>2</sub>O<sub>5</sub>=65.8, K<sub>2</sub>O =28. FYM = Rs. 100 q<sup>-1</sup>

## Plant nutrients uptake

### Macronutrients

Data presented in fig. 1. indicated significant variation in the nutrient uptake in wheat crop. A critical examination of the data revealed that the N, P and K uptake in wheat grain and straw was maximum under STCR based IPNS treatment for target yield of 35q ha<sup>-1</sup> (T<sub>8</sub>) and minimum in control (T<sub>1</sub>). N uptake in wheat grain and straw varied from 25.4 kg ha<sup>-1</sup> and 11.8 kg ha<sup>-1</sup> under control (T<sub>1</sub>) to 75.3 kg ha<sup>-1</sup> and 35.9 kg ha<sup>-1</sup> under target yield 35 q ha<sup>-1</sup> with FYM @ 5 t ha<sup>-1</sup> (T<sub>8</sub>), respectively. P uptake in wheat grain and straw varied from 4.04 kg ha<sup>-1</sup> and 2 kg ha<sup>-1</sup> in control (T<sub>1</sub>) to 11.8 kg ha<sup>-1</sup> and 8.13 kg ha<sup>-1</sup> in target yield 35 q ha<sup>-1</sup> with FYM @ 5 t ha<sup>-1</sup> (T<sub>8</sub>), respectively. K uptake in wheat grain and straw varied from 4.55 kg ha<sup>-1</sup> and 16.4 kg ha<sup>-1</sup> in control (T<sub>1</sub>) to 14.6 kg ha<sup>-1</sup> to 42 kg ha<sup>-1</sup> in target yield 35 q ha<sup>-1</sup> with FYM @ 5 t ha<sup>-1</sup> (T<sub>8</sub>), respectively. Supplementation of NPK fertilizers as per STCR approach with and without FYM significantly enhanced N, P and K uptake in wheat crop over general recommended dose, ZBNF treatment, farmers' practice and control. The significant increase in nitrogen, phosphorus and potassium uptake in grain and straw of wheat under STCR-IPNS treatments could be due to improved nutrient supplying capacity and flush of available nutrients. Simultaneous application of organic manure and chemical fertilizers results in the faster decomposition of chemical fertilizers compared to the organic manure, hence resulting in the increased nutrient availability to the plants. During the time of organic manure decomposition, many organic acids are released which break down the complex nitrogenous compounds gradually, providing a steady supply of nitrogen, phosphorus, and potassium throughout the crop growth period thereby, increasing wheat production and uptake of nutrients from soil system. (Singh *et al.*, 2020). Similar findings have been reported by (Choudhary and Dixit, 2022) and (Kumari *et al.*, 2022) on wheat crop.

Comment [WU4]: clarify



**Fig. 1. Effect of different recommendation of fertilizer application approaches on nitrogen, phosphorus and potassium uptake (kg ha<sup>-1</sup>) by wheat**

### Micronutrients

The results suggested that the different approaches of fertilizer application manifested a significant effect on the uptake of micronutrients by wheat over control (Fig. 2). The uptake of micronutrients was found to be maximum in the treatment target yield of 35 q ha<sup>-1</sup> with FYM (T<sub>8</sub>) and minimum in control (T<sub>1</sub>). All the STCR based target yield treatments with or without FYM were statistically superior over general recommended dose, ZBNF treatment, farmers' practice and control. Micronutrients uptake under treatments target yield of 25 q ha<sup>-1</sup> with 5 t ha<sup>-1</sup> FYM (T<sub>6</sub>), target yield of 35 q ha<sup>-1</sup> (T<sub>7</sub>) and target yield of 35 q ha<sup>-1</sup> with 5 t ha<sup>-1</sup> FYM (T<sub>8</sub>) was statistically higher compared to soil test based (T<sub>4</sub>). FYM is a good source of nutrients and growth promoting substances and higher uptake of these micronutrients in treatments supplied with FYM might be attributed to higher content of these micronutrients present in FYM, presence of higher microbial and enzymatic activity which stimulated the root growth and resulted in higher uptake (Laxminarayana and Patiram, 2006). Moreover, addition of organic manures lower the soil pH due to the release of various acids upon its breakdown, which enhances the bioavailability of

micronutrients compared to the sole use of inorganic fertilizers (Dhaliwal *et al.*, 2023). Similar type of results were observed by (Naidu *et al.*, 2008) and (Nandapure *et al.*, 2011).




Fig. 2. Effect of different recommendation of fertilizer application approaches on iron, manganese, zinc and copper uptake (g ha<sup>-1</sup>) by wheat

### Conclusion

The findings of present study concluded that, the combined application of FYM and fertilizers under the STCR-based IPNS approach not only led to higher wheat yields but also resulted in the highest nutrients uptake in both wheat grains and straw. The STCR-based target yield approach maximizes economic yields by supplying nutrients precisely according to crop requirements, thereby enhancing fertilizer use efficiency and conserving expensive fertilizers. This approach can significantly benefit the resource-poor farmers by not only boosting yields but also reducing the cultivation costs.

[Can add review which supports present study](#)  
[“Use of Soil Test Crop Response Approach in Direct Seeded Rice](#)

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

**Vidvavathi G Yadahalli\* and M R Kammar,**

**J Krishi Vigyan 2017, 6(1) : 213-216**

**DOI : 10.5958/2349-4433.2017.00082.4**

**Nutrient management in Bt. cotton through STCR**

**Approach, Vidvavathi, G. Yadahalli1, Ravi M.V2, G.S. Yadahalli1 and Shivaleela S. Kukanur2**

*Eco. Env. & Cons. 20 (1) : 2014; pp. (337-340)*

*Copyright@ EM International*

*ISSN 0971-765X*

Formatted: Font: (Default) Times New Roman

Formatted: Font: 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: 12 pt

## References

Black, C. A. (1965). Methods of soil analysis. Part II. Chemical and mineralogical properties. American Society of Agronomy, Madison, Wisconsin, USA.

Choudhary, D. & Dixit, S. P. (2022). Long term effect of STCR-based targeted yield application of fertilizers on productivity, profitability and nutrient uptake by wheat (*Triticum aestivum* L.) in an acid Alfisol. *Annals of Plant & Soil Research*, 24(3), 373-377.

Dhaliwal, S. S., Sharma, V., Shukla, A. K., Gupta, R. K., Verma, V., Kaur, M., Behera, S. K. & Singh, P. (2023). Residual Effect of Organic and Inorganic Fertilizers on Growth, Yield and Nutrient Uptake in Wheat under a Basmati Rice–Wheat Cropping System in North-Western India. *Agriculture*, 13(3), 556-573.

Gomez, K. A. and Gomez, A. A. (1984). Statistical Procedure for Agricultural Research. John Wiley and Sons, New York

Gooding, M. J. and Shewry, P.R. (2022). Wheat: environment Food and Health, Wiley

Jackson, M. L. (1967). Soil Chemical Analysis. Prentice Hall of India, Limited, New Delhi. P 219-221

Jackson, M. L. (1973). Soil Chemical Analysis. Prentice Hall Inc. Englewood Cliffs, New Jersey, USA.

Kumar, A., Singh, A. K., Rai, A., Gangwar., Kumar, M., Pareek, N., Kapil, S. T., Chanu, N. B., Padhi, G. K., Panda, P. P., Gupta, R. K. & Kumar, S. (2023). Productivity and economics of wheat under STCR based nutrient management in alluvial soil. *Journal of AgriSearch*, 10(4), 217-220.

Kumari, N., Kharia, S. K., Mandeewal, R. L., Rohj, M. K., Yadav, A. & Kumari, S. (2022). Effect of Soil Test Crop Response Approach on Yield, Nutrient Content and Uptake by Wheat (*Triticum aestivum* L.). *International Journal of Plant & Soil Science*, 34(14), 1-6.

Laxminarayana, K. & Patiram, (2006). Effect of integrated use of inorganic, biological and organic manures on rice productivity and soil fertility in ultisols of Mizoram. *Journal of Indian Society of Soil Science*, 2, 213-220.

- Naidu, L. G. K., Ramamurthy, V., Kumar, R., Sidhu, G. S. & Kumar, R. (2008). Soil based fertilizer recommendations. *Indian Journal of Fertilizers*,4, 47-56.
- Nandapure, S. P., Sonune, B. A., Gabhane, V. V., Katkar, R. N. & Patil, R. T. (2011). Influence of long term fertilization on micronutrients availability, their uptake and productivity of sorghum-wheat sequence under semi-arid condition on a Vertisol. *Crop Research*,42, 35-39.
- Rao, S. & Srivastava, S. (2000). Soil test based fertilizer use-a must for sustainable agriculture. *Fertilizer News*, 45, 25-38.
- Sellamuthu, K. M., Santhi, R., Maragatham, S. & Dey, P. (2015). Validation of soil test and yield target based fertilizer prescription model for wheat on Inceptisol. *Research on Crops*,16, 53-58.
- Singh, H., Ingle, S. R., Pratap, T., Raizada, S., Singh, P. K., Singh, R. & Parihar, A. K. S. (2020). Effect of integrated nutrient management on nitrogen content, uptake and quality of wheat (*Triticum aestivum* L.) under partially reclaimed sodic soil. *The Pharma Innovation Journal*, 9(5), 299-301.
- Singh, K. N., Raju, N. S., Rao, A. S., Rathore, A., Srivastava, S., Samanta, R. K. & Maji, A. K. (2005). Prescribing optimum dose of nutrients for targeted yield through soil fertility map in Andhra Pradesh (AP). *Journal of Indian Society of Agricultural Statistics*,59, 131-140.
- Singh, Y. V., Dey, P., Meena, R. & Varma, S. K. (2016). Effect of soil test based fertilizer application on yield and economics of chickpea in Inceptisol. *Annals of Plant and Soil Research*, 18, 409-412.
- Tiwari, L. B., Mishra, D. D. & Gupta, J. (2013). Evaluation of integrated nutrient management options in rice (*Oryza sativa*)- wheat (*Triticum aestivum*) cropping system in reclaimed sodic land. *Indian Journal of Agronomy*, 58, 137-14.
- Trough, E. (1960). Fifty years of soil testing. Transactions 7<sup>th</sup> International Congress of Soil Science 3: 46- 57
- Verma, T. S., Sharma, S. K., Kumar, P., Suri, V. K., Sandal, S., Murlidharudu, Y. & Sankhyan, N. K. (2007). Soil Test Crop Response: An approach for fertilizer recommendation based on target yield concept: A success story in Himachal Pradesh, Technical Bulletin, Department of Soil Science, CSK Himachal Pradesh Krishi Vishvavidyalaya, Palampur, India.