

Effect of N and K fertilization on vegetative growth of *Acroclinium* (*Helipterum roseum*)

Abstract

The present investigation entitled "Effect of N and K fertilization on vegetative growth of *Acroclinium* (*Helipterum roseum*)" was carried out at the laboratory of floriculture and experimental farm of Mata Gujri college, Fatehgarh Sahib, Punjab. The experiment was laid out in randomized block design (RBD) with twelve treatments and three replications. The treatments were T₁ i.e., 125 kg/acre urea + 60 kg/acre MOP, T₂ i.e., 175 kg/acre urea + 60 kg/acre MOP, T₃ i.e., 225 kg/acre urea + 60 kg/acre MOP, T₄ i.e., 125 kg/acre urea + 80 kg/acre MOP, T₅ i.e., 175 kg/acre urea + 80 kg/acre MOP, T₆ i.e., 225 kg/acre urea + 80 kg/acre MOP, T₇ i.e., 125 kg/acre urea + 0 kg/acre MOP, T₈ i.e., 175 kg/acre urea + 0 kg/acre MOP, T₉ i.e., 225 kg/acre urea + 0 kg/acre MOP, T₁₀ i.e., 0 kg/acre urea + 80 kg/acre MOP, T₁₁ i.e., 0 kg/acre urea + 60 kg/acre MOP and T₁₂ i.e., control. Observations were recorded for vegetative, flowering and seed yield parameters. Among all treatments maximum plant height (44.86 cm and 59.91 cm), leaf length (5.01 cm and 5.33 cm), plant spread (17.68 cm² and 32.37 cm²) and diameter of main stem (6.13 mm and 10.77 mm) were found maximum in T₃ i.e., 225 kg/acre urea + 60 kg/acre MOP and maximum number of leaves per stem (37.28 and 54.98) and number of leaves per plant (645.67 and 1055.13) was observed in T₉ i.e., 225 kg/acre urea + 0 kg/acre MOP at 45 and 90 days respectively. Best vegetative parameters were recorded in 225 kg/acre urea + 60 kg/acre MOP and 225 kg/acre urea + 0 kg/acre MOP are best dosage for *Acroclinium* flower.

Keywords: *Acroclinium*, Urea, MOP, Fertilizer

Introduction

The growing demand for floricultural products as a result of rising living standards and increased awareness among people of the need to live in an environmentally friendly environment has made it a profitable agribusiness in recent years. The plant *Acroclinium*, botanically known as *Helipterum roseum*, is frequently referred to as "paper daisies" or "everlasting flowers". The genus *Acroclinium*, which is native to Australia and belongs to the family Asteraceae of flowering plants, is well-known for both its ecological resilience and ornamental value.

The herb *Helipterum roseum* blooms from late spring to early autumn, adding a delicate and charming touch to gardens and floral arrangements. Its ability to retain its colour even after drying makes it a favourite for crafting everlasting bouquets. *Acroclinium* can adapt to different climates, but it generally thrives well in warm and sunny conditions. It is well-suited for regions with a Mediterranean climate or similar, with dry summers and mild winters. This flower prefers well-drained soil and plenty of sunlight to bloom beautifully.

MATERIAL AND METHODS

The present investigation was conducted at Research Farm, Mata Gujri College, Fatehgarh Sahib, Punjab during 2023-2024. Field of experimental site lies at 30.6435° North latitude and 76.3970° East longitudes. The altitude of the location is 246 meter above the mean sea level. The nursery of *Acroclinium* were obtained from the Biocarve seeds, Dhablan (Patiala). The experiment was laid out in Randomized Block Design (RBD) with twelve treatments three replications. The treatments were T₁ i.e., 125 kg/acre urea + 60 kg/acre MOP, T₂ i.e., 175 kg/acre urea + 60 kg/acre MOP, T₃ i.e., 225 kg/acre urea + 60 kg/acre MOP, T₄ i.e., 125 kg/acre urea + 80 kg/acre MOP, T₅ i.e., 175 kg/acre urea + 80 kg/acre MOP, T₆ i.e., 225 kg/acre urea + 80 kg/acre MOP, T₇ i.e., 125 kg/acre urea + 0 kg/acre MOP, T₈ i.e., 175 kg/acre urea + 0 kg/acre MOP, T₉ i.e., 225 kg/acre urea + 0 kg/acre MOP, T₁₀ i.e., 0 kg/acre urea + 80 kg/acre MOP, T₁₁ i.e., 0 kg/acre urea + 60 kg/acre MOP and T₁₂ i.e., control.

RESULTS AND DISCUSSIONS

Plant height (cm)

The maximum plant height was observed in T₃ i.e., (urea 225 kg + 60 kg MOP per acre) (59.91cm) which was statistically superior than other treatments. When nitrogen and potash are applied in combination, the plant receives a more balanced nutritional supply, resulting in robust leaf growth, strong stems, and improved plant structure. This combination provides for greater plant height since nitrogen promotes general growth and potash supports strong stem development. This finding is similar with the results of Ahmed *et al.* (2015) in gladiolus plant. Sowmyamala (2013), Gawade *et al.* (2016) in gaillardia. Minimum plant height was observed in T₁₂ i.e., control (48.15 cm), which was statistically inferior due to the insufficient availability of nutrients to plant during critical period for its luxuriant growth. The present findings are also found in Maheta (2015) in china aster, Acharya and Dashora (2004), Sharma *et al.* (2006) in marigold and Joshi (2014), in chrysanthemum.

Number of leaves per stem

There was significant effect on the number of leaves per stem of *Acroclinium*. The maximum number of leaves were observed in T₉ i.e., (urea 225 kg per acre + 0 kg MOP per acre) (37.28), which was statistically at par with T₈ i.e., (175 kg urea + 0 kg MOP per acre) (36.39). Number of leaves per stem increased linearly with corresponding increase in nitrogen doses. This may be attributed to the fact that nitrogen is an essential part of chlorophyll and nucleic acids, which might have played major role in promoting plant growth. Higher nitrogen concentrations can improve a leaf cell size, number, and overall production of leaves, as well as other aspects of the plant's vegetative growth (Meyer *et al.*, 1973; Sigedar *et al.*, 1991). Similar result was found by Aruna *et al.* (2007) in crossandra. Minimum number of leaves per stem was observed in T₁₂ i.e., control (42.98) which was statistically inferior than other treatments. At the time of critical growth, the availability of N is less due to which it produces less number of leaves per stem. Similar results were found by Grewal *et al.* (2004) in plants of chrysanthemum.

Number of leaves per plant

The maximum number of leaves per plant were observed in T₉ i.e., (urea 225 kg + 0 kg MOP per acre) (1055.13), which was statistically superior. Which might be due to the reason that the nitrogen is a component of chlorophyll, the pigment that gives leaves their green coloration, and it is necessary for photosynthesis, the process by which plants manufacture food. When plants acquire enough nitrogen, they can grow more leaves because cell division and expansion are stimulated, resulting in greater leaf area and total leaf production. Similar results were found by Gupta *et al.* (1999) and Chadha *et al.* (1999) in marigold and Devi *et al.* (2002) in carnation. The minimum number of leaves per plant was found in T₁₂ i.e., control (799.31) which was statistically inferior. The reduction in number of leaves per plant could be due to the lack of sufficient nutrients at critical stage of plant growth. Similar results were found by Agrawal *et al.* (2002) and Seharwat *et al.* (2003) in marigold.

Leaf length (cm)

Result of experiment disclose that, compared to all other treatments tested, T₃ i.e., (225 kg urea +60 kg MOP per acre) (5.33cm), produced highest leaf length. When nitrogen and potash applications are combined, they can work synergistically to promote floral crop growth and development, including increased leaf length. A balanced supply of essential minerals can promote healthy plant growth, resulting in longer and more robust leaves in flower yields. Similar findings are recorded in Gladiolus by Muhammad *et.al* (2020) in chrysanthemum by Teja *et al.* (2015-16). In treatment T₁₂ i.e., control (3.92cm) the minimum leaf length was observed which was statistically inferior. In the absence of enough nutrients, plants are likely to suffer suboptimal growth, resulting in shorter leaf length and overall weaker plant development as compared to plants that receive the appropriate nutrients. Same results found by Maheta (2015) in china aster.

Plant spread (cm²)

The maximum plant spread was observed in T₃ i.e., (225 kg urea + 60 kg MOP per acre) (32.37 cm²), which is statistically at par with T₆ i.e., (225 kg urea + 80 kg MOP per acre) (31.77 cm²). This might be due to increased synthesis of protein and protoplasm as nitrogen is an essential part of nucleic acid which plays a vital role in promoting plant growth (Wandleigh, 1957). Potassium is a vital mineral that aids in the production of starch and sugar in plant tissues. It stimulates the growth of weak developing sections, regulates stomatal movement, and balances water relationships. Potassium also plays a crucial role in carbohydrate formation and starch translocation, which leads to better plant development. In combination with nitrogen and potassium, there was a significant improvement in plant spread. Seharwat *et al.* (2003) similarly found that increasing N and K levels improved growth rates. The minimum plant spread was seen in T₁₂ i.e., control (25.49 cm²) which was significantly statically inferior. Due to insufficient amount of nutrients availability to plant during critical period of time for its development may cause a decline in plant spread. Similar results found in Acharya and Dashora (2004), in chrysanthemum.

Comment [VK1]: Use significantly than statically.

Diameter of main stem (mm)

The maximum diameter of main stem was observed in T₃ i.e., (225 kg urea + 60 kg MOP per acre) (10.7 mm). The increase in stem diameter might be due to the fact that nitrogen being a

constituent of protein, nucleic acids and nucleotides which are essential for the different metabolic function of plants (Bijimol and Singh 2001). Whereas potassium is an essential constituent which helps the formation of starch and sugar in plant body. It promotes the growth of tender growing parts, adjust stomata movement and balance water relationship. Similar results were also obtained by Khalaj and Edrisi et al. (2012) in tuberose, Dhaked et al. (2013) in calendula. The minimum diameter of main stem was found in T₁₂ i.e., control (6.74 mm). Insufficient amount of nutrients availability to plant during critical times for its luxuriant growth may cause a decline in diameter of main stem. Similar result was found by Sharma et al. (2011) in marigold.

Comment [VK2]: Not there in the reference

Formatted: Highlight

Table 1: Effect of N and K fertilization on vegetative growth, *Acroclinium (Helipterum roseum)*

TREATMENTS	Plant height (cm)	Number of leaves per stem	Number of leaves per plant	Leaf length (cm)	Plant spread (cm ²)	Diameter of main stem (mm)
T ₁ 125 kg/acre urea + 60 kg/acre MOP	55.49	47.27	860.08	4.24	26.98	7.20
T ₂ 175 kg/acre urea + 60 kg/acre MOP	56.81	49.02	866.96	4.50	29.02	7.81
T ₃ 225 kg/acre urea + 60 kg/acre MOP	59.91	50.42	910.92	5.33	32.37	10.77
T ₄ 125 kg/acre urea + 80 kg/acre MOP	55.45	48.69	891.21	4.36	27.24	7.83
T ₅ 175 kg/acre urea + 80 kg/acre MOP	56.50	49.60	902.48	4.42	28.09	8.35
T ₆ 225 kg/acre urea + 80 kg/acre MOP	57.49	51.02	954.46	5.09	31.77	9.42

T ₇ 125 kg/acre urea + 0 kg/acre MOP	53.49	48.00	858.46	4.15	27.57	7.58
T ₈ 175 kg/acre urea + 0 kg/acre MOP	54.12	51.81	906.46	4.37	29.66	7.42
T ₉ 225 kg/acre urea + 0 kg/acre MOP	54.93	54.98	1055.13	4.36	30.26	7.73
T ₁₀ 0 kg/acre urea + 80 kg/acre MOP	52.96	46.25	851.48	4.42	26.48	7.89
T ₁₁ 0 kg/acre urea + 60 kg/acre MOP	51.48	45.48	819.13	4.21	26.78	7.05
T ₁₂ Control	48.15	42.98	799.31	3.92	25.49	6.74
SEm±	0.24	0.39	1.05	0.04	0.21	0.09
CD _{0.05}	0.70	1.14	3.09	0.13	0.61	0.28

CONCLUSION

From the present study it can be concluded that T₃ @ 225 kg/acre urea + 60 kg/acre MOP performed best in various vegetative parameters such as plant height (59.91 cm), leaf length (5.33 cm), plant spread (32.37 cm²) and diameter of main stem (10.77 mm). While the highest dosage of urea i.e., T₉ @ 225 kg/acre urea + 0 kg/acre MOP has reported the best performance in number of leaves per stem (54.98) and number of leaves per plant (1055.13).

From the results, it can be concluded that 225 kg/acre urea + 60 kg/acre MOP and 225 kg/acre urea + 0 kg/acre MOP are best dosage for *Helipterum roseum*.

REFERENCES

Abhipsa, P., Palai, S.K. and Nath, M.R. 2018. Effect of source of nitrogen on growth and yield of African marigold (*Tagetes erecta* L.). The Pharma Innovative J. 2018. 7(7):917-921

Formatted: Highlight

Acharya, M. M. and Dashora, L. K. 2004. Response of graded levels of nitrogen and phosphorus on vegetative growth and flowering in African marigold (*Tagetes erecta* Linn.). J. Ornamental Hort., 7(2): 179-183.

Comment [VK3]: Follow author guidelines for reference arrangement
Sample reference
1.Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. Clin Exp Allergy. 2002;32(4):489-98.

Formatted: Highlight

Ahmad I, Ahmad T, Zafar M S and Nadeem A. 2007. Response of an elite cultivar of zinnia (*Zinnia elegans* cv. 'Giant Dahlia Flowered') to varying levels of nitrogenous fertilizer. Sarhad Journal of Agriculture. 23(2): 309-312.

Ahmed, S., S. Sultana, M.A. Kawochar, S. Naznin and F.U. Tuli. 2015. Effect of Corm Size and Nitrogen on the Growth and Flowering of Gladiolus (*Gladiolus grandifloras* L.). Emer. Life Sci. Res., 1(1), 18-25.

Dhaked R, Satish C and Srivastava R. 2013. Effect of spacing and levels of nitrogen on growth, flowering and yield of Calendula (*Calendula officinalis*) single type. Pantnagar Journal of Research 11:365-368.

Dorajeerao, A.V.D. 2009. Standardisation of Production Technology in Garland Chrysanthemum (*Chrysanthemum coronarium* L.). Ph.D. Thesis. Dept. of Horticulture, University of Agricultural Sciences, Dharwad (Karnataka).

Formatted: Highlight

Gawade, N. V., Bhalekar, S. G., Katwate, S. M., and Wadekar, V. D. 2016. Studies on different genotypes of gaillardia (*Gaillardia pulchella* L.) for quantitative and qualitative performance. Int. J. Curr. Microbiol. App. Sci. 7 (03): 1030 - 1039.

Grewal H S, Kumar R and Singh H. 2004. Effect of nitrogen, planting time and pinching on flower production in chrysanthemum (*Dendranthema grandiflora* Ramat.) cv. 'Flirt'. Journal of Ornamental Horticulture. 7(2): 189-192.

Khalaj MA, Edrisi B and Amiri M.2012. Effect of nitrogen and plant spacing on nutrient uptake, yield and growth of tuberose (*Polianthes tuberosa* L.). Journal of Ornamental and Horticultural Plants 2:45-54. Khan FN, Rahman MM and Karim AS. 2012. Effects of nitrogen and potassium on growth and yield of gladiolus corms. Journal of Agricultural Research 37:607-616.

Lehri, S.M., A.A. Kurd, M.A. Rind, and N.A. Bangulzai. 2011. The response of Gladiolus April 2020 | Volume 36 | Issue 2 | Page 401 Sarhad Journal of Agriculture tristis L. to N and P2O5 fertilizers. Sarhad J. Agric., 27(2): 185-188.

P. Ravi Teja et al.2016. Effect of graded levels of nitrogen and potassium on growth and flower yield of annual chrysanthemum (*Chrysanthemum coronarium* L.) Plant Archives Vol.17 No.2,2017 pp. 1371-1376.

Salve, S. D. 2017. Studies on growth and yield performance of promising genotypes of gaillardia (*Gaillardia pulchella* L.) M. Sc. (Hort.) thesis submitted to Mahatma Phule Krishi Vidyapeeth, Rahuri (Maharashtra).

Sehrawat, S.K., D.S. Dahiya, Sukhbir Singh and G. S. Rana, 2003. Growth, flowering and corm production in gladiolus as influenced by NPK application. Haryana Journal of Horticulture Sciences, 32 (3-4): 222-224.

Sharma, D.P., Patel, Manish and Nishith Gupta. 2006. Influence of nitrogen, phosphorus and pinching on vegetative growth and floral attributes in African marigold (*Tagetes erecta* Linn.) J. of Ornamental Hort. 9(1): 25-28.

Formatted: Highlight

Formatted: Font color: Red

Formatted: Highlight

Formatted: Highlight

Formatted: Indent: Left: 0", Hanging: 0.5", Space After: 0 pt, Line spacing: Double, Pattern: Clear (White)

UNDER PEER REVIEW

