

Original Research Article

Effects of ethylenediaminetetraacetic acid and some chlorides of divalent metals on the initial velocity of crude peroxidase from watermelon rind

ABSTRACT

Aim: This study investigated the effects of ethylenediaminetetraacetic acid (EDTA) and some chlorides of divalent metals on the initial velocity of crude peroxidase from watermelon rind

Study design: *In vitro* enzyme assay.

Place and Duration of Study: Department of Biochemistry, Faculty of Life Sciences, Ambrose Alli University, Ekpoma, Edo State, Nigeria between September 2023 to November 2023.

Methodology: The kinetics of the oxidation of Guaiacol by the crude peroxidase from the rind of watermelon in the presence of varying concentrations of EDTA and the chloride salts of iron, copper, magnesium and calcium was determined spectrophotometrically by monitoring the oxidation of Guaiacol to produce a brown tetraguaiacol at a wavelength of 470nm. The various salt concentrations were varied between 0.5mM and 3Mm. Each of the reaction mixtures used in the kinetic study comprised; 2.3 mL of 0.6 M sodium acetate buffer (pH 5.4), 0.2 mL of 0.02 mM Guaiacol, 0.1 mL of crude extract, 0.2 mL of varying concentration chloride salt, 0.2 mL of 2 mM of H₂O₂ added last to start the reaction. The final concentration of H₂O₂ in the 3 mL assay was 0.25 mM. The total volume of the reaction mixture was 3 mL. The absorbance was read every 2 seconds for one minute after adding hydrogen peroxide using a stop-clock. The control had no metal ion but replaced with distilled water

Results: Results showed that EDTA reduced the activity of the enzyme in a concentration dependent manner. CuCl₂ and FeCl₂ activated the enzyme and FeCl₂ being a better activator within the salt concentration range of 0.5mM to 3 mM.

Conclusion: These findings are of great importance to industries in understanding the mechanism of action of peroxidase from the rind of watermelon, especially as the search for cheap and alternative sources of peroxidases continues.

Keywords: [Watermelon Rind, Guaiacol, EDTA, Divalent metallic chloride]

1. INTRODUCTION

Peroxidases are oxidoreductases, they are produced by different plants and microorganisms. They have an iron porphyrin ring which catalyzes the oxidation of various organic substrates [1]. These ubiquitous oxidoreductases use hydrogen peroxide or alkyl peroxides as oxidants [2]. Peroxidases are considered thermo-stable enzymes [3]. However, they are readily inactivated by hydrogen peroxide [4]. Peroxidases have various applications such as in the synthesis of chemicals, medicine, and in the analysis of food, clinical and environmental

samples [5]. Peroxidase reactions can be monitored with the use of a compound called guaiacol (2-methoxyphenol). Guaiacol can be oxidized to produce a brown product called tetraguaiacol. This tetraguaiacol can be detected and quantified by a spectrophotometer at a wavelength of 470nm. [6]. Effects of Some Metal Chlorides on the Initial Reaction Rate of Crude Peroxidase from Watermelon Peels has been investigated previously [5]. This study aims at investigating the kinetics of peroxidase from the rind of watermelon using guaiacol as substrate. The effect of EDTA and some divalent metallic chlorides on the kinetics of peroxidase from watermelon rind could provide important information to aid their industrial application as alternative sources of peroxidase.

2. MATERIAL AND METHODS

2.1 Materials

Guaiacol, dimethyl sulphoxide, hydrogen peroxide (30 %), sodium acetate, acetic acid, disodium hydrogen phosphate, and sodium dihydrogen were purchased from SchauLab S.L. (Spain) and Loba Chimine Pot. Ltd. (India). Other reagents were all analytical grades and purchased from Sigma-Aldrich (Dorset, Poole, United Kingdom). All kinetic measurements were carried out using a UV-780 recording spectrophotometer.

2.2 Methods

2.2.1 Collection of Plants Materials

Watermelon (*Citrullus lanatus*) was purchased from a local market at Ekpoma, Esan West Local Government Area, Edo State, Nigeria. They were washed with distilled water in the laboratory.

2.2.2. Preparation of Crude Enzyme

10 g of rind from the watermelon fruit was weighed and washed with distilled water. This was then followed by homogenization in a blender using 100 mL of 0.1 M sodium phosphate buffer of pH 7.0. It was then filtered using a muslin cloth. The filtrate was centrifuged (Centrifuge 800B, Pec-Medical U.S.A) at 4000 rpm for 30 minutes. The supernatant was then decanted into a plain sample container and properly labeled as "crude extract". This was stored frozen in the freezer for analysis.

2.2.2 Estimation of Guaiacol Oxidation by Crude Peroxidase Isolated from watermelon rind with Salt Concentration

The kinetics of the oxidation of Guaiacol by the crude peroxidase from the rind of watermelon in the presence of varying concentrations of EDTA and the chloride salts of iron, copper, magnesium and calcium was determined spectrophotometrically by monitoring the oxidation of Guaiacol to produce a brown tetraguaiacol at a wavelength of 470nm. The various salt concentrations were varied between 0.5mM and 3mM. Each of the reaction mixtures used in the kinetic study comprised of: 2.3 mL of 0.6 M sodium acetate buffer (pH 5.4), 0.2 mL of 0.02 mM Guaiacol, 0.1 mL of crude extract, 0.2 mL of varying concentration chloride salt, 0.2 mL of 2mM of H₂O₂ added last to start the reaction. The final concentration of H₂O₂ in the 3mL assay was 0.25 mM. The total volume of the reaction mixture was 3mL. The absorbance was read every 2 seconds for one minute after adding hydrogen peroxide using a stop-clock. The control had no metal ion but replaced with distilled water

2.2.3. Determination of Initial reaction rate (Vo)

The initial reaction rate of the crude peroxidase from the rind of watermelon was determined by calculating the slope of the line for the first part of the data in the graph of absorbance versus time (i.e., Δ absorbance/second). The slope was then divided by the molar absorptivity for Guaiacol oxidation product ($\epsilon = 26,000 \text{ M}^{-1}\text{cm}^{-1}$), multiplied by the sample path length (1.00 cm for cuvette used). The result was expressed in mM/second. All assays were done in five replicates. The effects of varying concentrations of the salts were determined graphically using the mean values obtained per assay.

3. RESULTS AND DISCUSSION

Figure 1 shows the effect of varying concentrations of EDTA on the initial velocity (Vo) of crude peroxidase from rind of watermelon fruit in the oxidation of Guaiacol. Results showed that increasing the concentration of EDTA proportionately decreased the initial velocity (Vo) of the enzyme within a salt concentration range of 0.5 - 3 mM when compared with control which had no EDTA. Results from this study is similar to previous studies which showed that the inhibition of horseradish peroxidase activity by EDTA in iodide oxidation occurs in a concentration dependent manner [7]. Similarly, other studies [8] have also shown that EDTA is a concentration-dependent inhibitor of peroxidase from *P. stratiotes* leaf. EDTA usually results in inactivation of many metalloenzymes [9]. The observed reduction of the activity of peroxidase from rind of watermelon in a concentration dependent manner may be due to the ability of EDTA to sequester Fe^{3+} and Ca^{2+} which are required for structural and functional stability of the enzyme and thus its catalytic ability [8].

Figure 2 shows the effect of varying concentrations of FeCl_2 on the initial velocity (Vo) of crude peroxidase from rind of watermelon fruit in the oxidation of Guaiacol. Results showed that FeCl_2 significantly increased the activity of the peroxidase at low concentration (0.5mM) of the salt. However, progressively increasing the concentration of FeCl_2 up to 3mM proportionately decreased the activity of the peroxidase. The increase in peroxidase activity recorded at a concentration of 0.5 mM was similar to the findings in previous studies [10] where peroxidase activity increased significantly in gills, liver, kidney and brain of Fish (*Cirrhinamrigalaafter*) exposure to iron as compared to a control group. It was however observed that higher concentrations (>0.5mM) proportionately inhibited the activity of the peroxidase. This finding is similar to results from previous studies [11] which have shown that Fe^{2+} at a concentration range from 0 to 0.75 mmol/L improved the activity of bromelain, and the best Fe^{2+} concentration was found as 0.5 mmol/L. When the concentration of Fe^{2+} was higher than 0.75 mmol/L, the activity of the enzyme was inhibited, and the extent of inhibition was increased with increasing Fe^{2+} concentration.




Figure 1: Effect of varying concentrations of EDTA on the initial velocity of peroxidase from watermelon rind in the oxidation of Guaiacol




Figure 2: Effect of varying concentrations of FeCl₂ on the initial velocity of peroxidase from watermelon rind in the oxidation of Guaiacol.




Figure 3: Effect of varying concentrations of CuCl₂ on the initial velocity of peroxidase from watermelon rind in the oxidation of Guaiacol

Result in Figure 3 shows that increasing CuCl₂ concentration increased the activity of the peroxidase from the rind of watermelon when compared with the control with no metal ion. A peak average activity of the enzyme as a salt concentration of 1.5mM. Higher concentration of the salt led to reduced enzyme activity. Peroxidase activities has been previously shown to increase with increasing concentration of copper [12]

A comparative study of the effects of a low concentration (0.5 mM) of EDTA, CuCl₂ and FeCl₂ (Figure 4) on peroxidase activity from rind of watermelon shows that FeCl₂ is the most efficient activator of the enzyme at that concentration.




Figure 4: Comparative effect of 0.5 mM of EDTA, CuCl₂ and FeCl₂ on the initial velocity of peroxidase from watermelon rind in the oxidation of Guaiacol.

4. CONCLUSION

Results from this study have established the presence of peroxidases in rinds of the watermelon fruit. It has also been established that chlorides of Fe and Cu are activators of this peroxidase with Fe showing a better activating effect. EDTA was also shown to be an inhibitor of the enzyme. The similarity in the activities of the peroxidase from the rind of watermelon compared with other peroxidases in the presence of EDTA, Fe, and Cu is crucial in the search for alternative source of peroxidase.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

REFERENCES

1. Ali, H. and Shaker, K. (2023). extraction, purification and characterization of peroxidase from okra (*abelmoschus esculentus*). IOP Conference Series Earth and Environmental Science, 1158(9), 092009.
2. Kumar, A. (2024). Simple and efficient method for the extraction, isolation, and purification of peroxidase enzyme from wheat bran. *Biofuels Bioproducts and Biorefining*, 18(2), 425-438.
3. Al-Mentafji, H., Al-Fahdawi, M., & Al-farras, A. (2019). extraction, and purification of peroxidase enzyme from *peganum harmala* seeds. *International Journal of Drug Delivery Technology*, 9(04), 689-692.
4. Odilichukwu, O., Okpashi, V., Bayim, B., Oscar, E., Nwanguma, B., & Uzoma, N. (2021). Optimised isolation, partial-purification and profiling of peroxidase extracted from

saccharomyces cerevisiae of fresh palm wine. African Journal of Biological Sciences, 3(4), 88.

5. Iniaghe, O.M, Ibukun, O. Evaluation of the Effects of Some Metal Chlorides on the Initial Reaction Rate of Crude Peroxidase from Watermelon Peels. Asian Journal of Biotechnology and Bioresource Technology. 2023; 9(4): 44-49.

6. Mitsch, M, White, B. A Simple, Quantitative Peroxidase Assay Demonstrating Enzyme Inhibition with L-cysteine. Advances in Biology Laboratory Education. 2020. Publication of the Association for Biology Laboratory Education. 41 (44)

7. Enachi, E., Grigore-Gurgu, L., Aprodu, I., Stănciuc, N., Dalmadi, I., Bahrim, G., ... & Croitoru, C. (2018). extraction, purification and processing stability of peroxidase from plums (*prunus domestica*). International Journal of Food Properties, 21(1), 2744-2757.

8. Arise, R.O, Osundahunsi, B.O, and Adewale, A.A. Stability Profile of Pistia stratiotes Leaf Peroxidase. Nigerian Journal of Basic and Applied Science. 2018. 26(1): 96-105.

9. Peeters, H. Protides of the Biological Fluids. Proceedings of the Twenty-Ninth Colloquium 2014. 1981 (29). Elsevier.

10. Sabir, K., Javed, M., Latif, F, Ayub, M. Effect of Various Levels of Iron on Peroxidase Activity in the Fish, *Cirrhinamrigala*. 2018. Pakistan Journal of Zoology 13:13-18.

11. Meng, L., Yi-gang, Y., Xiao-yu, W., Bo-xi, Y., and Xing-long, X. Effects of iron (II) (Fe^{2+}) and iron (III) (Fe^{3+}) ions on the activity and stability bromelain. Modern Food Science and Technology. 2017. 33(8), 176–181.

12. Domingos, V.D., Boaro, C.S.F, Cataneo, A.C., Lobo, T.F. Martins, D. Effect of copper on the activity of pirogalol peroxidase in *Myriophyllum aquaticum* plants cultivated in nutritives solution. Planta daninha 2005; 23(2): 375-380