

# Efficacy of Ayurveda-based Ischemia Reversal Program in Management of Patients with Inducible Ischemia

## ABSTRACT

**Aim:** Despite comprehensive management recommendations for ischemic heart disease, the prevalence of ischemic heart disease continues to rise. Similar actions have been observed in various herbal drugs, making them promising targets for developing new therapeutic options for treating ischemic heart disease. The Ischemia Reversal Program addresses this issue by combining *Panchakarma* with nutrition therapy.

**Methods:** A retrospective, observational study was conducted at a single center in Maharashtra, India. The study duration was 2018 to 2022. A total of 84 patients aged 18 years and above diagnosed with inducible ischemic heart disease by stress test participated in the ischemia Reversal Program and were included in the study. Day 1, day 90, and 1 year 1 day data were compared.

**Results:** Mean age of the study population was  $58.83 \pm 12.29$  years. Males comprised 29 (34.5%) patients of the study population. Mean weight decreased significantly (day 1:  $66.43 \pm 11.71$  kg, day 90:  $61.04 \pm 9.99$  kg, and 1 year 1 day:  $61.95 \pm 10.92$  kg). Body mass index decreased significantly (day 1:  $25.87 \pm 4.55$ , day 90:  $23.84 \pm 3.87$ , 1 year 1 day:  $24.12 \pm 4.09$ ). Also,  $Vo_2$  max increased significantly (day 1:  $14.59 \pm 6.99$ , day 90:  $25.83 \pm 8.47$ , and 1 year 1 day:  $25.46 \pm 9.05$ ).

**Conclusion:** Ayurvedic *Panchakarma* in the form of the Ischemia Reversal Program showed significant short-term and long-term results in management of patients diagnosed with ischemic heart disease patients.

*Keywords:* Ayurveda, ischemic heart disease, Panchakarma, stress test

## 1. INTRODUCTION

Cardiovascular disease, predominantly comprising ischemic heart disease and stroke, is a leading cause of mortality in India. A National Mortality Study from 2000 to 2015 revealed that in the age group 30–69 years, 0.9 million cardiovascular deaths were attributed to ischemic heart disease. Furthermore, the probability of death due to ischemic heart disease in this age group increased from 10.4% to 13.1% in men and from 4.8% to 6.6% on women between 2000 and 2015 in India [1].

Pharmacotherapy alone is the mainstay treatment strategy, however poor adherence to medication and high costs significantly impedes effective and economical treatment of ischemic heart disease in India. Therefore, there is a need to explore novel therapeutic options to combat this disease [2]. The therapeutic role of drugs used in treatment of ischemic heart disease include balancing oxygen demand and supply to the heart, reducing blood pressure, decreasing platelet aggregation, exerting hypolipidemic action, and providing antioxidant effects [3]. Similar actions have been found in several herbal drugs, making them interesting targets for new therapeutic options for treatment of ischemic heart disease [2].

Ayurveda, a traditional Indian medicine system, is used by many physicians to treat various diseases. *Panchakarma* and allied therapies are integral to Ayurveda and provide relief for numerous diseases. The Ischemia Reversal Program (IRP) which combines *Panchakarma* and allied therapies is one such integrative approach for managing ischemic heart disease [4]. This study aims to assess the efficacy of Ayurvedic *Panchakarma* therapy combined with lifestyle modifications in the managing patients with inducible ischemia.

## 2. MATERIAL AND METHODS

### 2.1 Study Design and Patient Population

A retrospective, observational, single center study was conducted from 2018 to 2022 in Maharashtra, India. A total of 84 patients aged 18 years and above, diagnosed with inducible ischemia on stress test or 6 min walk test were included in this study. Patients with resting unstable angina, significant arrhythmia, or normal stress test according to the modified Bruce protocol were excluded from the study. The procedures were conducted in accordance with the Declaration of Helsinki [5], Good Clinical Practice [6], and applicable regulatory requirements.

### 2.2 Procedure

The IRP is a combination of *Panchakarma* and allied therapy. This therapy is performed on the patients after a light breakfast. One session of the procedure lasted approximately 65–75 mins. It is a 3-step procedure, the techniques used under this program are *Snehana*, *Swedana*, and *Basti*. *Snehana* is a massage in a centripetal manner with upper strokes. Further details are given in **Table 1 and Figure 1** and has also been described in previous literature [2]. The patients followed a diet plan of 1200 calories/day. The diet kits consumed are displayed in **Figure 2**.

**Table 1. Study treatment: Ischemia Reversal Program (IRP)**

Step	Type of therapy	Herbs used for therapy	Duration
Snehana	Massage	100 ml [80% Sesame oil + 20% Lavender oil]	30–35 mins
Swedana	Passive heat therapy	Dashmool (collection of 10 herbal roots) with steam at $\leq 40^{\circ}\text{C}$	10–15 mins + 3–4 mins of post procedure relaxation
Basti	Per rectal drug administration using a rectal solution	Like warm GHA decoction 100 ml	15 mins

**Figure 1. Graphical representation of Ischemia Reversal Program (IRP)**






Fig 2. Diet kit

### 2.3 Data Collection

Data for patient demographics, anthropometrics, Vo2 max and medications were collected from patient medical records. Data from day 1 were compared with data of day 90. Data of only those patients who had completed a total of 14 sessions was collected and analysed.

### 2.4 Statistical Analysis

All patient data were collected and coded in a Microsoft Excel sheet. Software R 3.4.4 was used to analyze data. Continuous data are expressed as the mean  $\pm$  standard deviation, whereas categorical data are expressed as number (frequency). Paired t-test was used to analyze the difference in various parameters at baseline and 90-day follow-up. A p-value  $<0.05$  was considered as statistically significant.

## 3. RESULTS

### 3.1 Patient demographics

The mean age of the patient population was  $58.83 \pm 12.29$  years. Males comprised 29 (34.5%) patients of the study population. Mean weight decreased significantly (day 1:  $66.43 \pm 11.71$  kg, day 90:  $61.04 \pm 9.99$  kg, and 1 year 1 day:  $61.95 \pm 10.92$  kg). Body mass index decreased significantly (day1:  $25.87 \pm 4.55$ , day 90:  $23.84 \pm 3.87$ , 1 year 1 day:  $24.12 \pm 4.09$ ). Vo2 max increased significantly (day 1:  $14.59 \pm 6.99$ , day 90:  $25.83 \pm 8.47$ , and 1 year 1 day:  $25.46 \pm 9.05$ ). The demographics are detailed in **Table 2**.

**Table 2.** Demographics of patient population

Variable	Day 1	Day 90	1 Year 1 Day	p value
Mean age, years		$58.83 \pm 12.29$		
Males, n (%)		29 (34.5%)		
Weight, kg	$66.43 \pm 11.71$	$61.04 \pm 9.99$	$61.95 \pm 10.92$	0.00
Body mass index	$25.87 \pm 4.55$	$23.84 \pm 3.87$	$24.12 \pm 4.09$	0.00
Vo2 max	$14.59 \pm 6.99$	$25.83 \pm 8.47$	$25.46 \pm 9.05$	0.00
Heart rate, bpm	$84.38 \pm 14.26$	$77.85 \pm 11.36$	$77.60 \pm 11.18$	0.00
Systolic blood pressure, mmHg	$132.00 \pm 18.33$	$128.54 \pm 20.09$	$117.19 \pm 27.79$	0.00

Diastolic blood pressure, mmHg      93.60 ± 24.37      79.27 ± 13.98      80.35 ± 13.06      0.00

*All data are expressed as number (percentage) or mean ± standard deviation.*

### 3.2 Vo2 max according to weight and body mass index

Vo2 max increased for low-risk (day 1: 24.98 ± 5.20, day 90: 32.55 ± 3.78 and 1 year 1 day: 31.26 ± 6.18), intermediate risk (day 1: 14.36 ± 1.74, day 90: 28.67 ± 7.31, and 1 year 1 day: 28.20 ± 8.19), and severe risk (day 1: 8.67 ± 1.89, day 90: 19.39 ± 6.83, and 1 year 1 day: 19.64 ± 7.76) patients. Weight according to Vo2 max decreased for low-risk (day 1: 69.39 ± 40.44 kg, day 90: 64.33 ± 8.21 kg, and 1 year 1 day: 65.07 ± 9.33 kg), intermediate-risk (day 1: 65.80 ± 10.43 kg, day 90: 59.76 ± 8.10 kg, and 1 year 1 day: 61.33 ± 8.96 kg), and severe-risk (day 1: 65.25 ± 13.11 kg, day 90: 60.23 ± 11.86 kg, and 1 year 1 day: 60.66 ± 12.83 kg) patients. Body mass index according to Vo2 max decreased for low-risk (day 1: 25.92 ± 4.86, day 90: 24.07 ± 3.93, 1 year 1 day: 24.17 ± 4.15), intermediate-risk (day 1: 26.29 ± 3.91, day 90: 23.87 ± 2.95, 1 year 1 day: 24.49 ± 3.43), and high risk (day 1: 25.48 ± 4.85, day 90: 23.68 ± 4.48, and 1 year 1 day: 23.76 ± 4.53) patients. The Vo2 max according to weight and body mass index is detailed in **Table 3**.

Vo2 max classification	Vo2 max			Weight, kg			Body mass index		
	Day 1	Day 90	1 Year 1 Day	Day 1	Day 90	1 Year 1 Day	Day 1	Day 90	1 Year 1 Day
Low-risk	24.98 ± 5.20	32.55 ± 3.78	31.26 ± 6.18	69.39 ± 40.44	64.33 ± 8.21	65.07 ± 9.33	25.92 ± 4.86	24.07 ± 3.93	24.17 ± 4.15
Intermediate-risk	14.36 ± 1.74	28.67 ± 7.31	28.20 ± 8.19	65.80 ± 10.43	59.76 ± 8.10	61.33 ± 8.96	26.29 ± 3.91	23.87 ± 2.95	24.49 ± 3.43
Severe-risk	8.67 ± 1.89	19.39 ± 6.83	19.64 ± 7.76	65.25 ± 13.11	60.23 ± 11.86	60.66 ± 12.83	25.48 ± 4.85	23.68 ± 4.48	23.76 ± 4.53

**Table 3. Vo2 max according to weight and body mass index**

*All data are expressed as mean ± standard deviation.*

### 3.3 Body mass index according to Vo2 max and weight

Vo2 max increased according to normal body mass index (day 1: 14.23 ± 7.45, day 90: 24.47 ± 8.37, and 1 year 1 day: 23.73 ± 8.55), overweight body mass index (day 1: 15.62 ± 7.17, day 90: 27.69 ± 8.76, and 1 year 1 day: 27.70 ± 9.61), obese 1 body mass index (day 1: 15.49 ± 4.19, day 90: 24.87 ± 7.31, and 1 year 1 day: 26.50 ± 6.96), obese 2 body mass index (day 1: 13.58 ± 1.11, day 90: 33.75 ± 1.95, and 1 year 1 day: 33.37 ± 1.90), and morbid body mass index (day 1: 13.08 ± 6.42, day 90: 24.89 ± 7.79, and 1 year 1 day: 23.79 ± 9.40). Weight decreased according to normal body mass index (day 1: 59.60 ± 6.52 kg, day 90: 56.24 ± 5.97 kg, and 1 year 1 day: 56.95 ± 6.94 kg), overweight body mass index (day 1: 65.55 ± 7.25 kg, day 90: 59.22 ± 6.29 kg, and 1 year 1 day: 60.36 ± 1.09 kg), obese 1 body mass index (day 1: 81.37 ± 8.29 kg, day 90: 73.50 ± 8.99 kg, and 1 year 1 day: 76.15 ± 8.79 kg), obese 2 body mass index (day 1: 83.27 ± 2.62 kg, day 90: 73.30 ± 2.97 kg, and 1 year 1 day: 73.57 ± 4.61 kg), and morbid body mass index (day 1: 84.41 ± 10.55 kg, day 90: 75.60 ± 11.10 kg, and 1 year 1 day: 75.80 ± 14.22 kg) patients. Body mass index decreased in normal body mass index (day 1: 22.62 ± 1.82, day 90: 21.43 ± 1.69, and 1 year 1 day: 21.69 ± 2.10), overweight body mass index (day 1: 26.20 ± 0.87, day 90: 23.77 ± 1.35, and 1 year 1 day: 24.06 ± 1.54), obese 1 body mass index (day 1: 29.17 ± 0.42, day 90: 26.37 ± 0.89, and 1 year 1 day: 27.24 ± 0.95), obese 2 body mass index (day 1: 30.95 ± 0.66, day 90: 27.25 ± 0.93, and 1 year 1 day: 27.34 ± 1.54), and morbid body mass index (day 1: 13.08 ± 6.42, day 90: 32.20 ± 4.02, and 1 year 1 day: 32.21 ± 5.11) patients. The body mass index according to Vo2 max and weight are demonstrated in **Table 4**.

**Table 4. Body mass index according to Vo2 max and weight**

Body mass index classification	Vo2 max			Weight, kg			Body mass index		
	Day 1	Day 90	1 Year 1 Day	Day 1	Day 90	1 Year 1 Day	Day 1	Day 90	1 Year 1 Day
Normal	14.23 ± 7.45	24.47 ± 8.37	23.73 ± 8.55	59.60 ± 6.52	56.24 ± 5.97	56.95 ± 6.94	22.62 ± 1.82	21.43 ± 1.69	21.69 ± 2.10
Overweight	15.62 ± 7.17	27.69 ± 8.76	27.70 ± 9.61	65.55 ± 7.25	59.22 ± 6.29	60.36 ± 1.09	26.20 ± 0.87	23.77 ± 1.35	24.06 ± 1.54
Obese 1	15.49 ± 4.19	24.87 ± 7.31	26.50 ± 6.96	81.37 ± 8.29	73.50 ± 8.99	76.15 ± 8.79	29.17 ± 0.42	26.37 ± 0.89	27.24 ± 0.95
Obese 2	13.58 ± 1.11	33.75 ± 1.95	33.37 ± 1.90	83.27 ± 2.62	73.30 ± 2.97	73.57 ± 4.61	30.95 ± 0.66	27.25 ± 0.93	27.34 ± 1.54
Morbid	13.08 ± 6.42	24.89 ± 7.79	23.79 ± 9.40	84.41 ± 10.55	75.60 ± 11.10	75.80 ± 14.22	35.92 ± 3.29	32.20 ± 4.02	32.21 ± 5.11

All data are expressed as mean ± standard deviation.

### 3.4 Medication adherence

Adherence to medications such as diuretics decreased by 80.95%, calcium channel blockers by 80.00%, antiplatelets decreased by 71.88%, statins by 65.22%, and beta blockers by 62.96% after 1 year 1 day. Patient adherence to medication is further elaborated in **Table 5**.

**Table 5. Medication adherence**

Medicine Name	Day 1 (n)	Day 90 (n)	1 Year 1 Day (n)	Day 1 to 90 (%)	Day 90 to 1 Year 1 Day (%)	Day 1 to 1 Year 1 Day (%)
Angiotensin-converting enzyme inhibitors	6	4	3	-33.33	-25.00	-50.0
Anticoagulant	51	51	42	0.00	-17.65	-17.65
Anti-platelets	32	21	9	-34.38	-57.14	-71.88
Beta-blockers	27	21	10	-22.22	-52.38	-62.96
Calcium channel blockers	15	6	3	-60.00	-50.00	-80.00
Diuretic	21	13	4	-38.10	-69.23	-80.95
Nitrate	21	15	11	-28.57	-26.67	-47.62
Statin	46	26	16	-43.48	-38.46	-65.22

## DISCUSSION

Despite extensive management guidelines for ischemic heart disease, its prevalence continues to rise. Conventional medicines address ischemic heart disease by correcting the imbalance between myocardial oxygen demand and supply, lowering blood pressure, reducing platelet aggregation, exerting hypolipidemic effects, and providing antioxidant benefits. Various herbal extracts have demonstrated similar properties, suggesting that Ayurveda could offer viable therapeutic options for managing ischemic heart disease. The IRP comprises 3 steps: *Snehana*, *Swedana*, and *Basti*. The proposed mechanisms of action include reduction of sympathetic over activity through *Swedana*,

decreased resting preload to reduce myocardial oxygen through *Swedana*, and nitric oxide release from the endothelium facilitated by a decoction of *TribulusTerrestris*, *Curcumin* and *Phyllanthusembelica*[2].

Ischemic heart disease is the leading cause of morbidity and mortality among women. This disease affects women and men at the same rate; however, the risk is greater among elderly women [1]. This observation is reflected in the current study as 65.5% of the study patients are women. A hypothesis for this may be that younger women especially, are subjected to bias that leads to lower rates of diagnosis and inferior management [7]. The Prospective Urban Rural Epidemiology (PURE) study has documented lower preventive management of cardiovascular risk factors in women compared to men [8]. Body mass index and high blood pressure are among other risk factors for ischemic heart disease. The mean body mass index decreased from  $25.87 \pm 4.55$  to  $23.84 \pm 3.87$  at day 90 and  $24.12 \pm 4.09$  after 1 year 1 day. Similarly, literature states that every 20 mmHg increase in systolic blood pressure doubles the risk of cardiovascular mortality [9]. In the current study systolic blood pressure decreased by 15 mmHg after 1 year 1 day.

Vo<sub>2</sub> max measures the maximum oxygen that can be utilized during exercise. Ischemic heart disease patients suffer from diastolic dysfunction, hence Vo<sub>2</sub> max is reduced in such cases which manifest clinically as reduced exercise/work capacity. In the present study, the Vo<sub>2</sub> max at the 90-day follow-up increased from  $14.59 \pm 6.99$  to  $25.83 \pm 8.47$  at the 90-day follow-up. This is in line with similar earlier studies that reported increase of Vo<sub>2</sub> max from  $20.29 \pm 6.72$  to  $29.40 \pm 6.71$  [2],  $20.74 \pm 7.25$  to  $29.69 \pm 6.62$  [4],  $17.82 \pm 7.23$  to  $26.65 \pm 6.14$  [10], and  $18.14 \pm 7.82$  to  $27.88 \pm 7.31$  [11] after 90 days.

## CONCLUSION

The study assesses the efficacy of Ayurvedic Panchakarma therapy combined with lifestyle modifications in the managing patients with inducible ischemia. Ayurvedic Panchakarma in the form of IRP therapy showed significant short-term and long-term results in management of patients diagnosed with ischemic heart disease patients.

## Study limitations

The study has a few limitations to consider. First, its retrospective design may limit the depth of insights compared to prospective studies, which could strengthen the existing predominantly retrospective literature. Second, the relatively small sample size restricts the generalizability of the study outcomes.

## Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

**Consent:**

All patients provided written informed consent for collection, analysis, and publication of their data.

**DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

**REFERENCES**

1. Ke C, Gupta R, Xavier D, Prabhakaran D, Mathur P, Kalkonde YV, et al. Million Death Study Collaborators. Divergent trends in ischaemic heart disease and stroke mortality in India from 2000 to 2015: a nationally representative mortality study. *Lancet Glob Health*. 2018 Aug;6(8):e914-e923.
2. Sane R, Sugwekar V, Nadapude A, Hande A, Depe G, Mandole R. Study of efficacy of ischemia reversal program (IRP) in ischemic heart disease (IHD) patients with VO<sub>2</sub>max and Duke's treadmill score. *International Journal of Basic & Clinical Pharmacology*. Aug 2018;6(8):1642-7.
3. Youssef G. Drugs used for treatment of ischemic heart disease. *Angiol Open Access*. 2021 9:258.
4. Sane R, Gond B, Raje G, Walzade k, Badre A, Mandole R. Ischemia Reversal Program (IRP) in patients suffering from ischemic heart disease (IHD) with known history of hypertension: A retrospective study. *Journal of Ayurveda Medical Sciences*. 2018;3(2):377-83.
5. General Assembly of the World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *J Am Coll Dent* 2014; 81(3):14–8.
6. Dixon JR Jr. The International Conference on Harmonization Good Clinical Practice guideline. *Qual Assur* 1998; 6(2):65–74.
7. Kiran G, Mohan I, Kaur M, Ahuja S, Gupta S, Gupta R. Escalating ischemic heart disease burden among women in India: Insights from GBD, NCDRisC and NFHS reports. *Am J PrevCardiol*. 2020 Jul 17;2:100035.
8. Walli-Attei M, Joseph P, Rosengren, Chow C, Rangarajan S, Lear SA, et al. Variations between women and men in risk factors, treatments, cardiovascular disease incidence, and death in 27 high income, middle-income and low-income countries (PURE): a prospective cohort study. *Lancet* 2020;396(10244):97–109.
9. Gupta R. Convergence in urban–rural prevalence of hypertension in India. *J Hum Hypertens* 2016; 30: 79–82.
10. Sane S, Wadekar A, Shinde K, Furia H, Upadhyay P, Mandole R. Understanding the role of ayurveda based Ischemia Reversal Program and low carbohydrate diet in reduction of risk of heart disease. *Asian Journal of Cardiology Research*. 2019;2(1):1-8.
11. Sane R, Ghadigaonkar P, Kharat A, Yadav KS, Mahajan S, Mandole R. Efficacy of Ischemia Reversal Program (IRP) in elderly patients of ischemic heart disease with known history of hypertension. *Asian Journal of Cardiology Research*. 2018;1(1):1-8.