

**“A study of insecticidal effects of leaf extracts of *Aegle marmelos* (L.) *Correa* and *Ricinus communis* (L.) on *Callosobruchus chinensis* (L.)”**

**Abstract:-**

Laboratory experiments were conducted to evaluate the efficiency of leaf powders of *Aegle marmelos* and *Ricinus communis* against the pulse beetle, *Callosobruchus chinensis*, which infests stored pulse seeds. Pulses specially stored for the purpose of human consumption should not be treated with toxic chemicals because residues of these chemical compounds may pose a serious risk to human health. Stored pulses are primarily attacked by a major insect pest particularly the pulse beetle, *Callosobruchus chinensis*. Though chemicals provide effective control of *Callosobruchus* sp., but the efficiency of the chemicals mostly depends on the mode of exposure. The maximum effectiveness is given when the insecticides are provided by gustatory method, but majority of these are toxic to human. Many synthetic insecticides have been found effective against stored product pests but proved to be hazardous to men and domestic animals. The over reliance on and ~~non-judicious~~ non-judicious use of synthetic pesticides especially insecticides since last four decades led to wide spectrum of pest problem like pest resistance to chemicals, resurgence of pests, residues in food and soil and risks to human and animal health, besides environmental pollution. The methanol and chloroform fraction of *Aegle marmelos* and *Ricinus communis* proved effective for the control of *Callosobruchus chinensis*. The findings suggest that plant extracted bio-pesticides may play a major role in insect pest control program in near future.

**Keywords:-**

*Aegle marmelos* and *Ricinus communis*, leaf powders, *Callosobruchus chinensis*, pulse protection, biopesticides.

**Introduction:-**

Pulses are an important source of nutrition for billions of people around the world. The terms “legumes” and “pulses” are interchangeable because all pulses are considered legumes but not all legumes are considered pulses. Pulses belong to leguminosae family and include those species that are consumed by human beings and domestic animals, commonly in the form of dry grains and does not include groundnut (*Arachis hypogaea*) and soybean (*Glycine max*) which are grown mainly for edible oil. The Food and Agriculture Organization

**Comment [U1]:** A format for abstract presentation is:

1. Introduction
2. Challenges/Problems
3. Aim
4. Methodology
5. Results
6. Conclusion
7. Recommendation/ suggestion

of the United Nations has declared 2016 as the International Year of Pulses with aims to enhance public awareness of the nutritional benefits of pulses as part of sustainable food production. This was intended to forward global food security and nutrition.

Pulses are rich source of many vitamins and minerals (iron, zinc, calcium, magnesium). The deficiency of various minerals in different parts of the world led to cardiovascular disease and imbalance in majority of the biological pathways. Pulses can provide adequate minerals required to fulfill nutritional requirement. Phenolic compounds are grouped into phenolic acids, tannins and flavonoids. Phenolic compounds and antioxidant activity also varied among different varieties of pulses.

Pulses are also being used by the processors worldwide for developing canned products as they are ready to use and the demand is expected to increase as they are having high shelf life (Warsame and Kimani 2014). Varieties with ease of preparation, processing efficiency and high yield of raw product were mostly preferred by the processors for thermal processing (Wassim et al. 1990; Hosfield et al. 2000).

Pulse varieties with uniform size and rapid expansion during soaking, high water holding capacity during processing and less splitting are required to be developed for canning. The effects of conventional processing methods (soaking, dehulling, boiling, pressure cooking, germination, fermentation etc.) on the levels of anti-nutritional factors (phytate, protein inhibitors, phenolics, tannins, lectins, saponins etc.) have been extensively evaluated, however, novel methods to eliminate these need to be developed (Patterson et al. 2017). Canning of pulses also reported to decrease phenolic content (Parmar et al. 2016).

Though the pulses are used in number of indigenous products but their functional attributes and potential health benefits have not been fully explored.

Pulses play a significant role in a country like India, and serve as an important part of the diet. They are considered to be as the “poor man’s meat” given that the consumption of dairy and animal products is relatively low among the poorest section of the country. They are also an important source of protein, minerals and fibre. The protein content in pulses is double the protein content of wheat, and three times more than that of rice. Additionally, pulses are used as green manure and contribute in improving soil health, offering the possibility for a mixed/ intercropping system (Rawal and Navarro, 2019).

Comment [U2]: Please, add a reference

Comment [U3]: A reference is needed here

Comment [U4]: Please, add a reference

Comment [U5]: Please, add a reference

Comment [U6]: Kindly adhere to the specified referencing style by this journal.

Comment [U7]: Join it to the previous paragraph

Comment [U8]: Insert a reference here

Pulses as a commodity group fit in all the Feed the Future Initiative themes aiming toward sustainable poverty and hunger reduction and enhancement of nutrition and health conditions alongside protection of the environment (Maredia, 2012).

Major pulses are grown chickpeas (Gram/Chana), Pigeon pea (Tur/Arhar), Mungbeans, Urdbeans (Blackgram), Masur (Lentil), Peas and various kinds of Beans (Minor pulses). The main regions with high productivity are Madhya Pradesh, Rajasthan, Maharashtra, Karnataka, Uttar Pradesh, Coastal Andhra Pradesh, Gujarat, Tamil Nadu, Jharkhand, Odisha, Chhattisgarh, Telangana, Bihar and West Bengal delta region.

The census projection report has further revealed that the proportion of the working age population between 15 and 59 years is likely to increase from 58% in 2001 to > 64 percent by 2021. Such a trend would make the country one of the youngest nations in the world. Thus, one of the India's competitive advantages is its demographic dividend.

Despite the imports, in 2019, the consumption of pulses in India amounted to 48 g per capita per day, slightly less than the 50 g per capita per day recommendation of the Indian Medical Research Council. One of the major hurdles in meeting self-sufficiency in pulses is policies that promote staple crop production, such as subsidies for fertilizers and credit and irrigation facilities that discourage the production of pulses and other legumes (FAO et al., 2020).

#### Material and Methods:-

#### Insect (*Callosobruchus chinensis*)-

*Callosobruchus chinensis* is a major pest of pulses in India. It is a holometabolic insect with the egg and adult stage found on grains and the pupal stages living inside the grain.

As the larva eats up the endosperm in most damaging stage of the life cycle. Adult beetles are 3-

4 mm in length, oval in shape, chocolate/reddish brown in colour, and have long erect antennae.

The female lays between 1-8 oval shaped and scale like egg grain. Each larva completes the life cycle in a separate chamber.

In India the insect breeds freely from March to November and hibernates in the larval stage during the winter. The adult emergence takes place from January to April. The pest causes maximum damage during February to August, when all its developmental stages exist simultaneously.

**Comment [U9]:** Join it to the previous paragraph

**Comment [U10]:** This statement is not linking to your review.

**Comment [U11]:** Your introduction should;  
1. mainly focused on *Aegle marmelos*, *Ricinus communis* and *Callosobruchus chinensis*  
2. Your justification for the experiment

**Comment [U12]:** Kindly re-cast this statement.

**Formatted:** Indent: First line: 0", Right: 0.62", Space Before: 0 pt

**Formatted:** Right: 0.62", Space Before: 0 pt

**Comment [U13]:** Add a reference

**Comment [U14]:** Add a reference

**Comment [U15]:** This explanation should be part of the INTRODUCTION. You only need to mention the insect's name and how they were collected under MATERIALS AND METHODS.

### Selection of plant-

*Aegle marmelos* and *Ricinus communis* was selected for the study because of its insecticidal and medicinal properties, and its abundance in the study area.

### Collection of plant material-

The indigenous plant *Aegle marmelos* leaves of family Rutaceae and plant *Ricinus communis* leaves of family Euphorbiaceae were collected from the roadside of **Laharpur Bhopal**, collection was done in winter season.

Comment [U16]: Laharpur

### Taxonomic position of the plant-

Identification of the collected plant was carried out in the Department of Botany of **the research center**.

Comment [U17]: ?

#### **Aegle Marmelos (Bael)-**

Kingdom-

Plantae Order

-

Sopindales Family -

Rutaceae

Sub family -

Aurantioideae Genus

-*Aegle*

Species-*marmelos*

Comment [U18]: Add reference

The biological and pharmacological activity against various chronic diseases such as cancer and cardiovascular and gastrointestinal disorders,

Comment [U19]: This statement is floating and incomplete.

Antioxidant, antiulcer, antidiabetic, anticancer, anti-hyperlipidaemic, anti-inflammatory, antimicrobial, anti-spermatogenic effects have also been reported on various animal models by the crude extracts of this plant. Every part of *Aegle marmelos*,

Comment [U20]: Kindly recast and add references.

Bael leaves could also be used as a potential biosorbent. **No** toxic lead ions were demonstrated to be removed from an aqueous solution by absorbing them into the bael leaves. An unusual fatty acid, 12-hydroxyoctadec-cis-9-enoic acid (ricinoleic acid), is present in the bael seed oil, which has the potential to be manufactured as biodiesel in the future. **S**tructure of bael exudate gum was also determined which could be further characterized for various industrial

applications. (Suvimol et al., 2008).

Comment [U21]: Recast

**Ricinus communis (castor):-**

Kingdom-

Plantae Division -

Magnoliophyta Class -

Magnoliopsida Subclass-

Rosidae

Order -

Euphorbiales Family -

Euphorbiaceae Genus-

*Ricinus*Species-*communis*

Comment [U22]: Add reference

The leaves ~~is~~ applied to boils and swelling, the hot leaves are applied over the abdomen of children to relieve flatulence. The leaves promote menstrual flow in women. Tender leaves ~~leaves are~~ cure pain in bladder. ~~the~~ leaves are also recommended to relieve headache and joint pains. (Jena J and Gupta, 2012.)

Comment [U23]: All these should be in the INTRODUCTION, not here.

For phytochemical analysis of the plants, the collected material after identification was used and a voucher specimen was procured in the herbarium sheet. The shade dried and powdered leaves were Soxhleted in 90% alcohol, rectified spirit and water respectively.

Comment [U24]: Put in parenthesis the voucher numbers

Comment [U25]: Do you mean successively or separately?

The extract thus obtained was kept in a glass vial and stored in the refrigerator; percentage yield was recorded in the tabular form (Table No-1). Percentage loss in weight after drying of the plant material was also recorded which showed 96.2% loss in weight.

Comment [U26]: Delete. This is not result section.

Comment [U27]: Delete. This is not result section.

The cold percolation method as mentioned by Abbott (1925) was followed for fresh plant leaves. The air dried plant material was powdered using electrical blender. 500 gm of powder was mixed with 300ml of n-hexane and kept for 72 hrs then filtered and it was stored in a reagent bottle. The powder was allowed to dry for 2hrs before pouring the other solvents-methanol and chloroform and kept for 72 hrs. The crude extracts thus obtained were filtered through Whatman's filter paper No. 1 and were evaporated to dryness in a rotary vacuum evaporator or water bath at the room temperature (40°C) and pressure (25-30 mmhg). Such dried and semisolid crude extract was stored in refrigerator the use i.e for bioassay test against *C. chinensis*.

Comment [U28]: ?

Comment [U29]: ?

Formatted: Superscript

Comment [U30]: ?

### Chemical analysis and identification of the compounds

Firstly, the crude extract of plant was defatted in n-hexane and extracted with methanol and chloroform. The concentrated solution was allowed to stand when a green yellow deposit was obtained.

For further identification and structural elucidation of plant extract, the purified sample was sent to SAIF, CDRI Lucknow for spectral analysis: where IR spectrum, UV spectrum, NMR and Mass spectrum were done.

On the basis of spectral data obtained from SAIF CDRI Lucknow and on comparing the data with authentic markers available finally, sesquiterpenelactone was identified.

### Study of *Aegle marmelos* and *Ricinus communis* leaf extract on *Callosobruchus chinensis*

The collection of insects was done from the infested grains.

20 beakers were taken. They were washed with distilled water and then dried. In each beaker 100 seeds of pulse (e.g. Urad bean) were taken.

In each beaker two drops of chemical: methanol and chloroform fractions of different concentrations e.g. 50 ppm, 100 ppm, 150 ppm, 200 ppm, 250 ppm were poured.

In each beaker 3 pairs of insects (3 males and 3 females) were introduced. The mouth of each beaker was covered with muslin cloth and tied with a rubber band and a control was kept.

After 7 days, the beakers were opened by removing the muslin cloth and the insects were taken out.

The eggs laid on the grains were counted and the beakers were covered with muslin cloths as before.

After 7 days, the beakers were opened for calculating the number of larva emerged from the eggs and then beakers were once again closed as before.

After another 7 days, once again the beakers were opened to calculate the number of pupae. Then beakers were once again closed with the muslin cloth.

Formatted: Line spacing: 1.5 lines

Formatted: Justified, Indent: Left: 0.56", Space Before: 0 pt, Line spacing: 1.5 lines

Formatted: Justified

Formatted: Justified, Right: 0.63", Space Before: 0 pt

Formatted: Right: 0.63", Space Before: 0 pt

Formatted: Justified, Right: 0.63", Space Before: 0 pt

After 7 days, again the beakers were opened to see the adult insects emerged from the pupae. They are recounted.

The seeds of each beaker were taken in a petri dish and were wetted with water to observe their germination.

These seeds which germinated were recounted.

The same experiment was repeated with the wrinkled seeds.

**Table No-1:**  
**Percentage loss in weight of the plant materials**

S. No.	Name of the plant	Wet weight of plant material in (gms)	Weight on drying of the plant material (gms)	Loss in weight on drying (gms)	Percentage loss in weight
1.	<i>Aegle marmelos</i> (Linn)	2000	180	1920	96.2%
2.	<i>Ricinus communis</i> (Linn)	2000	190	1932	98.4%

**Result and Discussion:-**

Hemanthapiriset al. (2021) reported highly significant repellency 96.91% and 98.99% of dried leaf and seed kernel extract of *C. thevetia* and *O. tenuiflorum* against red flour beetle after 24 hours of exposure which indicates that botanicals have high repellency properties. Likewise Gitahi et al. (2021).

According to Batish et al. 2008, Nerio et al. 2010. Several plant extracts have been well documented by researchers for both their insect repellent and insecticide effects. A study by Koon et al. (2007) showed that the use of impregnated jute bags with *Chenopodium ambrosioides* and aqueous extracts from *Lantana camara* significantly reduced damage to stored pulses of *Acanthoscelides obtectus* (Say) and *C. maculatus*. Those locally available plant extracts can be used as an inexpensive source of insecticide.

In the present investigation the repellent activity of *Aegle marmelos* leaf extract in five concentrations against *C. chinensis* at one hour interval showed maximum repellent activity (90%) in 250 ppm conc. of methanol and chloroform and minimum in control.

Formatted: Justified

Formatted: Justified, Right: 0.68", Space Before: 4 pt

Formatted: Justified, Right: 0.68", Space Before: 4 pt, Line spacing: 1.5 lines

Formatted: Justified, Line spacing: 1.5 lines

Formatted: Justified

Formatted: Left

Formatted: Font: Bold

Formatted: Normal, Indent: Left: 0.87", Space Before: 0.05 pt

Comment [U31]: Write them in full. However, how do they relate to your own plants?

Comment [U32]: How does this statement relate to your work?

In the present investigation the repellent activity of *Ricinus communis* leaf extract in five concentrations against *C. chinensis* at one hour interval showed maximum repellent activity (90%) in 250 ppm conc. of methanol and chloroform and minimum in control.

Aliyu and Ahmed (2006) reported that all non-edible oils caused larval and adult mortality to 100%, and also had ovipositional deterrence. The bruchids are controlled by oil extracted from some plant matter revealed that groundnut oil and *Mentha arvensis*(L), *M. spicata*(L), *M. piperata*(L) and *Cymbopogon nardus*(L) have an effect on bruchid.

In the present study of research work larval mortality was observed maximum in 250 ppm conc. of chloroform and methanol extract of *Aegle marmelos* and *Ricinus communis* leaf while it was minimum in the control.

Chakraborty et al., 2004 Based on total life cycle of *C. chinensis* on various pulses, the order of pulses were chickpea > black gram > green gram >. However, it is evident from the literature that higher seed weight and thick seed coat prolonged developmental period. In spite of higher nutrition, the development is retarded because of thick seed coat which creates obstruction for penetration of young larvae inside the seed and emergence of adult from the seed.

In the present study the leaf extract of experimental plant inhibited adult emergence. Minimum adults emerged in methanol and chloroform 250 ppm conc. and minimum in the control.

**Table No-2:**

**Statistical data of purified fraction of *Aegle marmelos* (AMF<sub>2</sub>)—methanol extract against pulse beetle (*Collosobruchus chinensis*) on fresh Urad bean seeds.**

Concentration	24 hr. larval mortality	Regression equation (y = a + bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	68	-3.669 ± 1.773 x	6.347	117.372	0.0243	0.156	L= 89.224 U = 147.830
100	92						
150	145						
200	158						
250	175						

Formatted: Indent: Left: 0.56"

Formatted: Font: Not Bold

Formatted: Heading 1, Justified, Space Before: 10.25 pt

Formatted: Heading 1, Justified, Indent: Left: 0", First line: 0", Right: 0", Space Before: 10.25 pt, Line spacing: single

Comment [U33]: Use either pulse beetle or *Collosobruchus chinensis*

Formatted: Font: Not Bold

TableNo.-3:

Statistical data of purified fraction of *Aegle marmelos*(ACF<sub>5</sub>) Chloroform extract against Pulse beetle (*Collosobruchus chinensis*) on fresh Urad bean seeds.

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	73	-3.775 ± 1.832 x	20.703	115.082	0.0243	0.156	L= 56.150 U=183.278
100	99						
150	119						
200	159						
250	195						
Control	6						

Formatted: Font: Not Bold

Formatted: Left, Indent: Left: 0.82"

Formatted: Heading 1, Indent: Left: 0.82", Right: 0.94", Space Before: 0 pt

Formatted: Indent: Left: 0.82", First line: 0"

Formatted: Font: Not Italic

Formatted: Font: Not Italic

TableNo.-4

Statistical data of purified fraction of *Aegle marmelos*( AMF<sub>5</sub>) methanol extract against Pulse beetle (*Collosobruchus chinensis*) on Wrinkled Urad bean seeds.

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	79	-3.82 ± 1.922 x	9.606	98.665	0.0246	0.157	L= 66.191 U =127.892
100	104						
150	149						
200	173						
250	197						
Control	6						

Comment [U34]: Kindly adjust other tables' titles.

**TableNo-5**

**Statistical data of purified fraction of *Aegle marmelos*( ACF<sub>5</sub>) Chloroform extract against Pulse beetle (*Collosobruchus chinensis*) on Wrinkled Urad bean seeds.**

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	71	-4.262 ± 2.109 x	11.656	104.906	0.0252	0.159	L= 71.848 U = 136.552
100	99						
150	142						
200	176						
250	199						
Control	6						

**TableNo-6**

**Statistical data of purified fraction of *Ricinus Communis*(RMF<sub>5</sub>) methanol extract against Pulse beetle (*Collosobruchus chinensis*) on fresh Gram/chickpea seeds.**

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	79	-3.873 ± 1.929 x	14.774	101.805	0.0246	0.157	L= 58.895 U=140.919
100	99						
150	144						
200	170						
250	199						
Control	9						

**TableNo-7**

**Statistical data of purified fraction of *Ricinus Communis*(RCF<sub>5</sub>) Chloroform extract against Pulse beetle (*Collosobruchus chinensis*) on fresh Gram/chickpea seeds.**

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	60	-5.018 ± 2.416 x	32.378	119.360	0.02689	0.164	L= 62.022 U=190.732
100	79						
150	128						
200	166						
250	209						
Control	9						

**TableNo.-8**

**Statistical data of purified fraction of *Ricinus Communis*(RMF<sub>5</sub>) methanol extract against Pulse beetle (*Collosobruchus chinensis*) on Wrinkled Gram/chickpea seeds.**

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	89	-4.317 ± 2.224 x	35.106	87.411	0.0259	0.161	L= 18.279 U=137.155
100	110						
150	149						
200	189						
250	223						
Control	9						

**TableNo.-9**

**Statistical data of purified fraction of *Ricinus Communis*(RCF<sub>5</sub>) Chloroform extract against Pulse beetle (*Collosobruchus chinensis*) on Wrinkled Gram/chickpea seed**

s.

Concentration	24 hr.larval mortality	Regression equation (y = a ± bx)	Chi - Square x <sup>2</sup> (n-1)	LC <sub>50</sub> (ppm)	Variance (V)	S.E.	Fiducial limits (ppm)
50	63	-4.880 ± 2.386 x	30.443	111.045	0.0268	0.164	L= 55.138 U=169.725
100	94						
150	134						
200	165						
250	216						
Control	9						

**References-**

**Aliyu, M** and Ahmed B I. 2006. Comparative efficacy of different rates of groundnut oil for the control of cowpea weevils *Callosobruchus maculatus*(F.) in stored cowpea (*Vigna unguiculata*(L) Walp). *Global Journal of Agricultural Sciences* 5(2): 123–26.

**Abbott, W.S.** (1925). A method of computing the effectiveness of an insecticide. *J. Eco. Entomol.*, 18:265-267.

**Batish, D R, Singh H P, Kohli R K and Kaur S.** 2008. Eucalyptus essential oil as a natural pesticide. *Forest Ecology and Management* 256(12): 2166–74.

**Chakraborty, SN, Chaudhar and Senapati. SK.** Correlation between seed parameters and relative susceptibility of mungbean genotypes (*Vigna radiata* L.) to *Callosobruchus chinensis* L. during storage. *Annals of Plant Protection Science* 2004;12(1):48-50.

**FAO, /IAEA.** Guidelines for Mass Rearing Aedes Mosquitoes. Version 1.0; IAEA: Vienna, Austria, 2020.

**Gitahi, SM, Piero MN, Mburu DN, Machocho AK.** Repellent effect of selected organic leaf extracts of *Tithonia diversifolia* (Hemsl.) A. Gray and *Vernonia lasiopus* (O. Hoffman) against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae). *Hindawi. The Scientific World Journal*, 2021. <https://doi.org/10.1155/2021/27718629>.

**Hosfield, GL, Uebersax MA, Occena LG (2000)** Technological and genetic improvements in dry bean quality and utilization. In: Proceedings of the Idaho Bean Workshop, University of Idaho, Moscow, 135–152.

**Hemantha, Piris BGB, Pakeerathan K, Sayanthan S.** Efficacy of botanical extracts against storage insect pests *Tribolium confusum* (Confused flour beetle) and *Sitophilus oryzae* (Rice weevil). Proceedings of 1st International electronic conference of entomology, 2021, 1-15.

**Jena, J and Gupta AK (2012).** *Ricinus communis* Linn: A phytopharmacological Review. Inter. J. Pharm. Pharmaceut. Sci., 4(4): 25-29.

**Koona, P, Tatchago V and Malaa D. 2007.** Impregnated bags for safer storage of legume grains in West and Central Africa. Journal of Stored Products Research 43(3):248–51.

**Maredia, M., 2012.** Global pulse production and consumption trends: the potential of pulses to achieve 'Feed the Future' food and nutritional security. In: Presented at Global Pulse Researchers Meeting, Rwanda, vols. 13–19. February.

**Nerio, L S, Olivero-Verbel J and Stashenko E. 2010.** Repellent activity of essential oils: a review. Bioresource Technology 101(1): 372–78.

**Parmar, N, Singh N, Kaur A, Viridi AS, Thakur S (2016)** Effect of canning on color, protein and phenolic profile of grains from kidney bean, field pea and chickpea. Food Res Int 89:526–532 .

**Patterson, CA, Curran J, Der T (2017)** Effect of processing on antinutrient compounds in pulses. Cereal Chem 94:2–10.

**Rawal, V and Navarro DK eds. 2019.** The Global Economy of Pulses. Rome, FAO.

**Suvinol C., Pranee, A., Suvinol, C., Pranee, A., (2008):** Bioactive compounds and volatile compounds of Thai baelfruit (*Aegle marmelos* L.) Correa) as a valuable source for functional food ingredients. International Food Research Journal., 15(3): 1-9.

**Warsame, AO, Kimani PM (2014)** Canning quality of new drought tolerant dry bean (*Phaseolus vulgaris* L.) lines. Am J Food Tech 9(6):311–317.

**Wassimi, NN, Hosfield GL, Uebersax MA (1990)** Inheritance of physico-chemical seed characters related to culinary quality in dry bean. J Am Soc Horticult Sci 115:492–499.