

Review Article

Effect of biological priming on metabolomic and molecular changes in response to drought stress in *Brassica juncea*

Abstract

Indian mustard (*Brassica juncea*) is a leading oilseed crop in India and plays a crucial role in the agricultural as well as oilseed home marketing system of the country. High oil and phytosterols, glucosinolate content make it a more valuable crop, and it is used for edible oil production, poultry feed. There, however, is always one major problem during the growth of trees and their productivity; drought. This paper discusses the use of bio-priming to compound drought resistance in Indian mustard. Bio-priming, which entails the use of various microorganisms in seed treatment is therefore modern system of farming. In the process of germination and plant growth it enhances the seed germination rate, seedling vigor and overall plant health through various metabolomic and molecular mechanisms. Works have demonstrated that bio-priming containing elements such as *Pseudomonas fluorescens*, *Bacillus subtilis*, and *Trichoderma harzianum* facilitates the uptake of sulfur, growth and improvement of tolerance to abiotic stresses. Furthermore, by impacting on expression of stress associated genes, bio-priming enhances the activity of the WRKY transcription factor as well as the production of abscisic acid (ABA) and jasmonic acid (JA), leading to drought tolerance. These first results based on drought tolerance genes help to priorities future breeding programs dedicated to the improvement of drought resistant cultivars. Bio-priming also stimulated the antioxidant defense pathway and improved the ability of plants to cope with oxidation pressure arising from drought stress. Overall, bio-priming is a relatively cost-effective, environmentally considerate method for enhancing drought tolerance in Indian mustard, thus being beneficial for sustainable farming and food production.

Keywords: abiotic stress, bio-priming, Brassica juncea, drought, metabolomics.

INTRODUCTION

Indian mustard (*Brassica juncea*), a member of the Brassicaceae family, is an important oilseed crop in India and plays a vital part in the country's agriculture. With an oil content ranging from 38% to 50% (Shakeel et al., 2019), it is highly valued and accounts for about 80% of the nation's total rapeseed-mustard production (Rajput et al., 2018). This important crop during the Rabi season is produced through interspecific hybridization and has a high oil content (37–42%) (V. V. Singh et al., 2020). India is a major producer of rapeseed-mustard, which is the third most important oilseed crop in the world and makes a considerable contribution to the oilseed economy of the nation (S. Kumar et al., 2020). Mustard is well known for being high in bioactive substances that have anti-inflammatory and antioxidant qualities. It also contains phytosterols, phenolic compounds, glucosinolates, and anti-carcinogenic substances (Theertha, et al., 2023). India, which lies in third place behind China and Canada in the world's mustard production scenario, is an important participant. Over 6.23 million hectares of mustard are cultivated in India, producing 9.34 million tons at a productivity of 1499 kg/ha (Bhardwaj et al., 2020). With 5.96 million hectares under cultivation and 8.32 million tonnes produced annually at a yield of 1397 kg/ha, the nation provides 28.3% of the world's rapeseed production (Basavanneppa & Kumar, 2020). In terms of the economy of oilseeds in India, mustard is grown in states such as Rajasthan, Uttar Pradesh, Madhya Pradesh, Punjab, Haryana, West Bengal, Assam, and Bihar (Sahu et al., 2020). It ranks second behind groundnut. Rajasthan is a prominent mustard-producing state that contributes significantly to India's overall mustard production, with Madhya Pradesh and Haryana following closely after (Basavanneppa & Kumar, 2020). Furthermore, while mustard is produced in other states in India, including Gujarat, Andhra Pradesh, Maharashtra,

Karnataka, and Uttar Pradesh, Madhya Pradesh makes up a substantial portion of the country's GDP (Shukla & Gupta, 2020). Drought is one of the major obstacles to Indian mustard growth and development, affecting yield and productivity. India's agriculture is highly dependent on its climate, which leaves it open to water scarcity and decreased crop growth as a result of weak monsoons (Nivetha et al., 2020). Drought-proofing techniques are therefore required because of its reduced ability to grow and produce seed in drought conditions (Singh et al., 2018). The significance of drought tolerance indices in the selection of high-yielding and drought-tolerant cultivars, such as PusaJaikisan, which demonstrated resilience to drought stress while sustaining high yields, has been highlighted by research on a variety of mustard genotypes (Kumari et al., 2020). In rainfed regions that are vulnerable to drought, increasing mustard's tolerance to water stress through plant breeding and agronomic techniques is essential for long-term productivity (Sahoo et al., 2019). Seed priming is a new generation technique that helps to resist the plants from abiotic stresses like drought. Priming has various approaches, among which there is a method where we use living micro-organisms such as bacteria, fungi, called as bio-priming (Kępczyński & Kępczyńska, 2023). Biological priming improves a plant's resistance to a range of stresses by inducing particular metabolomic reactions and molecular pathways (Irshad et al., 2023).

Biological priming

Priming is a conventional technique that is typically employed for solid crop stands and synchronized seedling growth, but in recent years, it has become a powerful tool for sustainable agriculture (Paul et al., 2022). There are different priming methods that effect the germination pattern of mustard such as hydro-priming, osmo-priming, hormonal priming and bio-priming (Thapa et al., 2022). Bio-priming, a seed treatment approach which combines a physiological (seed hydration) and biological (seed inoculation with a beneficial organism in order to protect seed) component of disease management is a novel seed treatment method (Reddy, 2012). By improving the adherence of bacteria to seeds, seed priming with living inoculums enhances rhizosphere colonization and increases plant resistance to unfavourable environmental conditions (Prasad et al., 2016). Biopriming, a practice where seeds are treated with biocontrol agents together with priming agents, has been used to control a number of soil- and seedborne diseases (Reddy, 2012). Chemical fungicides are not the only option for growers to control seed and seedling diseases; biological seed treatments offer an alternative option for this and hence eco-friendly and sustainable. Various bioinoculants such as *Pseudomonas fluorescens* and *Bacillus subtilis* are used as biopriming agents to increase the sulphur Use Efficiency in *Brassica juncea* (Singh et al., 2023). Other than these bioagent *Trichoderma asperellum* Th-14 is also use for promoting the plant growth in Indian mustard (Rawat et al., 2022). Mycorrhizal fungus-assisted biopriming of mustard plants has demonstrated encouraging outcomes in terms of improving plant growth and disease resistance. Studies have shown that the use of mustard oil cake, *Pseudomonas fluorescens*, and mycorrhizal fungi like *Glomus mosseae* and *Scutellospora erythropa* can successfully treat root-rot disorders brought on by pathogens like *Macrophomina phaseolina* (Neeraj & Singh, 2010). To increase the rate and uniformity of seed emergence and reduce numerous seedborne diseases, seed priming has been employed either alone or in conjunction with suitable fungicides and/or biocontrol agent (Sood et al., 2021). It has been demonstrated that biological priming, which involves soaking seeds in different substances such as gibberellic acid (GA), salicylic acid (SA), and abscisic acid (ABA), greatly increases the ability of plants such as rapeseed to withstand drought stress (Holness et al., 2023). The processes involved in bio priming in plants help to improve seed germination, seedling vigour, and overall plant growth. When plant growth-promoting microorganisms (PGPMs) are used in the biopriming process, various mechanisms are triggered (Deshmukh et al., 2020). These include the production of siderophores, the solubilization or mobilisation of soil nutrients, the induction of plant growth-promoting activities, and the synthesis of defense-related enzymes, phytoalexins, and beneficial biochemicals (Rhaman et al., 2020). Additionally, by promoting the synthesis of growth regulators, boosting nutrient uptake, and shielding plants from pathogens, bio priming with PGPMs like PGPR can result in increased seed quality, germination speed, vigour index, growth promotion, and disease resistance (Mitra et al., 2023).

Procedure for seed biopriming:




Figure 1: Process of Biopriming of mustard seed.

List of bio-priming agents used in mustard:

Table 1: bio-priming agents used in mustard

SL.NO.	BIO-PRIMING AGENTS	USES/BENEFITS	REFERENCE
1.	<i>Bacillus subtilis</i> and <i>Pseudomonas fluorescens</i>	It enhancing sulphur use efficiency and promoting sustainable production	(S. Singh et al., 2023)
2.	<i>Bacillus amyloliquefaciens</i> D5 ARV	Helps in improving germination rates when combined with a static magnetic field treatment to revive white mustard seed	(Jovičić-Petrović et al., 2021)
3.	<i>Trichoderma harzianum</i> and <i>Pseudomonas fluorescens</i>	Helps in managing pests and diseases in mustard crops and contributes increased yield and plant health	(Chauhan et al., 2018)
4.	<i>Rhizobium</i>	Rhizobium with vermiwash, neem leaf	(Sree et al., 2022)

		extract, tulsi leaf extract used. Rhizobium (10%) showed maximum growth and yield in yellow mustard	
5.	<i>Coniothyriumminitans</i> , <i>Aspergillusnidulans</i> , <i>Trichoderma harzianum</i> , <i>Pseudomonas chlororaphis</i>	It is effective in reducing carpogenic germination in Indian mustard	(Gupta, 2016)
6.	NSKE, <i>Metarhizium anisopliae</i> 2x10 ⁹ CFU and <i>Beauveria bassiana</i> 2x10 ⁹ CFU	Used as insecticide and protect against aphids in mustard	(Nayak et al., 2014)

Phases of priming and effects:

It is crucial for priming to involve a series of environmental stimuli (Tenenboim & Brotman, 2016). The process is initiated with a triggering stimulus and goes up to the time when the plant is exposed to a challenging stress called priming phase (Hilker et al., 2016). During this phase, the plant is kept in a state of readiness for attack because the levels of main and secondary metabolites, phytohormones, SA (salicylic acid), and JA (jasmonic acid) fluctuate slightly (Balmer et al., 2015). Stress-hardened plants move to the second level of stress-hardening known as post-challenge primed state through a second stress which triggers defence response mechanisms promptly (Shi et al., 2016). This includes de novo production of antimicrobial chemicals. A primed plant can revert to naïve state, however, if it inherits primed status from the parent plants, then plants can be trans generationally primed (Tugizimana et al., 2018).




Figure 2: Phases of Priming

Metabolomic and molecular changes due to bio-priming in mustard:

Priming can be caused by natural or artificial substances such certain agrochemicals, or by interactions between the host plants and helpful microbes (rhizobacteria, mycorrhizal fungi), diseases,

or virulent/avirulent microorganisms and plants that have these kinds of interactions experience long-lasting cellular and organismal reprogramming, and they molecularly "remember" these interactions (Tugizimana et al., 2018). Bio-priming in Indian Mustard have metabolomic changes as it involves seed priming, research reveals that the use of bioinoculants such as *Bacillus subtilis* and *Pseudomonas fluorescens* promotes sulfur usage and its uptake with increased levels of S content and enzymes responsible for S mineralization (S. Singh et al., 2023). Sulphur increases the amount of glucosinolate and the proportion of oil content, which is essential for the production of oil (Saleem et al., 2019). A lack of S causes an imbalance in the intake of nutrients, which in turn causes the loss of chlorophyll, stunted growth, and decreased agricultural yields (Waraich et al., 2021). Hence *B. subtilis* and *P. fluorescens* promotes the sulphur level that increases the oil content in mustard. Moreover, seed born plant hormones like GA3 and SA play vital role in enhancing several bio-chemical attributes of mustard crops including chlorophyll process and oil percentage in seed (Tulsi et al., 2018). Studies on growth promotion, disease suppression and biochemical changes in response to bio control agents like *Trichoderma harzianum*, *Pseudomonas fluorescens*, *Bacillus subtilis* were shown to affect growth, disease and tissue biochemical composition like dry matter, phenols, sugars, lipids in the leaves of Indian mustard and alter protein quantity (S. Sharma et al., 2010). Treatment of tomato plants with *Bacillus subtilis* against *Fusarium oxysporum* has been shown to boost lipid content (Ghonim et al., 1999). There is a demonstration after an experiment that in all cultivars of mustard, the total phenol concentration decreased after reaching a maximum at 60 DAS in healthy leaves and 80 DAS in ill leaves (Gupta et al., 1990). Furthermore, the studies on transgenic technology in *B. juncea* have also given fruitful results regarding the management of fatty acid environment in the seed, and also changing the nature of the seed itself to increase the yield of the oil (Sinha et al., 2007). The main aim of the present study involving two candidate genes was to reduce the VLCUFA (i. e. , C22: To this end, the levels of one of the undesirable fatty acids, namely, erucic acid, in the seed-oil of one Indian mustard cultivar (cv.) *B. juncea* cv. PCR7, a high-yielding variety and resistant to lodging and pod-shattering were altered through genetic engineering to change the carbon flux in the metabolic pathway of fatty acid biosynthesis accompanied by the corresponding changes in the expression of the respective First, the effects of over-expression of the novel acyl-ACP thioesterase of FatB originated from the MbFatB gene from *Diploknema* (*Madhuca*) *butyracea* on the VLCUFA content were examined in order to shorten the acyl chain-elongation process and increase C16 and C18 FAs in the seed-oil (Jha et al., 2006). This study concluded process to reduced erucic acid, increased beneficial fatty acids, and enhanced oil content. It has been noted that using plant growth-promoting bacteria in biopriming techniques is an eco-friendly way to increase seed germination in a variety of crops, highlighting the significance of microbial interactions in raising germination rates and seedling vigor (Rahaman et al., 2022). The studies using RAPD markers established that various mustard lines and hybrid have distinct genetic patterns suggesting diversified genetic base and have prospect of utilization in breeding program (R.A et al., 2013). In addition, polymorphic amplicons, alleles, and gene diversity among several Indian mustard genotypes have been discovered by genetic diversity assessments employing SSR markers, offering insights for the crop's genetic development (Shrivastav et al., 2023). The biochemical and molecular properties of Indian mustard, such as the concentration of sinigrin, the isozyme patterns of peroxidase and esterase, the intensity of seed protein bands, and the genetic diversity determined by SSR and ISSR markers, can all be affected by biopriming (Chaudhary et al., 2014). Furthermore, the cloning and function identification of a promoter in Indian mustard has shown strong expression activity under various stress conditions, suggesting a potential role in gene regulation for stress responses (Lang et al., 2017). When Indian mustard genotype (RH 781) was inoculated with *Albugo candida* led to a significant increase in ascorbic acid content, and the most significant changes in metabolite levels (ascorbic acid and glutathione) were observed in the 3rd leaf of both genotypes after 3 days of inoculation, with minimal changes in the 4th leaf compared to healthy leaves and other inoculated leaves on all days of inoculation (SAPNA1, 2011). Indian mustard's molecular alterations, which impact the genotypes' genetic variability and biochemical composition, are influenced by biopriming.

Effect and response to drought due to bio-priming on Indian mustard:

Drought modifies a number of physiological and anatomical factors that are essential for mustard crop physiology to adapt to the drought. Drought stress causes mustard plants to lose fresh weight, dry weight, and abscisic acid content (Fang et al., 2022). Drought stress also alters the protein composition, chlorophyll content, and activity of enzymes like catalase and peroxidase in mustard, resulting in physiological alterations that impact drought resistance (Zhu et al., 2021). Moreover, osmotic adaptations to display drought tolerance preserve the turgidity of cells in Brassica species, including mustard, emphasizing the significance of physiological features in maintaining resistance to water stress (Chandra & Kumar, 2022). Biopriming with drought-tolerant isolates such as *Trichoderma harzianum*, improves osmoregulation and induces physiological protection against oxidative damage in Indian mustard (Sahoo et al., 2019). *Trichoderma* as a seed treatment before sowing can induce tolerance to water stress in plants, highlighting its potential as a sustainable approach to improving crop resilience and could thrive in water stress conditions and positively impact mustard plant growth under such stress (K. K. Sharma & Singh, 2014). By delaying changes in net photosynthesis, stomatal conductance, and chlorophyll fluorescence brought on by drought, this strategy eventually makes plants more resilient to water stress. It has also been discovered that using bioregulators such as glycinebetaine and salicylic acid can improve drought tolerance and growth in mustard crops by lessening the detrimental effects of water stress (A. Singh & Meena, n.d.). Moreover, it has been noted that induced drought circumstances cause mustard types to mount a defence response. This reaction results in elevated antioxidant, detoxifying enzyme, and biochemical parameter levels that improve resistance to drought stress and Alternaria blight infection (Mallick., 2017). Furthermore, it has been researched that pre-treating mustard seedlings with hydrogen peroxide activates defense mechanisms, increasing membrane stability and resistance to drought-induced oxidative damage (Anwar, Hossain et al., 2013). It has been shown that biological priming—pre-treating plants like Indian mustard with hydrogen peroxide or β -aminobutyric acid (BABA)—increases their resistance to drought. Research has indicated that the use of hydrogen peroxide priming can initiate defense mechanisms, initiate ROS and MG detoxification processes, and sustain essential enzyme functions and redox equilibrium, thereby equipping seedlings to endure drought-induced oxidative stress (Rasul et al., 2021). Similarly, BABA administration has been associated with increased accumulation of ABA, which causes stomatal closure and stress gene expression to accelerate, improving plant resistance to drought stress without causing genetic changes (Fleming et al., 2019). Indian mustard genetic expression linked to drought resistance and tolerance has been demonstrated to be positively impacted by bio priming, more especially by the use of biostimulants such as *Ascophyllum nodosum* extract. It has been shown that the application of biostimulants can suppress stress-responsive negative growth regulators, like RESPONSIVE TO DESICCATION 26 (RD26), while preserving the expression of markers of the cell cycle, such as HISTONE H4 (HIS4), to facilitate the active progression of the cell cycle under drought stress (Holness et al., 2023).

Genes responsible for drought tolerance in Indian mustard

Various genes are that are either already present or incorporated in plant can result in drought tolerance.

Table 2: Different genes responsible for drought tolerance in mustard

Sl.No.	Gene	Importance	Reference
1.	BjRD26	A NAC transcription factor gene (NAM, ATAF1/2, and CUC2) implicated in the regulation of genes associated to stress and the response to drought stress.	(Tripathi et al., 2017)
2.	BjHSP17.8	This gene helps prevent damage to cellular proteins by encoding a little heat shock protein that is produced during	(Yadav et al., 2019)

		drought stress.	
3.	BjDREB2A	A transcription factor gene known as dehydration-responsive element-binding (DREB) that controls the expression of genes sensitive to stress in drought-prone environments.	(Luo et al., 2018)
4.	BjERF1	A gene that responds to ethylene and is involved in the regulation of genes that are sensitive to drought.	(Divya et al., 2010)

The genes responsible for drought tolerance in Indian mustard (*Brassica* species) have been identified through various studies. Several unique genes involved in drought tolerance have been identified by comparative transcriptome analysis, including genes related to photosynthesis, biosynthesis of glutathione, biosynthesis of IAA signal transduction, biosynthesis of amino acids, cysteine, and methionine, and glucosinolate (Kumari vedna, 2020). These genes are involved in the molecular aspects of drought stress in *Brassica* species and also useful in the selection of drought-resistant cultivars for the improvement of drought tolerance in breeding programs (Sharma et al., 2022). Further, genes associated with protein phosphatase 2C have been reported to be involved in drought stress and hence, it becomes important to know the genetic makeup of drought tolerance in Indian mustard (Zhu et al., 2022). IgWRKY50 and IgWRKY32, the genes of *Iris germanica*, are two genes which were found to be important for increasing drought resistance in Indian mustard (Zhang et al., 2022). The genes belonging to this superfamily of WRKY transcription factor are regulated by kind of stress conditions including PEG-6000, high temperature and ABA and reaches at its maximum level at 3 h in case of PEG-6000 treated plants, when overexpressed in both *Arabidopsis* plants, enhanced shooting with improved osmotic tolerance, higher germination rates, increased average root length, and increased stomatal closure and reduced water loss rates under drought stress conditions (Zhang et al., 2022). Moreover, the expression of the stress-responsive genes in the transgenic plants was also upregulated, proline content and soluble protein content in the plants also increased, soluble protein content in the plants also increased, together with SOD, POD and CAT activities, all of which accumulated in the transgenic plant to enhance the drought resistance of the plants (Kumari et al., 2019). Therefore, IgWRKY50 and IgWRKY32 genes could be prospective candidates for effective utilization in molecular breeding programs directed at improving DT in Indian Mustard. Transgenic mustard lines demonstrated increased antioxidant enzyme activity, decreased oxidative stress markers, and improved physiological traits, leading to better survival and performance under drought conditions. Over-expression of the chickpea Metallothionein1 (MT1) gene in mustard significantly enhances drought tolerance (Lal et al., 2024).

Effect of bio-priming on the expression of genes involved in drought tolerance in Indian mustard

Research have shown that bio-priming upregulates stress-responsive genes and signalling pathways, including those involved in ABA and JA production, which in turn affects the expression of genes linked to drought tolerance (Wang, Li, et al., 2021). Furthermore, it has been discovered that bio-priming preserves the shoot apical meristem's function and encourages active cell cycle progression during drought stress, which enhances plant resilience and growth (Rasul et al., 2021). Through the

activation of stress memory systems, reinforcement of cellular defence responses, and induction of phytohormone synthesis, bio priming plays a critical role in improving drought tolerance in plants. Research has demonstrated that, in drought-stressed environments, seed biopriming with bacteria resistant to drought can greatly increase crop yields (Lastochkina et al., 2023). Numerous seed priming methods, such as bio-priming with bacteria that promote plant growth, have been found to be effective ways to increase crops' resistance to stress, which will boost growth, dry matter accumulation, and overall plant performance in both ideal and drought-prone environments (CIG et al., 2022). It has been discovered that bio priming with particular bacterial strains raises antioxidant levels, including those of catalase, superoxide dismutase, and ascorbic peroxidase, all of which are essential in reducing the harmful consequences of oxidative stress brought on by dehydration (H. Kumar et al., n.d.). Additionally, it may result in the activation of genes linked to plant hormone signalling, cell wall remodelling, and stress defence, all of which improve the plant's overall drought tolerance in Indian mustard and increase its capacity to fend off the impacts of drought (Wang, et al., 2021). CabHLH10, a bHLH transcription factor that regulates yield attributes under drought stress by altering the expression of drought-responsive genes and photosynthetic efficiency genes, is one of the specific genes bio-primed for better drought tolerance in Indian mustard (Thakro et al., 2023). The development of drought-tolerant cultivars of Brassica species has also been aided by the identification of traits such as primary branches plant-1, secondary branches plant-1, siliquae plant-1, seeds siliqua-1, seed yield plant-1, and 1000-seed weight as trustworthy selection criteria for drought tolerance (Sharma et al., 2022). Additionally, differentially expressed genes involved in photosynthesis, glutathione biosynthesis, IAA signal transduction, amino acid biosynthesis, and ABA signal transduction pathways were found in Brassica napus through comparative transcriptome analysis. These genes support drought tolerance mechanisms in rapeseed cultivars (Zhu et al., 2022). Higher seed vigour index and germination energy under drought stress processes have been observed in rapeseed cultivars that use drought-tolerant bacteria for biopriming, such as the strain SH-8, to boost drought tolerance and germination potential in wheat seeds (Shaffique et al., 2023). The desirable genes could be enhanced by molecular markers like RAPD and SSRs; which can be used to investigate the genetic diversity among Indian mustard genotypes. This allows for the identification of diverse parents with high heritability and genetic advance, which can be used to improve yield and quality in early segregating generations (Kumar et al., 2013). Enhancing genes associated with plant growth and seed germination is one of the many advantages of bio priming. Based on research, biopriming entails introducing advantageous microbes, including Bacillus species, into seeds to enhance gene expression and seedling growth (Rakshit, 2018). The processes of seed biopriming include the release of defence-related enzymes, induction of plant growth-promoting activities, and solubilization of soil nutrients, all of which support increased gene expression and plant growth (Deshmukh et al., 2020). Furthermore, bio priming is regarded as a sustainable method of mitigating abiotic stresses- drought in crops while guaranteeing uniform stand development under stress situations (Prasad et al., 2016). It is an environmentally benign substitute for chemical fungicides. Overall, bio priming improves seed germination and seedling establishment in difficult settings by upregulating genes involved in antioxidant defence and DNA damage repair at the molecular level (Forti et al., 2020).

CONCLUSION:

Brassica juncea, also known as Indian mustard, is an important oilseed crop that plays a major role in the agricultural economy and food security of India. The crop's economic and health benefits are further enhanced by its high oil content and significant bioactive components. However, the crop's development and productivity are negatively impacted by the drought, which presents serious issues. Biological priming has become a viable approach to lessen these difficulties, including bio-priming with advantageous bacteria. This method induces particular metabolomic and molecular changes that improve seed germination, seedling vigor, and overall plant growth. It has been demonstrated that bio-priming increases drought tolerance through upregulating genes that respond to stress, improving physiological and biochemical characteristics, and encouraging advantageous interactions between microbes and plants. Research indicates that bio-priming can boost Indian mustard's resistance to drought stress, maintaining productivity and enhancing crop quality. These efforts are further supported by genetic and molecular breakthroughs that increase the crop's resistance to harsh

conditions, such as the discovery of genes resistant to drought and the application of transgenic technology. As a whole, there is a lot of potential to increase Indian mustard's productivity and resistance to drought by combining biological priming techniques with conventional breeding and genetic methods. This integrated approach guarantees this essential crop's sustained economic and nutritional contributions to India's agricultural sector in addition to promoting sustainable agriculture.

REFERENCES:

1. Balmer, A., Pastor, V., Gamir, J., Flors, V., & Mauch-Mani, B. (2015). The "prime-ome": Towards a holistic approach to priming. In *Trends in Plant Science* (Vol. 20, Issue 7, pp. 443–452). Elsevier Ltd. <https://doi.org/10.1016/j.tplants.2015.04.002>
2. Basavanneppa, M. A., & Kumar, B. A. (2020). Production and Economic Efficiencies as Influenced by Mustard Genotypes in Paddy Fallows of Tungabhadra Command Area of Karnataka. *International Journal of Current Microbiology and Applied Sciences*, 9(6), 2376–2380. <https://doi.org/10.20546/ijcmas.2020.906.291>
3. Bhardwaj, S., Solanki, N. S., & Nagar, C. (2020). Study of Phenological Parameters and Agrometeorological Indices of Indian Mustard (*Brassica juncea* L. Czern) Varieties under Different Sowing Dates. *International Journal of Current Microbiology and Applied Sciences*, 9(11), 3431–3436. <https://doi.org/10.20546/ijcmas.2020.911.409>
4. Chandra, K., & Kumar, A. (n.d.). *Impact of Water Deficit Condition on Osmoregulation of the Brassica Species*. www.irj.iars.info
5. Chauhan, A., Ranjan, A., & Jindal, T. (2018). *Biological Control Agents for Sustainable Agriculture, Safe Water and Soil Health* (pp. 71–83). https://doi.org/10.1007/978-3-319-58415-7_6
6. CHAUDHARY, J. N. (2014). *BIOCHEMICAL AND MOLECULAR CHARACTERIZATION OF INDIAN MUSTARD [Brassica juncea (L.) CZERN & COSS] GENOTYPES* (Doctoral dissertation, AAU, Anand).
7. CIG, F., ERMAN, M., INAL, B., BEKTAS, H., SONKURT, M., MIRZAPOUR, M., & CERITOGLU, M. (2022). Mitigation of Drought Stress in Wheat by Bio-priming by PGPB Containing ACC Deaminase Activity. *Ataturk University Journal of Agricultural Faculty*, 53(1), 51–57. <https://doi.org/10.54614/AUAF.2022.972753>
8. Deshmukh, A. J., Jaiman, R. S., Bambharolia, R. P., & Patil, V. A. (2020). Seed biopriming-a review. *International Journal of Economic Plants*, 7(1), 038-043.
9. Divya, K., et al. (2010). Ectopic expression of an ABA- and stress-inducible protein from *Brassica juncea* confers dehydration and salt tolerance to transgenic tobacco. *Plant Cell Reports*, 29(5), 459-469. doi: 10.1007/s00299-010-0835-3
10. Fang, S., Zhao, P., Tan, Z., Peng, Y., Xu, L., Jin, Y., Wei, F., Guo, L., & Yao, X. (2022). Combining Physio-Biochemical Characterization and Transcriptome Analysis Reveal the Responses to Varying Degrees of Drought Stress in *Brassica napus* L. *International Journal of Molecular Sciences*, 23(15). <https://doi.org/10.3390/ijms23158555>
11. Fleming, T. R., Fleming, C. C., Levy, C. C. B., Repiso, C., Hennequart, F., Nolasco, J. B., & Liu, F. (2019). Biostimulants enhance growth and drought tolerance in *Arabidopsis thaliana* and exhibit chemical priming action. *Annals of Applied Biology*, 174(2), 153–165. <https://doi.org/10.1111/aab.12482>
12. Forti, C., Shankar, A., Singh, A., Balestrazzi, A., Prasad, V., & Macovei, A. (2020). Hydropriming and biopriming improve medicago truncatula seed germination and upregulate dna repair and antioxidant genes. *Genes*, 11(3). <https://doi.org/10.3390/genes11030242>
13. Ghonim, M. I. (1999). Induction of systemic resistance against Fusarium wilt in tomato by seed treatment with the biocontrol agent *Bacillus subtilis*.
14. Gupta, S. K., Gupta, P. P., Yadava, T. P., & Kaushik, C. D. (1990). Metabolic changes in mustard due to *Alternaria* leaf blight. *Indian Phytopathology*, 43(1), 64-69.
15. Gupta, J. (2016). Efficacy of Antagonists on mycelium growth and carpogenic germination of sclerotia of *Sclerotinia sclerotiorum* on Indian mustard. *International Journal of Agricultural Invention*, 1(02), 190–193. <https://doi.org/10.46492/ijai/2016.1.2.12>
16. Hilker, M., Schwachtje, J., Baier, M., Balazadeh, S., Bäurle, I., Geiselhardt, S., Hinch, D. K., Kunze, R., Mueller-Roeber, B., Rillig, M. C., Rolff, J., Romeis, T., Schmölling, T., Steppuhn, A., van Dongen, J., Whitcomb, S. J., Wurst, S., Zuther, E., & Kopka, J. (2016). Priming and memory

- of stress responses in organisms lacking a nervous system. *Biological Reviews*, 91(4), 1118–1133. <https://doi.org/10.1111/brv.12215>
17. Holness, S., Bechtold, U., Mullineaux, P., Serino, G., & Vittorioso, P. (2023). Highlight Induced Transcriptional Priming against a Subsequent Drought Stress in *Arabidopsis thaliana*. *International Journal of Molecular Sciences*, 24(7). <https://doi.org/10.3390/ijms24076608>
 18. Irshad, K., Shaheed Siddiqui, Z., Chen, J., Rao, Y., Hamna Ansari, H., Wajid, D., Nida, K., & Wei, X. (2023). Bio-priming with salt tolerant endophytes improved crop tolerance to salt stress via modulating photosystem II and antioxidant activities in a sub-optimal environment. *Frontiers in Plant Science*, 14. <https://doi.org/10.3389/fpls.2023.1082480>
 19. Jha, J. K., Maiti, M. K., Bhattacharjee, A., Basu, A., Sen, P. C., & Sen, S. K. (2006). Cloning and functional expression of an acyl-ACP thioesteraseFatB type from *Diploknema* (Madhuca) butyracea seeds in *Escherichia coli*. *Plant Physiology and Biochemistry*, 44(11–12), 645–655. <https://doi.org/10.1016/j.plaphy.2006.09.017>
 20. Jovičić-Petrović, J., Karličić, V., Petrović, I., Ćirković, S., Ristić-Djurović, J. L., & Raičević, V. (2021). Biomagnetic Priming—Possible Strategy to Revitalize Old Mustard Seeds. *Bioelectromagnetics*, 42(3), 238–249. <https://doi.org/10.1002/bem.22328>
 21. Kępczyński, J., & Kępczyńska, E. (2023). Plant-Derived Smoke and Karrikin 1 in Seed Priming and Seed Biotechnology. In *Plants* (Vol. 12, Issue 12). MDPI. <https://doi.org/10.3390/plants12122378>
 22. Kumar, B., Pandey, A., & Singh, S. K. (2013). GENETIC DIVERSITY FOR AGRO-MORPHOLOGICAL AND OIL QUALITY TRAITS IN INDIAN MUSTARD (*BRASSICA JUNCEA* L. CZERN & COSS). In *N Save Nature to Survive* (Vol. 8, Issue 3). www.thebioscan.in
 23. Kumar, H., Verma, P., John, S. A., & Blaise, D. (n.d.). *Physiological, biochemical and molecular manifestations in response to seed priming with elicitors under drought in cotton*. <http://14.139.232.166/opstat/>
 24. Kumar, S., Yadav, K. G., Patel, S., Singh, P., & Kumar, N. (2020). Nutrient Management Improves Nutrition, Quality and Economic Feasibility of Indian Mustard (*Brassica juncea* L.). *International Journal of Current Microbiology and Applied Sciences*, 9(7), 735–746. <https://doi.org/10.20546/ijcmas.2020.907.085>
 25. Kumari vedna. (2020). *assessment of drought tolerance using drought tolerance indices and their inter relationship in mustard*.
 26. Kumari, A., Avtar, R., Jattan, A. N. M., & Rani, B. (2019). Screening for drought tolerance in Indian mustard (*Brassica juncea* L.) genotypes based on yield contributing characters and physiological parameters. *Journal of Oilseed Brassica*, 10(1), 1-7.
 27. Lal, S., Kumar, V., Gupta, U., Sushma, Shirke, P. A., & Sanyal, I. (2024). Overexpression of the chickpea Metallothionein 1 (MT1) gene enhances drought tolerance in mustard (*Brassica juncea* L.). *Plant Cell, Tissue and Organ Culture (PCTOC)*, 157(1), 6.
 28. Lastochkina, O., Yakupova, A., Avtushenko, I., Lastochkin, A., & Yuldashev, R. (2023). Effect of Seed Priming with Endophytic *Bacillus subtilis* on Some Physio-Biochemical Parameters of Two Wheat Varieties Exposed to Drought after Selective Herbicide Application. *Plants*, 12(8). <https://doi.org/10.3390/plants12081724>
 29. Lang, Minglin., An, Chunju., Wu, Dingding., Chu, Yuan., Song, He., Zeng, Liu. "Constitutive high-expression strong-priming Indian mustard promoter BjCET1 and application thereof." null (2017).
 30. Luo, Y., Dong, D., Su, Y., Wang, X., Peng, Y., Peng, J., & Zhou, C. (2018). Transcriptome analysis of *Brassica juncea* var. *tumida* Tsen responses to *Plasmodiophora brassicae* primed by the biocontrol strain *Zhihengliuella aestuarii*. *Functional and Integrative Genomics*, 18(3), 301–314. <https://doi.org/10.1007/s10142-018-0593-0>
 31. Mallick, S. A., et al. "AUGMENTATION OF BIOCHEMICAL DEFENSE AGAINST ALTERNARIA BLIGHT OF MUSTARD BY INDUCTION OF DROUGHT STRESS." *Journal of Plant Pathology*, vol. 99, no. 1, 2017, pp. 47–60. JSTOR, <http://www.jstor.org/stable/44280572>. Accessed 16 June 2024.
 32. Mitra, D., Pellegrini, M., Olatunbosun, A. N., Mondal, R., Del Gallo, M., Chattaraj, S., ... & Panneerselvam, P. (2023). Seed priming with microbial inoculants for enhanced crop yield. In *Microbial Inoculants* (pp. 99-123). Academic Press.
 33. Mohammad, Anwar, Hossain., Masayuki, Fujita. "Hydrogen Peroxide Priming Stimulates Drought Tolerance in Mustard (*Brassicajuncea* L.) Seedlings." 4 (2013). doi: 10.5376/PGT.2013.04.0020

34. Monika, Sood., Vipul, Kumar., Ruby, Rawal. "Seed biopriming a novel method to control seed borne diseases of crops." null (2021):181-223. doi: 10.1016/B978-0-12-822919-4.00008-9
35. Nayak, U. S., Mishra, I., & Mishra, B. K. (2014). Field evaluation of bio-intensive IPM modules against important insect pests of mustard under North Coastal Plain Zone of Odisha. In *Journal of Biological Control* (Vol. 28, Issue 4).
36. Neeraj, & Singh, K. (2010). *Cyamopsis tetragonoloba* L Taub inoculated with arbuscular mycorrhiza and *Pseudomonas fluorescens* and treated with mustard oil cake overcome *Macrophomina* root-rot losses. *Biology and Fertility of Soils*, 46(3), 237–245. <https://doi.org/10.1007/s00374-009-0422-7>
37. Nivetha, N., Kiruthika, A., Asha, A. D., Lavanya, A. K., Vikram, K. V., Manjunatha, B. S., & Paul, S. (2020). Osmotolerant Rhizobacteria Improve Seedling Vigour and Plant Growth of Mustard under Water Scarcity. *International Journal of Current Microbiology and Applied Sciences*, 9(12), 1928–1937. <https://doi.org/10.20546/ijcmas.2020.912.229>
38. Paul, S., Dey, S. & Kundu, R. Seed priming: an emerging tool towards sustainable agriculture. *Plant Growth Regul* 97, 215–234 (2022). <https://doi.org/10.1007/s10725-021-00761-1>
39. Prasad, S. R., Kamble, U. R., Sripathy, K. V., Bhaskar, K. U., & Singh, D. P. (2016a). Seed biopriming for biotic and abiotic stress management. In *Microbial Inoculants in Sustainable Agricultural Productivity: Vol. 1: Research Perspectives* (pp. 211–228). Springer India. https://doi.org/10.1007/978-81-322-2647-5_12
40. Prasad, S. R., Kamble, U. R., Sripathy, K. V., Bhaskar, K. U., & Singh, D. P. (2016b). Seed biopriming for biotic and abiotic stress management. In *Microbial Inoculants in Sustainable Agricultural Productivity: Vol. 1: Research Perspectives* (pp. 211–228). Springer India. https://doi.org/10.1007/978-81-322-2647-5_12
41. Poyil, Theertha, (2023) "Bioactive Compounds of Mustard, its Role in Consumer Health and in the Development of Potential Functional Foods." *Current Nutrition & Food Science* 19.9: 950-960.
42. Rahaman, S., Ramana, C., Rao, A. S., & Reddy, B. R. (2022). Field Evaluation and Economic Analysis of Manual Drawn Rotor Weeder for Small Farms. *Economic Affairs (New Delhi)*, 67(4), 415–421. <https://doi.org/10.46852/0424-2513.4.2022.7>
43. Rajput, R. K., Singh, S., Varma, J., Rajput, P., Singh, M., Nath, S., Ravindra, C., & Rajput, K. (2018). Effect of different levels of nitrogen and sulphur on growth and yield of Indian mustard (*Brassica juncea* (L.) Czern and Coss.) in salt affected soil. ~ 1053 ~ *Journal of Pharmacognosy and Phytochemistry*, 7(1).
44. Rakshit, A. (2018). Impact assessment of bio priming mediated nutrient use efficiency for climate resilient agriculture. In *Climate Change and Agriculture in India: Impact and Adaptation* (pp. 56–68). Springer International Publishing. https://doi.org/10.1007/978-3-319-90086-5_6
45. Rasul, F., Gupta, S., Olas, J. J., Gechev, T., Sujeeth, N., & Mueller-Roeber, B. (2021). Priming with a seaweed extract strongly improves drought tolerance in arabis. *International Journal of Molecular Sciences*, 22(3), 1–28. <https://doi.org/10.3390/ijms22031469>
46. R.A., Gami., R.M., Chauhan., A., Parihar., S.D., Solanki., V., H., Kanbi. "Molecular characterization in mustard [*Brassica juncea* L. Czern&Coss.]." *International Journal of Agriculture, Environment and Biotechnology*, 6 (2013):61-67.
47. Rawat, monika. , Rawat, L. , B. T. , C. S. (2022). Comparative studies of the effect of microbial inoculants and inorganic chemicals on growth, yield, 247 yield contributing traits and disease suppression in two varieties of mustard green (*Brassica juncea*L.) under open field conditions in mid hills of U., & *PANTNAGAR JOURNAL OF RESEARCH* (Vol. 20, Issue 2).
48. Reddy, P. P. (2012). Bio-priming of Seeds. In *Recent advances in crop protection* (pp. 83–90). Springer India. https://doi.org/10.1007/978-81-322-0723-8_6
49. Rhaman, M. S., Imran, S., Rauf, F., Khatun, M., Baskin, C. C., Murata, Y., & Hasanuzzaman, M. (2020). *plants Seed Priming with Phytohormones: An Effective Approach for the Mitigation of Abiotic Stress*. <https://doi.org/10.3390/plants10010>
50. Sahoo, C. R., Dash, M., & Acharya, N. (2019). Biochemical Responses of Indian Mustard to Water Stress. *International Journal of Current Microbiology and Applied Sciences*, 8(02), 1711–1718. <https://doi.org/10.20546/ijcmas.2019.802.201>
51. Sahu, A., Salam, J. L., Verma, S., & Samuel, S. (2020). Combining Ability and Heterosis for Seed Yield and its Attributing Traits in Indian Mustard (*Brassica juncea* L. Czern& Coss).

- International Journal of Current Microbiology and Applied Sciences*, 9(7), 720–727. <https://doi.org/10.20546/ijcmas.2020.907.083>
52. Saleem, M., Elahi, E., Gandahi, A. W., Bhatti, S. M., Ibrahim, H., & Ali, M. (2019). Effect of sulphur application on growth, oil content and yield of sunflower. *Sarhad Journal of Agriculture*, 35(4). <https://doi.org/10.17582/JOURNAL.SJA/2019/35.4.1198.1203>
 53. SAPNA1, S. J. , V. J. and A. S. R. (2011). *Biochemical changes in antioxidant pathway metabolites in Indian mustard leaves due to white rust infection.*
 54. Singh, V. V., Garg, P., Meena, H. S., & Meena, M. L. (2018). Drought stress response of Indian mustard (*Brassica juncea* L) genotypes.
 55. Shaffique, S., Imran, M., Kang, S. M., Khan, M. A., Asaf, S., Kim, W. C., & Lee, I. J. (2023). Seed Bio-priming of wheat with a novel bacterial strain to modulate drought stress in Daegu, South Korea. *Frontiers in Plant Science*, 14. <https://doi.org/10.3389/fpls.2023.1118941>
 56. Shakeel, A., Khan, A. A., & Ahmad, G. (2019). The potential of thermal power plant fly ash to promote the growth of Indian mustard (*Brassica juncea*) in agricultural soils. *SN Applied Sciences*, 1(4). <https://doi.org/10.1007/s42452-019-0404-9>
 57. Sharma, K. K., & Singh, U. S. (2014). Induction of water stress tolerance of mustard plants using *Trichoderma* as biological seed treatment. In *Journal of Applied and Natural Science* (Vol. 6, Issue 2). www.ansfoundation.org
 58. Sharma, S., Singh, J., Munshi, G. D., & Munshi, S. K. (2010). Biochemical changes associated with application of biocontrol agents on Indian mustard leaves from plants infected with *Alternaria* blight. *Archives of Phytopathology and Plant Protection*, 43(4), 315–323. <https://doi.org/10.1080/03235400701804109>
 59. Sharma, A., Kumari, V., & Rana, A. (2022). Genetic variability studies on drought tolerance using agro-morphological and yield contributing traits in rapeseed-mustard. *International Journal of Bio-resource and Stress Management*, 13(7), 771-779.
 60. Shi, L., Tang, X., & Tang, G. (2016). GUIDE-Seq to Detect Genome-wide Double-Stranded Breaks in Plants. In *Trends in Plant Science* (Vol. 21, Issue 10, pp. 815–818). Elsevier Ltd. <https://doi.org/10.1016/j.tplants.2016.08.005>
 61. Shrivastav, A., Tripathi, M. K., Tiwari, S., Tripathi, N., Tiwari, P. N., Bimal, S. S., Rajpoot, P., & Chauhan, S. (2023). Evaluation of Genetic Diversity in Indian Mustard (*Brassica juncea* var. *rugosa*) Employing SSR Molecular Markers. *PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY*, 10–21. <https://doi.org/10.56557/pcbmb/2023/v24i3-48245>
 62. Shukla, P., & Gupta, J. K. (2020). Study of Cost and Profit Strategies for the Mustard Crop in Madhya Pradesh. *Current Journal of Applied Science and Technology*, 79–87. <https://doi.org/10.9734/cjast/2020/v39i2730922>
 63. Singh, A., & Meena, R. S. (n.d.). *EFFECT OF FOLIAR SPRAY OF BIOREGULATORS AND IRRIGATION ON DRY MATTER ACCUMULATION OF MUSTARD (Brassica juncea L.)*.
 64. Singh, S., Sarkar, D., Rakesh, S., Singh, R. K., & Rakshit, A. (2023). Bioprimering with *Bacillus subtilis* Enhanced the Sulphur Use Efficiency of Indian Mustard under Graded Levels of Sulphur Fertilization. *Agronomy*, 13(4). <https://doi.org/10.3390/agronomy13040974>
 65. Singh, V. V., beer, B., Meena, H. S., Kulshrestha, S., Dubey, M., Gurjar, N., Garg, P., Meena, M. L., & Rai, P. K. (2020). Heterosis and Combining Ability Analysis for Yield and Yield Attributes in Indian Mustard (*Brassica juncea* L.). *International Journal of Current Microbiology and Applied Sciences*, 9(3), 1622–1632. <https://doi.org/10.20546/ijcmas.2020.903.190>
 66. Sinha, S., Jha, J. K., Maiti, M. K., Basu, A., Mukhopadhyay, U. K., & Sen, S. K. (2007). Metabolic engineering of fatty acid biosynthesis in Indian mustard (*Brassica juncea*) improves nutritional quality of seed oil. *Plant Biotechnology Reports*, 1(4), 185–197. <https://doi.org/10.1007/s11816-007-0032-5>
 67. Sree, B. K., Bara, B. M., & Pal, A. K. (2022). Effect of Various Presowing Seed Treatments on Growth and Yield Parameters of Yellow Mustard (*Sinapis alba*.L) Variety (ISP -186). *International Journal of Environment and Climate Change*, 837–843. <https://doi.org/10.9734/ijecc/2022/v12i1030871>
 68. Tenenboim, H., & Brotman, Y. (2016). Omic Relief for the Biotically Stressed: Metabolomics of Plant Biotic Interactions. In *Trends in Plant Science* (Vol. 21, Issue 9, pp. 781–791). Elsevier Ltd. <https://doi.org/10.1016/j.tplants.2016.04.009>

69. Thakro, V., Malik, N., Basu, U., Srivastava, R., Narnoliya, L., Daware, A., Varshney, N., Mohanty, J. K., Bajaj, D., Dwivedi, V., Tripathi, S., Jha, U. C., Dixit, G. P., Singh, A. K., Tyagi, A. K., Upadhyaya, H. D., & Parida, S. K. (2023). A superior gene allele involved in abscisic acid signaling enhances drought tolerance and yield in chickpea. *Plant Physiology*, 191(3), 1884–1912. <https://doi.org/10.1093/plphys/kiac550>
70. Thapa, S., Baral, B., Shrestha, M., & Chandra Dahal, Dr. K. (2022). EFFECT OF DIFFERENT PRIMING METHODS ON GERMINATION BEHAVIOUR OF BROADLEAF MUSTARD CV. MARPHA CHAUDA PAATE. *Tropical Agrobiodiversity*, 3(2), 52–59. <https://doi.org/10.26480/trab.02.2022.52.59>
71. Tripathi, P., et al. (2017). Functional characterization of Brassica juncea BjSOS3 and BjSOS2 homologs in plant salt tolerance. *Frontiers in Plant Science*, 8, 650. doi: 10.3389/fpls.2017.00650
72. Tugizimana, F., Mhlongo, M. I., Piater, L. A., & Dubery, I. A. (2018). Metabolomics in plant priming research: The way forward? In *International Journal of Molecular Sciences* (Vol. 19, Issue 6). MDPI AG. <https://doi.org/10.3390/ijms19061759>
73. Tulsı, Ram, Yadav., Pradip, Kumar, Saini., Ravi, Yadav., Raj, Bhahadur. "Effect of seed priming with plant growth regulators on physiological changes of Indian mustard (Brassica juncea L. Czern&Coss)." *Journal of Pharmacognosy and Phytochemistry*, 7 (2018):355-358.
74. Wang, X., Chen, J., Ge, J., Huang, M., Cai, J., Zhou, Q., Dai, T., Mur, L. A. J., & Jiang, D. (2021). The different root apex zones contribute to drought priming induced tolerance to a reoccurring drought stress in wheat. *Crop Journal*, 9(5), 1088–1097. <https://doi.org/10.1016/j.cj.2020.11.008>
75. Wang, X., Li, Q., Xie, J., Huang, M., Cai, J., Zhou, Q., Dai, T., & Jiang, D. (2021). Abscisic acid and jasmonic acid are involved in drought priming-induced tolerance to drought in wheat. *Crop Journal*, 9(1), 120–132. <https://doi.org/10.1016/j.cj.2020.06.002>
76. Waraich, E. A., Hussain, A., Ahmad, Z., Ahmad, M., & Barutçular, C. (2021). Foliar application of sulfur improved growth, yield and physiological attributes of canola (*brassica napus* L.) under heat stress conditions. *Journal of Plant Nutrition*, 45(3), 369–379. <https://doi.org/10.1080/01904167.2021.1985138>
77. Yadav, S. K., et al. (2019). Heat shock proteins in Brassica juncea: Genome-wide identification, phylogeny, structure, and expression profiling under heat stress. *Journal of Experimental Botany*, 70(18), 4771-4786. doi: 10.1093/jxb/erz216
78. Zhang, J., Huang, D., Zhao, X., Zhang, M., Wang, Q., Hou, X., Di, D., Su, B., Wang, S., & Sun, P. (2022). Drought-responsive WRKY transcription factor genes IgWRKY50 and IgWRKY32 from *Iris germanica* enhance drought resistance in transgenic Arabidopsis. *Frontiers in Plant Science*, 13. <https://doi.org/10.3389/fpls.2022.983600>
79. Zhu, B., Xu, H., Guo, X., Lu, J., Liu, X., & Zhang, T. (2022a). *Comparative analysis of drought responsive transcriptome in Brassica napus genotypes with contrasting drought tolerance under different potassium levels*. <https://doi.org/10.21203/rs.3.rs-2077417/v1>
80. Zhu, B., Xu, H., Guo, X., Lu, J., Liu, X., & Zhang, T. (2022b). *Comparative analysis of drought responsive transcriptome in Brassica napus genotypes with contrasting drought tolerance under different potassium levels*. <https://doi.org/10.21203/rs.3.rs-2077417/v1>
81. Zhu, J., Cai, D., Wang, J., Cao, J., Wen, Y., He, J., Zhao, L., Wang, D., & Zhang, S. (2021). Physiological and anatomical changes in two rapeseed (*Brassica napus* L.) genotypes under drought stress conditions. *Oil Crop Science*, 6(2), 97–104. <https://doi.org/10.1016/j.ocsci.2021.04.003>