

Original Research Article

Pre-harvest application of NAA, Ca, and B on fruit drop, and yield of Thailand ber (*Ziziphus mauritiana*)

ABSTRACT

The present study was carried out to investigate the beneficial effects of the pre-harvest spray of NAA, calcium, and boron on fruit drop and yield of Thailand ber (*Ziziphus mauritiana*) at the Experimental Farm, Department of Horticulture, AAU, Jorhat-13 during 2022-2023. The treatment T₄ i.e. spraying with NAA @ 20 ppm + B @ 0.5% resulted in maximum fruit set (4.68%), lesser fruit drop (53.28%), the highest yield per tree (18.12 kg/tree), yield per hectare (24.16 t), the maximum thickness of pulp (1.94 mm) and pulp weight (62.25 g). Spraying of NAA 20 ppm in combination with Boron 5% is the most profitable approach for minimizing fruit drop and maximizing production.

Keywords: Calcium, Boron, Fruit drop, Fruit retention, NAA.

INTRODUCTION

One of the oldest and most popular fruits native to China and India is the ber (*Zizyphus mauritiana* Lamk.), which is in the family Rhamnaceae and genus *Zizyphus*. Genus *Zizyphus* consists of 50 genera and more than 600 species (Pareek, 1983). Ber fruits are packed with vitamins, carotenoids, pectin, and minerals. They have higher ascorbic acid content than citrus and are second only to guava among tropical fruits (Meena et al., 2022). In recent years, Thailand developed a crossbreed (green apple x jujube) variety of ber called “Thailand ber” or “Thai Apple ber” that is rapidly gaining popularity worldwide. The new cultivar has an apple-like appearance (Mathangi & Maran, 2020). Thailand ber plants bear large round to ovate fruits. It outperforms in terms of both quality and economic benefits. As a result, Thailand ber cultivation is growing in popularity across India, especially in Assam. Round green and roundish ovate reddish varieties are commonly grown in Assam (Sarma, 2022). It also bears fruit within 6 to 8 months of planting and is very nutritious. Delicious fruits have a crunchy texture and are very juicy and sweet.

Fruit drop is the separation of premature fruits from the attached plant. When pollination occurs in ber, there is a significant loss of blooms and ovaries due to a number of circumstances. The majority of fruit drop occurs at an early stage of fruit development (pea stage). Numerous factors, including hormonal instability, embryonic abortion, and unfavourable weather are responsible for immature fruit drop in ber(Singh et al., 1991) .

The growth regulators and micronutrients play a significant role in vegetative propagation, inducing seedlessness, increasing fruit set, preventing pre-harvest fruit drop, regulating flowering, controlling fruit size, thinning flowers and fruits, and improving fruit quality and yield in many fruit crops. Mineral nutrients also play a role in the flowering and fruiting processes, pollen germination, cell division, and glucose metabolism. Exogenous application of plant growth regulators can effectively prevent fruit drop in ber.

Keeping all the above-mentioned facts and issues into consideration, the present investigation has been undertaken with the objective to study the influence of NAA, Ca, and B on premature fruit drop and yield in Thailand Ber

MATERIALS AND METHOD

The present investigation was carried out at the Experimental Farm, Department of Horticulture, Assam Agricultural University, Jorhat 13 during 2022- 2023. The standing crop of Thailand Ber (*Ziziphus mauritiana*) was selected for this experiment. The crop was sprayed with seven treatments viz., T₀- Control, T₁-NAA @ 20ppm, T₂-Borax @ 0.5%, T₃-CaCl₂ @ 1%, T₄-NAA @ 20ppm + Borax @ 0.5%, T₅-NAA @ 20ppm + CaCl₂ @ 1%, T₆-NAA @ 20ppm + Borax @ 0.5% + CaCl₂ @ 1%. The foliar application of the treatments was done thrice at 45-day intervals after flower initiation.

Using a pneumatic foot sprayer equipped with a nozzle, the spraying was carried out in the afternoon between 3 and 5 PM. Each tree's five uniform branches were tagged. Before the first spray, the flower buds in the chosen cymes were counted. At seven days following the second spray, the number of little fruits that had been set on each marked branch was counted.

The per cent fruit set was calculated by dividing the number of fruits by the number of flowers. The number of buds/fruitlets retained on each branch was counted at 15-day intervals after foliar application. The average number of fruit dropped from each branch was calculated in percent drop by subtracting the number of fruitlets at the initial from the number of fruit retention at harvest and dividing it by the number of fruitlets at the initial.

The fruits' length and girth was recorded using a vernier caliper and expressed in cm. The fruit volume was calculated using the water displacement method using a measuring cylinder and expressed in cc. Fruit weights were measured in grams using an electronic balance. To calculate the yield per plant in kg, the total number of fruits in each individual plant was recorded and the average fruit weight of the plant was multiplied by that number. The yield per plant was multiplied by the total number of plants for each treatment and expressed in kg/ha.

The experiment was laid out in a Randomized Block Design with seven treatments and four replications. Observations obtained during field trials along with data acquired from laboratory investigations were subject to analysis of variance (ANOVA). 'F' values were used to calculate the significance and non-significance of the variance (Panse & Sukhatme, 1985). Critical differences at a 0.05% probability level were calculated only when the 'F' value was significant.

RESULTS AND DISCUSSION

Flowering and Fruiting Parameter

Foliar applications of CaCl₂ and NAA in combination required lesser time to reach maturity than in other treatments. Calcium binds with pectin, producing calcium pectate which increases middle lamella rigidity and resistance to enzymes that break down pectin, including polygalacturonase (Grant et al., 1973). These results are in agreement with prior reports of Schlegel and Schonherr (2002) on apples and Abbasi et al. (2010) on tomatoes.

NAA in association with boron significantly influenced the flowering and fruit set in ber as boron levels affected the pollen tube growth and pollen germination (Ganie et al., 2013). The effects of NAA on plant growth are strongly influenced by the time of application and concentration levels. NAA enhances fruit set by preventing pre-harvest fruit drop by reducing the impact of ethylene (Yuan & Carbaugh., 2007). Boron regulates metabolic processes involved in the translocation of carbohydrates, the formation of cell walls, and the synthesis of RNA (Ram & Bose, 2000). These factors led to an improvement in flowering and fruit set in ber during the experimentation. These results are in accordance with the reports of Shukla (2011), Helal et al. (2019), and Badal and Tripathi (2021)

Table 1. Days from flowering to harvesting, flower/plant, fruit set, fruits/plant

| Treatments | Days from flowering to harvesting | Flower per plant | Fruit set % | Fruits per plant |
|--|--|-------------------------|--------------------|-------------------------|
| T ₀ : Control | 125.50 | 14090.00 | 3.99 | 241.25 |
| T ₁ : NAA @ 20ppm | 103.75 | 15337.50 | 4.23 | 304.75 |
| T ₂ : Borax @ 0.5% | 105.00 | 16342.50 | 4.48 | 308.00 |
| T ₃ : CaCl ₂ @ 1% | 102.50 | 16340.00 | 4.26 | 300.50 |
| T ₄ : NAA @ 20ppm + Borax @ 0.5% | 103.25 | 17197.50 | 4.68 | 346.25 |
| T ₅ : NAA @ 20ppm + CaCl ₂ @ 1% | 99.50 | 16160.00 | 4.48 | 325.25 |
| T ₆ : NAA @ 20ppm + Borax @ 0.5% + CaCl ₂ @ 1% | 102.25 | 16975.00 | 4.57 | 328.00 |
| Mean | 105.96 | 16063.21 | 4.43 | 307.75 |
| S.Ed (±) | 1.414 | 4.062 | 0.019 | 1.621 |

| | | | | |
|-----------|-------|-------|-------|-------|
| CD (0.05) | 2.971 | 8.535 | 0.041 | 3.382 |
|-----------|-------|-------|-------|-------|

Fruit Drop and Fruit Retention

Fruit growth and development after fruit set is largely dependent on photosynthates supplied by leaves, and an increase in sink ability in fruit via increased levels of endogenous growth promoters, which regulate the mobility of nutrients and photosynthates from the source (leaves) to the sink (fruit), resulting in increased fruit growth and development and decreased fruit drop (Emongor & Murr, 2001).

NAA and B when applied alone or in combination enhanced the fruit retention thereby preventing fruit drop. NAA's beneficial role in fruit retention could be attributed to its involvement in cell division, cell elongation, and increased volume of intercellular spaces in mesocarpic cells, which could have improved plant health and resulted in a healthy fruit, which was ultimately beneficial in fruit retention (Kaur & Kaur, 2017). The exogenous application of NAA might delay the abscission of the fruit's outer layer, thereby increasing fruit retention. It might also have enhanced auxin levels in plants which prevented fruit drop.

Boron reduced fruit drop and increased fruit retention as it played an important role in increasing pollen viability and fertilization, carbohydrate translocation, and synthesis of auxin that delayed the formation of the abscission layer in the earlier stages of fruit development (Wojcik & Wojcik, 2006). The results conform with the reports of Gill and Bal (2009), Sharif et al. (2016), Tirkey et al. (2018), Singh and Singh (2019) and Badal and Tripathi (2021)

Table 2. Fruit drop, fruit retention, yield/tree, and yield/ha

| Treatments | Fruit drop % | Fruit retention % | Yield/tree e (kg) | Yield/h a (t) |
|---|-------------------------|------------------------------|------------------------------|--------------------------|
| T ₀ : Control | 62.80 | 37.20 | 12.98 | 17.30 |
| T ₁ : NAA @ 20ppm | 54.46 | 45.54 | 16.25 | 21.66 |
| T ₂ : Borax @ 0.5% | 54.17 | 45.83 | 16.50 | 21.99 |
| T ₃ : CaCl ₂ @ 1% | 56.84 | 43.16 | 15.81 | 21.08 |
| T ₄ : NAA @ 20ppm + Borax @ 0.5% | 53.28 | 46.72 | 18.12 | 24.16 |
| T ₅ : NAA @ 20ppm + CaCl ₂ @ 1% | 57.74 | 42.26 | 17.50 | 23.33 |
| T ₆ : NAA @ 20ppm + Borax @ 0.5% + CaCl ₂ @ | 57.71 | 42.29 | 17.60 | 23.46 |

| | | | | |
|----------------|-------|-------|-------|-------|
| 1% | | | | |
| Mean | 56.71 | 43.29 | 16.39 | 21.85 |
| S.Ed (\pm) | 0.537 | 0.536 | 0.232 | 0.308 |
| CD (0.05) | 1.130 | 1.127 | 0.487 | 0.647 |

Yield and Yield Attributing Parameters

The increase in yield owing to NAA and Boron applied in combination is associated with a high rate of enzymatic activity as well as biosynthesis of auxin, as well as an increase in the number and size of fruit, which ultimately increased the yield per plant (Singh & Singh, 2019). As a direct correlation between the auxin content and fruit growth has been reported by Krishnamoorthy (1981), the application of NAA at the preharvest stage may have increased the auxin level in fruits, which ultimately may have helped in the improvement of cell size and subsequently fruit size.

Calcium chloride has the greatest impact on fruit length and volume because it increases cell size and the strength of the carbohydrate sink (Qeyami et al., 2020). The major roles that boron played in increasing fruit size and weight in the current experiment can be attributed to its ideal dose, which may have generated the desired improvement due to its involvement in hormone metabolism, accelerated cell division, elongation, and expansion of cells and it may have aided in the mobilization of food materials to the fruits, thereby boosting their size and weight (Singh & Singh, 2019). The increased fruit weight might be related to the strengthening of the middle lamella and, as a result, the cell wall, which might have increased the free passage of solutes to the fruits (Kumar et al., 2013). The results are in accordance with the reports of Phookan et al. (1991), Londe et al. (2020), Dixit et al. (2013), and Arora and Singh (2014)

Table 3. Fruit length, fruit girth, fruit volume, and fruit weight

| Treatments | Fruit length (cm) | Fruit girth (cm) | Fruit volume (cc) | Fruit weight (g) |
|---|-------------------|------------------|-------------------|------------------|
| T ₀ : Control | 4.92 | 4.50 | 54.75 | 52.25 |
| T ₁ : NAA @ 20ppm | 5.00 | 4.83 | 66.00 | 65.25 |
| T ₂ : Borax @ 0.5% | 4.92 | 5.60 | 64.75 | 63.75 |
| T ₃ : CaCl ₂ @ 1% | 5.65 | 4.80 | 68.00 | 64.25 |
| T ₄ : NAA @ 20ppm + Borax @ 0.5% | 5.38 | 4.70 | 62.25 | 65.25 |
| T ₅ : NAA @ 20ppm + CaCl ₂ @ 1% | 5.25 | 4.88 | 60.75 | 61.25 |

| | | | | |
|--|-------|-------|-------|-------|
| T ₆ : NAA @ 20ppm + Borax @ 0.5% + CaCl ₂ @ 1% | 5.20 | 4.75 | 61.25 | 57.25 |
| Mean | 5.19 | 4.86 | 62.54 | 61.32 |
| S.Ed (±) | 0.241 | 0.229 | 2.523 | 1.856 |
| CD (0.05) | 0.495 | 0.486 | 5.070 | 3.899 |

Physical Characteristics of Fruit

Combined application of NAA and boron had a stimulatory effect on the plant metabolism which enhanced the thickness and weight of pulp. Boron, either alone or in combination, also contributed to the maximum increase in pulp by accelerating the transport of photosynthates from the leaf to the fruits (Dugger, 1983). Similar results were reported by Tirkey et al. (2018), Gaur et al. (2014).

The decrease in stone weight in NAA-treated fruits could be attributed to auxins inducing a parthenocarpic effect, resulting in decreased stone weight (Singh et al., 2001). NAA possibly enhanced solid deposition, which increased cell size by increasing water storage in intracellular space, which might have reduced stone weight (Kumar et al., 2013). The application of boron increased the pulp weight while decreasing the stone weight, resulting in a high pulp/stone ratio. Similar results were reported by Bankar and Prasad (1990), Bal et al. (1986), Singh et al. (2016), and Brahmachari and Kumar (1997).

Table 4. Thickness of pulp, pulp weight, stone weight, and pulp stone ratio

| Treatments | Thickness of pulp (cm) | Pulp weight (g) | Stone weight (g) | Pulp stone ratio |
|--|------------------------|-----------------|------------------|------------------|
| T ₀ : Control | 1.70 | 47.50 | 4.08 | 11.66 |
| T ₁ : NAA @ 20ppm | 1.90 | 61.75 | 3.25 | 19.23 |
| T ₂ : Borax @ 0.5% | 1.85 | 61.00 | 3.00 | 20.33 |
| T ₃ : CaCl ₂ @ 1% | 1.85 | 60.75 | 3.00 | 20.25 |
| T ₄ : NAA @ 20ppm + Borax @ 0.5% | 1.94 | 62.25 | 3.25 | 19.44 |
| T ₅ : NAA @ 20ppm + CaCl ₂ @ 1% | 1.85 | 58.50 | 3.50 | 17.27 |
| T ₆ : NAA @ 20ppm + Borax @ 0.5% + CaCl ₂ @ 1% | 1.75 | 54.25 | 3.00 | 18.08 |
| Mean | 1.83 | 58.00 | 3.30 | 18.04 |
| S.Ed (±) | 0.051 | 2.021 | 0.251 | 1.541 |
| CD (0.05) | 0.105 | 4.245 | 0.527 | 3.238 |

Fruit Surface Colour

All fruits reached a point of maximum ripeness, followed quickly by decomposition. Colour can help determine the ripeness of fruits. It is also necessary to evaluate how pre- and post-harvest treatments affect fruit colour. Colour features include hue, which is a visual phenomenon in which a region appears to have characteristics of one or more proportions of the four recognized colors - red, yellow, green, and blue and so hue angle value is the actual observed colour (Schanda, 2007). The angle between 0° and 360° of the colour wheel is represented by hue. The saturation level of colour is measured by chroma.

The L* (lightness) axis is 0 for black and 100 for white. L* value ranged from 61.12 (darkest) in T₄ to 65.21 (lightest) in T₀. For a* axis, positive values are shown as red, negative values as green, and 0 is the neutral value. The a* value ranged from -9.08 (less green) in T₃ to -12.08 (more green) in T₄. The negative value of 'a*' indicates the greenness of ber fruits. For the b* axis, positive values are shown as yellow, negative values as blue, and 0 is the neutral value. The b* value ranged from 43.54 (less yellow) in T₂ to 45.42 (slightly more yellow) in T₁. The a* and b* values of all the fruits lied in the second quadrant with H* value ranging from -1.30 in T₄ to -1.37 in T₃.

The hue angle ranged from 105° in T₄ and 102° in T₃ which indicated that the green fruits had a tinge of yellow colour in them. C* value was in the range of 44.64 (less intense) in T₂ to 46.48 (more intense) in T₁. A study of fruit skin colour in the apple variety 'Granny Smith, revealed that colour changes were primarily caused by variations in green pigment concentrations (chlorophylls) and were almost independent of yellow pigment concentrations (carotenoids and flavonoids). Since the human eye is particularly sensitive to the presence of green colour, the consequent yellower fruit hue might have been partially caused by decreased chlorophyll levels unmasking yellow pigments (Mussini et al., 1985).

Table 5. CIE L*, CIE a*, CIE b*, CIE H* and CIE C* value

| Treatments | CIE L* | CIE a* | CIE b* | CIE H* | CIE C* |
|---|-----------|-----------|-----------|-----------|-----------|
| T ₀ : Control | 65.21 | -10.00 | 43.30 | -1.35 | 46.44 |
| T ₁ : NAA @ 20ppm | 62.78 | -9.82 | 45.42 | -1.36 | 46.48 |
| T ₂ : Borax @ 0.5% | 64.09 | -9.67 | 43.54 | -1.35 | 44.64 |
| T ₃ : CaCl ₂ @ 1% | 63.62 | -9.08 | 44.12 | -1.37 | 45.09 |
| T ₄ : NAA @ 20ppm + Borax @ 0.5% | 61.12 | -12.08 | 44.29 | -1.30 | 45.91 |

| | | | | | |
|--|-------|--------|-------|-------|-------|
| T ₅ : NAA @ 20ppm + CaCl ₂ @ 1% | 63.62 | -11.13 | 44.11 | -1.32 | 45.51 |
| T ₆ : NAA @ 20ppm + Borax @ 0.5% + CaCl ₂ @ 1% | 62.55 | -9.92 | 44.50 | -1.35 | 45.63 |
| Mean | 63.28 | -10.24 | 44.47 | -1.34 | 45.67 |
| S.Ed (±) | 1.890 | 1.368 | 1.217 | 0.021 | 1.200 |
| CD (0.05) | NS | NS | NS | 0.045 | NS |

CONCLUSION

The treatment T₄ (NAA @ 20 ppm + B @ 0.5 %) was best in terms of fruit set, fruit drop, and various yield parameters. It is evident from the data that using exogenous plant growth regulators and nutrients can yield positive outcomes if the proper spraying interval and dose of growth regulator and nutrients are employed. Growth regulators, micro and macronutrients show economic viability and ease of administration for farmers with low concentration and one-time application. Therefore it can be recommended for practice to increase production. In Assam, areas under Thailand ber cultivation is increasing due to its high demand. Fruit drop is a limiting factor as it causes yield losses. Foliar application of PGR's and nutrients can be utilized for improved production.

REFERENCES

- Abbasi, N., Zafar, L., Khan, H., & Qureshi, A. (2010). Effects of naphthalene acetic acid and calcium chloride application on nutrient uptake, growth, yield and post harvest performance of tomato fruit. *Pakistan Journal of Botany*, 45, 1581-1587.
- Arora, R., & Singh, S. (2014). Effect of growth regulators on quality of ber (*Zizyphus mauritiana* Lamk.) cv. Umran. *Agricultural Science Digest*, 34(2), 102-106.
- Badal, D. S., & Tripathi, V. K. (2021). Influence of Foliar Feeding of NAA and Boron on Growth, Flowering, Fruiting, and Yield of Winter Season Guava (*Psidium guajava* L.) cv. L-49. *Biological Forum – An International Journal*, 13(3), 387-391.
- Bal, J. S., Singh, S. N., & Randhawa, J. S. (1986). Response of naphthalene acetic acid spray at fruit set and slow growth phase in ber fruits (*Zizyphus mauritiana* Lamk.). *Journal of Research, Punjab Agricultural University*(23), 569-572.
- Bankar, G. S., & Prasad, R. N. (1990). Effect of gibberellic acid and NAA on fruit set and quality of fruit in ber cv. gola. *Progressive Horticulture*, 22(1-4), 60-62.
- Brahmachari, V. S., & Kumar, R. (1997). Effect of foliar spray of mineral nutrients on fruit set retention and cracking of litchi fruit. *Haryana Journal of Horticultural Sciences*, 26(3), 177-180.
- Dixit, A., Shaw, S. S., & Pal, V. (2013). Effect of micronutrients and plant growth regulators on fruiting of litchi. *Hortflora Research Spectrum*, 2(1), 77-80.
- Dugger, W. M. (1983). Boron in plant metabolism. In *Encyclopedia of Plant Physiology*. (Vol. 15B). Berlin, New York.: Springer-verlag.

- Emongor, V. E., & Murr, D. P. (2001). Effect of Benzyladenine on Fruit Set, Quality, and Vegetative Growth of 'Empire' Apples. *East African agricultural and forestry journal*, 67(1-2), 83-91. <https://doi.org/10.1080/00128325.2001.11663340>
- Ganie, M. A., Akhter, F., Bhat, M. A., Malik, A. R., Junaid, J. M., Shah, M. A., Bhat, A. H., & Bhat, T. A. (2013). Boron — a critical nutrient element for plant growth and productivity with reference to temperate fruits. *Current Science*, 104(1), 76-85. <http://www.jstor.org/stable/24110665>
- Gaur, B., Beer, K., Hada, T. S., Kanth, N., & Syamal, M. M. (2014). Studies on the effect of foliar application of nutrients and GA3 on fruit yield and quality of winter Season Guava. . *International Journal of Environmental Sciences*, 6, 479-483.
- Gill, P. P. S., & Bal, J. S. (2009). Effect of growth regulator and nutrients spray on control of fruit drop, fruit size and quality of ber under sub-montane zone of Punjab. *Journal of Horticultural Sciences*, 4(2), 161-163.
- Grant, G. T., Morris, E. R., Rees, D. A., Smith, P. J. C., & Thom, D. (1973). Biological interactions between polysaccharides and divalent cations: The egg-box model. *FEBS Letters*, 32(1), 195-198. [https://doi.org/https://doi.org/10.1016/0014-5793\(73\)80770-7](https://doi.org/https://doi.org/10.1016/0014-5793(73)80770-7)
- Helal, E. M., Ashour, N. E., Merwad, M. M., & Mansour, A. E. M. (2019). Effect of Some Growth Regulators and Boron on Fruiting and Quality of Orange. *Middle East Journal of Agriculture Research*, 8(2), 594-599.
- Kaur, S., & Kaur, A. (2017). Effect of growth regulators on yield and quality of winter Guava cv. Allahabad Safeda. . *International Journal of Advanced Research*, 5(9), 98-103.
- Krishnamoorthy, H. N. (1981). *Plant growth substances including applications in agriculture*. Tata McGraw Hill Publishing Co. Ltd.
- Kumar, R., Tiwari, R., & Kumawat, B. R. (2013). Quantitative and qualitative enhancement in guava (*Psidium guajava* L.) cv. Chittidar through foliar feeding. *International Journal of Agricultural Sciences*, 9(1), 177-181.
- Londe, M., Kshirsagar, D. B., & Shinde, S. R. (2020). Effect of plant growth regulators and micronutrients on growth and fruit setting, fruit retention, and yield of tomato (*Solanum lycopersicum* L.) during summer season. *Multilogic in science*, 10.
- Mathangi, S., & Maran, J. P. (2020). A study on Apple ber to identify the suitability of new product development. *Plant Science Today*, 7(1), 61-69.
- Meena, V. S., Gora, J. S., Singh, A., Ram, C., Meena, N. K., Pratibha, R., Y., Basile, B., & Kumar, P. (2022). Underutilized Fruit Crops of Indian Arid and Semi-Arid Regions: Importance, Conservation and Utilization Strategies. *Horticulturae*, 8(2), 171.
- Mussini, E., Correa, N., & Crespo, G. (1985). Evolution of pigments in Granny Smith apple fruits. *Phyton - Annales Rei Botanicae*, 45, 79-84.
- Panse, V. G., & Sukhatme, P. V. (1985). *Statistical methods for Agricultural workers* (4th ed.). Indian Council of Agricultural Research.
- Pareek, O. P. (1983). *The Ber*. Indian Council of Agricultural Research.
- Phookan, D. B., Shadeque, A., & Baruah, P. J. (1991). Effect of plant growth regulators on yield and quality of tomato. *Vegetable Science*, 18(1), 93-96.
- Qeyami, M., Bajpay, A., & Jailani, A. W. (2020). Effects of Calcium and Potassium Application on Growth, Yield and Quality of Apple (*Malus x domestica* Borkh.) cv. Red Delicious. *Indian Journal of Pure & Applied Biosciences*, 8(3), 574-584.
- Ram, R. A., & Bose, T. K. (2000). Effect of foliar application of magnesium and micronutrients on growth, yield and fruit quality of mandarin orange (*Citrus reticulata* Blanco). *Indian Journal of Horticulture*, 57(3), 215-220.
- Sarma, B. (2022). *Canopy and Micronutrient Management of Thailand ber (Ziziphus mauritiana Lamk.)* Assam Agricultural University].
- Schanda, J. (Ed.). (2007). *Colorimetry: Understanding the CIE System*. (1st ed.). Wiley-Interscience.

- Schlegel, T. K., & Schonherr, J. (2002). Penetration of calcium chloride into apple fruits as affected by stage of fruit development. *Acta Horticulturae*, 594, 527-533.
<https://doi.org/10.17660/ActaHortic.2002.594.69>
- Sharif, N., Abbas, M. M., Memon, N., & Javaid, M. (2016). Comparative evaluation of naphthalene acetic acid and urea for preventing premature fruit drop and improving fruit yield and quality in ber (*Ziziphus mauritiana* Lamk.) cv. Suffon. *Journal of Agricultural Research*, 54, 55-62.
- Shukla, A. (2011). Effect of foliar application of calcium and boron on growth, productivity and quality of Indian gooseberry (*Embllica officinalis*). *Indian Journal of Agricultural Sciences*, 81(7), 628-632.
- Singh, N., Kaur, A., & Gill, S. B. (2016). Effect of foliar application of zinc and boron on yield and fruit quality of litchi cv. Dehradun. *International Journal of Developmental Research*, 6(7), 8686-8688.
- Singh, R., Godara, N. R., Singh, R., & Dahiya, S. S. (2001). Responses of foliar application of growth regulators and nutrients on ber cv. Umran. *Haryana Journal of Horticultural Sciences*, 30(384), 161-164.
- Singh, S., & Singh, J. P. (2019). Effect of foliar sprays of NAA and boron on flowering, fruiting, fruit retention and yield of litchi (*Litchi chinensis* Sonn.). *International Journal of Chemical Studies* 7(4), 1995-1999.
- Singh, Z., Dhillon, B. S., & Sandhu, A. S. (1991). Relationship of embryo degeneration with fruit drop and its pattern in different cultivars of ber. *Indian Journal of Horticulture*, 48, 277-281.
- Tirkey, N. R., Kanpure, R. N., Kachouli, B. K., Bhandari, J., & Patidar, D. K. (2018). Effect of foliar nutrition of Zinc sulphate, borax and NAA on yield and quality of Guava (*Psidium guajava* L.) cv. Allahabad Safeda. *International Journal of Chemical Studies*, 6(4), 2295-2298.
- Wojcik, P., & Wojcik, M. (2006). Effect of boron fertilization on sweet cherry tree yield and fruit quality. *Journal of Plant Nutrition*, 29(10), 1755-1766.
- Yuan, R., & Carbaugh, D. H. (2007). Effects of NAA, AVG and 1-MCP on ethylene biosynthesis, preharvest fruit drop, fruit maturity, and quality of 'Golden supreme' and 'Golden delicious' apples. *HortScience*, 42, 101-105.