

CHEMICAL CHARACTERIZATION OF SEAWEEDS FROM BIRI, NORTHERN SAMAR, PHILIPPINES

ABSTRACT

This research was carried out in an attempt to ascertain the phytochemical screening on selected seaweed species from different barangays in Biri, Northern Samar, Philippines using a descriptive research approach. It also aimed to identify the seaweed species in the study area and determine the three most and least-frequently appearing species respectively, which were subjected to phytochemical screening and the determination of extracts and its physicochemical constituents as conducted from November 2022 to April, 2023. Seaweed samples were collected, identified, and authenticated by an expert. Specimens were preserved in jars containing 10% formaldehyde and seawater in equal volumes. Fresh samples of the three most frequently appearing and the three least frequently appearing species were used for the screening procedures. Results revealed a total of 44 seaweed species present in selected barangays of Biri, Northern Samar. The most-frequently appearing species were *Ulva reticulata*, *Hormophysacuneiformis*, and *Turbinariaconoides*, while the least-frequently appearing species were *Hydropuntiaedulis*, *Laurenciapapillosa*, and *Tricleocarpafragilis*. Chemical characterization of these species confirmed the presence of alkaloids, saponin, phenolic compounds, and anthraquinones, although not all of them contained these metabolites. Lipids and proteins were absent in all species tested. Meanwhile, the ash content of the samples is relatively high, implying that high amounts of minerals may be found in seaweeds. Thus, the researchers recommend the replication of this research in other areas of the province to document seaweed species and their potential applications in the functional food and nutraceutical industries. This will help develop natural products with economic or commercial significance to the residents. Likewise, more specific tests for lipids and proteins in seaweeds may be used in future investigations.

KEYWORDS: *chemical characterization, seaweeds, chemical constituents, Biri, Northern Samar*

1.0 INTRODUCTION

Seaweeds, or marine macroalgae, are non-flowering, primitive photosynthetic macrophytes found in tidal areas of the seas and oceans that make up 71% of the world's surface [Rao, *et al.*, 2018]. The main groups of seaweeds are brown (*Phaeophyta*), red (*Rhodophyta*), and green (*Chlorophyta*) [MacArtain, *et al.*, 2007]. Moreover, seaweeds can be found worldwide but are restricted by light, nutrition, and a suitable substrate,

[Muth & Congdon, 2019]. Furthermore, a number of seaweed species are edible, and many are also of commercial importance to humans [Britannica, 2022].

Meanwhile, the increasing global population and human activities have negatively impacted the natural environment that leads to shortage of bioresources for the growing population [Jagtap & Meena, 2022]. This has sparked an interest in finding novel foods that will meet the needs of a growing population and offer some health benefits [Granato, *et al.*, 2020; Peñalver, 2020]. Moreover, seaweeds represent a huge resource for mankind, and as the demands grow for special drugs, nutraceuticals, cosmetic products, and functional foods, there is a strong momentum for the exploration of marine biological resources in general, and seaweeds in particular, for novel compounds with health benefits [Qin, 2018]. Thus, recent increased interest in seaweed is motivated by attention generated in their bioactive components that have potential applications in the functional food and nutraceutical industries [Nazarudin, *et al.*, 2021].

The phytochemicals from marine algae are extensively used in various industries such as food, confectionery, textile, dairy, pharmaceutical, and paper, mostly as gelling, stabilizing, and thickening agents [Mohy El-Din & El-Ahwany, 2016]. Seaweeds are the richest source of proteins, lipids, carbohydrates, minerals, vitamins (A, B, C, and Niacin), and antioxidants. They also serve as low-calorie food and are valued as food supplements for people [Rao, *et al.*, 2007]. Generally, seaweeds are known to contain medicinally rich metabolites that include steroids, phenols, tannins, saponins, flavonoids, terpenoids, and glycosides, which have been extensively studied and used in the pharmaceutical industry [Nazarudin, *et al.*, 2021].

Therefore, this present study focuses on the phytochemical screening of selected species of seaweeds in Biri, Northern Samar, specifically, to determine the active chemical constituents of three (3) most-frequently appearing and three (3) least-frequently appearing seaweeds in terms of: alkaloids, anthraquinones, cardiac glycosides, phenolic compounds, and saponin. Furthermore, it will determine the physicochemical activities of the three (3) most-frequently appearing and least-frequently appearing seaweeds in terms of: lipids, proteins, and ash content. No such investigation has yet been done in Biri, Northern Samar, hence, this study.

3.0 MATERIALS AND METHODS

3.1 Area of Study

Northern Samar is situated in the Eastern Visayas region. Its capital is the municipality of Catarman. It is bordered, clockwise from the North, by the Philippine Sea, Eastern Samar, Samar, Samar Sea, and San Bernardino Strait. The province is composed of 24 municipalities (Phil Atlas, 2022).

The island of Biri [Latitude: 12°41'N (12.6816); Longitude: 124°22' (124.3619)], a 5th class municipality, is located in its northernmost tip, facing the Pacific Ocean to the east, the famous San Bernardino Strait to the north, and to either Lavezares or San Jose, both Northern Samar's coastal towns. As of 2020, the current population of the municipality of Biri is 11,274 (Phil Atlas, 2020)., the source of livelihood were farming, fishing, aquaculture, and tourism. The municipality comprises of eight (8) barangays, four (4) of which are located in separate islands.




Figure 1. Map of Study Location

3.2 Sampling Technique

Purposive sampling was applied in the collection of seaweed species of seaweeds utilized for the phytochemical analyses. Specimen for the phytochemical screening were collected purposely, the basis for the collection were the three most frequently occurring seaweeds and the least three occurring seaweeds in the municipality of Biri, Northern Samar.

3.3. Data Collection

The researchers were able to secure a Prior Informed Consent (PIC) from the Municipal Agriculture Office (MAO) and from the municipal mayor and the barangay captain of each sampling site to conduct the study in their area. The researchers went to each barangay for two (2) weekends to collect seaweeds, and each was tallied to determine the three most and three least frequently appearing seaweeds.

3.3.1 Specimen Collection

Specimens were collected by hand-picking and using a knife. Underwater camera for photo documentation and a field notebook were used to record the data for each specimen to serve as guide in identification. Specimens were washed with seawater to remove adhering sediments and impurities, stored in glass jars, and were brought to the College of Science for identification.

3.3.2 Preservation of the Specimens

Each specimen collected was stored in glass jars and soaked in formaldehyde solution at a ratio of 3.7% formaldehyde:96.3% seawater.

3.3.3 Identification of the Specimens

Seaweed samples were initially identified *in situ*, then, to confirm the authenticity of the seaweed samples, it was brought to the College of Science, University of Eastern Philippines

3.4. Seaweed Extraction

The six (6) selected seaweed samples were rinsed using seawater to remove dirt, epiphytes, sediments, and other debris to make sure that it will not affect the result of the phytochemical screening and other chemical test. Manual juicer/ extractor were used for the pure extract, the extracts were then placed and contained in a clean sterile jar for phytochemical screening and characterization.

3.5 Phytochemical Screening

The chemical constituents of the three most- and three least-frequently appearing seaweeds were determined using the following procedures:

3.5.1 Tests for Alkaloids

In this test, the Dragendorff's reagent and Mayer's reagents were used in determining the presence of alkaloids. A positive result was indicated by the presence of an orange precipitate in Dragendorff's reagent, and a white precipitate with the Mayer's reagent (Guevarra, 2005 as cited by Cruz, 2014). Confirmatory tests for alkaloids were also performed using the protocols of Guevarra (2005).

3.5.2 Tests for Anthraquinone

10mL of 10% ammonia solution, few milliliters of filtrate was added and shaken vigorously for 30 seconds. A pink, violet, or red color indicated a positive result of anthraquinone (Shaikh &Patil, 2020).

3.5.3 Test for Cardiac Glycosides

Keller Killiani Test was performed wherein a solution of 0.5ml, containing glacial acetic acid and 2-3 drops of ferric chloride, was mixed with 2 mL of extract. The appearance of deep blue color at the junction of two liquids indicated the presence of cardiac glycosides (Kancherla, *et al.*, 2019).

3.5.4 Test for Phenolic Compounds

Ferric chloride Test was utilized, where ferric chloride solution was added drop wise in a test tube containing 3 mL of the given sample. The color blue, green, violet, or red indicates the presence of phenolic compounds (Vedantu, 2023).

3.5.5 Test for Saponin

An equivalent of 10 g of seaweed extract was loaded to a capillary tube by immersing the tube to a height of 10 mm in the seaweed extract. Likewise, another capillary tube was loaded with distilled water. After sometime, the heights of the liquids in the two tubes were compared. If the level of the seaweed extract in the capillary tube is half or less than in the other tube containing water, then the presence of saponin may be inferred. To confirm its presence in the extract, the frothing test was administered, and a layer of foam of approximately 1cm indicated the presence of saponin (Guevarra, 2005; Cruz, 2014).

3.6 Physicochemical Activities

The different physico-chemical activities of the three (3) most and least frequently appearing seaweeds were determined using the following procedures:

Test for Lipids

Utilizing the emulsion test, a sample was added to 2cm³ of ethanol, and was shaken well, allowed to settle in a test tube rack for 2 minutes, then, any clear liquid was emptied into a test tube containing 2cm³ of distilled water. A layer of milky-white emulsion at the top of the solution indicated the presence of lipids (Guevarra, 2005).

3.6.1. Test for Proteins

The Biuret test was utilized. One to two milliliters of the seaweed solution, egg albumin, and de-ionized water was added in clean and dry test tubes. Then, 1-2 mL of Biuret reagent was added to all the test tubes, shaken well, and allowed to stand for 5 minutes. If the solution turns from blue to deep purple, proteins are present (Aryal, 2022).

3.6.2. Ash content

Oven-dried samples were weighed, and 3 grams of sample was placed in each of 3 separate crucibles. The crucibles were then heated using a

portable gas stove until the sample became completely ash. It was allowed to cool in a desiccator and weighed [Patilan, 201]. Ash content was calculated using the formula below:

$$\text{Ash content} = \frac{\text{weight of ash (g)}}{\text{weight of sample (g)}} \times 100$$

RESULTS AND DISCUSSION

Species Composition of Seaweeds in Biri, Northern Samar

There were forty-four (44) identified species of seaweeds classified into three (3) classes (Chlorophyceae [17 species], Phaeophyceae [13 species], and Rhodophyceae [14 species]). Moreover, the seaweeds were also classified into twenty-one (21) families.

Of this number, the three (3) most-frequently appearing and widely distributed in all of the sampling sites were chosen as samples by the researchers, and they were *Ulva reticulata*, *Hormophysacuneiformis*, and *Turbinariaconoides*. On the other hand, the three least-frequently appearing or those found in only one or two of the sampling sites, were likewise chosen to undergo phytochemical screening and test for the physicochemical constituents. The least frequently appearing seaweed species were *Hydropuntiaedulis*, *Laurenciapapillosa*, and *Tricleocarpafragilis*.

Table 1. **Species Composition of Seaweeds in Selected Brangays of Biri, Northern Samar, Philippines**

Class	Seaweed Species	Sampling Sites				
		1	2	3	4	5
Chlorophyceae	<i>Anadyomeneplicata</i> C. Agardh	x	x	x	/	/
	<i>Avrainvilleaerecta</i> (Berkeley) A. Gepp& E. Gepp	/	x	x	x	x
	<i>Boergeseniaforbesii</i> (Harvey) J. Feldmann	/	/	x	x	/
	<i>Bornetellasphaerica</i> (Zanardini) Solms-Laubach	/	/	x	x	/
	<i>Caulerparachypus</i> Harvey	x	x	x	x	/
	<i>Caulerparacemosa</i> (Forsskål) J. Agardh	/	/	x	/	x
	<i>Caulerpaserrulata</i> (Forsskål) J. Agardh	/	/	x	/	x
	<i>Caulerpataxifolia</i> (Vahl) C. Agardh	/	/	x	x	x
	<i>Chaetomorphacrassa</i> (C. Agardh) Kutzing	/	/	/	/	/
	<i>Dictyosphaeriaversluysii</i> Weber-van Bosse	x	/	x	x	/
	<i>Halicorynewrightii</i> Harvey	x	x	x	x	/
	<i>Halimeda macroloba</i> Decaisne	/	/	/	/	/
	<i>Halimedaopuntia</i> (Linnaeus) Lamoureux	/	/	/	/	/
	<i>Ulva clathrata</i> (Roth)	/	/	x	x	x
	<i>Ulva intestinalis</i> Linnaeus	x	/	x	x	/
	<i>Ulva lactuca</i> Linnaeus	/	/	/	/	/
	<i>Ulva reticulata</i> Forsskål	/	/	/	/	/

Phaeophyceae	<i>Canistrocarpuscervicornis</i> (Kützing) De Paula and De Clerk	/	/	x	/	/
	<i>Dictyotadichotoma</i> (Hudson) Lamoureux	x	/	x	x	x
	<i>Hormophysacuneiformis</i> (J.F, Gmelin) P.C. Sliva	/	/	/	/	/
	<i>Padinagymnospora</i> (Kützing) Sonder	/	/	/	x	/
	<i>Padina minor</i> Yamada	/	/	/	/	/
	<i>Sargassumaquifolium</i>	x	/	x	/	/
	<i>Sargassumcinctum</i>	/	/	x	/	/
	<i>Sargassumgracillimum</i> Reinbold	x	/	x	x	/
	<i>Sargassumhemiphyllum</i> (Turner) C. Agardh	/	/	x	x	x
	<i>Sargassumoligocystum</i> Montagne	/	/	/	/	/
	<i>Sargassumpolycystum</i> C.A. Agardh	/	/	/	x	x
	<i>Sargassumsiliculosum</i> J. Agardh	/	/	x	x	/
	<i>Turbinariaconoides</i> (Turner) J. Agardh	/	/	/	/	/
Rhodophyceae	<i>Acanthaporaspicifera</i> (Vahl) Borgesen	x	/	/	x	/
	<i>Amphiroafofiacea</i> J.V. Lamouroux	/	/	/	/	/
	<i>Coelothrixirregularis</i> (Harvey) Borgesen	/	x	x	x	x
	<i>Eucheumadenticulatum</i> (N.L.Burman) Collins and Harvey	x	x	/	/	x
	<i>Euthoracristata</i> (C. Agardh) J. Agardh	x	x	x	/	x
	<i>Gelidiellaacerosa</i> (Forsskål) Feldmann and Hamel	/	/	x	/	x
	<i>Gracilariasalicornia</i> (C. Agardh) Dawson – with four (4) haplotypes	/	/	/	/	/
	<i>Hydropuntiaedulis</i> (S.G. Gmelin) Gurgel&Fredericq	/	/	x	x	x
	<i>Hypneaservicornis</i> J. Agardh	x	/	/	x	x
	<i>Laurenciapapillosa</i> (C. Agardh) Greville	x	/	x	/	x
	<i>Laurenciatronoi</i> Ganzon-Fortes	/	/	x	/	x
	<i>Mastophorarosea</i> (C. Agardh) Setchell	/	x	x	x	/
	<i>Portieriahornemannii</i> (Lyngbye) P.C. Silva	/	/	x	x	x
	<i>Tricleocarpafragilis</i> (Linnaeus) Huisman& R.A. Townsend	x	/	/	x	x

Legend:
/ - present
x - absent

Sampling sites:
1 - Barangay Progress
2 - Barangay Pio del Pilar
3 - Barangay McArthur
4 - Barangay San Pedro
5 - Barangay San Antonio




Figure 2. The frequently occurring and the least occurring seaweeds for Phytochemical screening. (A). *Ulva reticulata*; (B). *Hormophysacuneiformis*; (C). *Turbinariaconoides*; (D). *Hydropuntiaedulis*; (E).

Phytochemical Screening Results

Results of the phytochemical screening reveal that phenolic compounds were present in all species tested, while saponin was present only in 4 of the samples. Moreover, alkaloids were detected only in *Hormophysacuneiformis* and *Turbinariaconoides*. On the other hand, anthraquinones were present only in *Tricleocarpafragilis*, while all 6 seaweed species tested negative for cardiac glycosides. These results imply that the coastal waters of the municipality of Biri, Northern Samar, specifically the sample barangays, has a diversity of seaweed species which could be harnessed economically through their possession of phytochemicals that could be the bases for drug discovery and development aside from the common use as food or source of alginates.

Table 2. Summary result of phytochemical screening of the three most- and three least-frequently appearing seaweeds in Biri, Northern Samar

SECONDARY METABOLITE TEST	SEAWEED EXTRACTS					
	1	2	3	4	5	6
ALKALOID						
Dragendorff's Test	+	+	+	+	+	+
Mayer's Test	-	+	+	-	-	-
Confirmatory Test	-	+	+	-	-	-
ANTHRAQUINONE	-	-	-	-	-	+
CARDIAC GLYCOSIDES	-	-	-	-	-	-
PHENOLIC COMPOUNDS	+	+	+	+	+	+
SAPONIN						

Capillary Test	-	-	+	+	+	+
Frothing Test	-	-	+	+	+	+

LEGEND:

- + = Tested compound is present in the seaweed extract
- = Tested compound is absent in the seaweed extract

- 1 = *Ulva reticulata*
- 2 = *Hormophysacuneiformis*
- 3 = *Turbinariaconoides*
- 4 = *Hydropuntiaedulis*
- 5 = *Laurenciapapillosa*
- 6 = *Tricleocarpafragilis*

Physicochemical Activities of Seaweed Extracts

The six identified seaweed species were subjected to tests to determine their physicochemical activities or constituents, specifically lipids, proteins, and ash content. Results reveal that lipids and proteins were absent in all of the species tested. However, test for the ash content revealed a relatively high percentage of ash content. This

result implies that there is a potentially high amount of minerals.

Table 3. Summary result of the determination of physicochemical activities of three most and three least frequently appearing seaweeds in Biri, Northern Samar

PHYSICOCHEMICAL ACTIVITIES	SEAWEED EXTRACT					
	1	2	3	4	5	6
LIPID EMULSION TEST	-	-	-	-	-	-
PROTEIN BIURET TEST	-	-	-	-	-	-
ASH CONTENT	22.60%	29.20%	33.07%	33.96%	38.37%	63.17%

LEGEND:

- + = Tested compound is present in the seaweed extract
- = Tested compound is absent in the seaweed extract

- 1 = *Ulva reticulata* 6= *Tricleocarpafragilis*
- 2 = *Hormophysacuneiformis*
- 3 = *Turbinariaconoides*
- 4 = *Hydropuntiaedulis*
- 5 = *Laurenciapapillosa*

CONCLUSION

Utilizing the purposive sampling technique, this descriptive research to set to assess the seaweed resources in the Municipality of Biri, Northern Samar, Philippines. From the results, the three most - and the three least-frequently appearing seaweeds were collected, identified, and subjected to phytochemical screening and determination of its physicochemical constituents.

There were 44 species of seaweeds in five sampling sites in Biri, Northern Samar. These belong to three classes, and 21 families, implying that Biri, Northern Samar harbors a diversity of species of seaweeds.

The three most-frequently appearing seaweeds are: *Ulva reticulata*, *Hormophysacuneiformis*, and *Turbinariaconoides*. Meanwhile, the three least-frequently appearing seaweeds are: *Hydropuntiaedulis*, *Laurenciapapillosa*, and *Tricleocarpafragilis*.

Although not all secondary metabolites were present in any one seaweed species, the most commonly recorded secondary metabolite were phenolic compounds, while the least common metabolite was anthraquinone, which was present in only one sample. This implies that seaweed species available in the study area contain several chemical constituents or secondary metabolites which could have economic significance or potential applications for drug discovery and development.

In terms of the physicochemical activities of the seaweed samples, lipids and proteins turned out to be negative in all samples; however, the ash content of all the samples were relatively high, which implies that the seaweed species tested have a potentially rich mineral content.

Considering the limited tests done in this research, it is recommended that the seaweed extracts may be subjected to further testing to verify the existence of other chemical constituents or secondary metabolites that could have potential medical applications and/or can be used to create natural products.

The use of other methods of preparations, tests, and analyses for phytochemical screening and physicochemical constituent determination should be done to further test the availability of secondary metabolites and physicochemical constituents on seaweeds.

Consent

As per international standards or university standards, respondents' written consent has been collected and preserved by the author(s).

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and

source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

- Abobo, Tom Jericho L. 2018. Species Composition of Seaweeds in Palapag, Northern Samar. Undergraduate Thesis College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Alkaloid. Britannica. 2022. <https://www.britannica.com/science/alkaloid> Accessed: October 24, 2022
- Alu'datt, M.H., T. Rababah, M.N. Alhamad, G.J. Al-Rabadi, C.C. Tranchant, A. Almajwal, & A. Inteaz. 2017. Occurrence, types, properties and interactions of phenolic compounds with other food constituents in oil bearing plants. Critical Reviews in Food Science and Nutrition. 58(18): 3209-3218. doi:<https://doi.org/10.1080/10408398.2017.1391169> Retrieved: October 27, 2022
- Anthraquinone. PubChem. 2022. <https://pubchem.ncbi.nlm.nih.gov/compound/Anthraquinone> Retrieved: October 24, 2022
- Aryal, Sagar. 2022. Biuret Test for Protein- Definition, Principle, Procedure, Results, Uses. Microbe Notes. <https://microbenotes.com/biuret-test-for-protein/> Accessed: November 8, 2022

- Babenko, L.M., O.E. Smirnov, K.O. Romanenko, O.K. Trunova, I.V. Kosakivska. 2017. Phenolic compounds in plants: biogenesis and functions. Ukr. Biochem. J. 91(3): 5-18. doi: <https://doi.org/10.15407/ubj91.03.005>. Accessed: October 27, 2022
- Baldia, Susana F., Mark Timothy C, Kabling, Connie Anne L. Gabriel, Angelique Nadine D.G. Abeleda, Rose Celine A. Aguinaldo, Ken Joseph E. Clemente, & Paciente A. Cordero Jr. 2017. Marine macroalgal diversity assessment of Biri Island and Dalupirit Island, Northern Samar, Philippines. ActaManilana: 65, pp. 29-38. doi:<https://doi.org/10.53603/actamanil.65.2017.vmbh5085> Retrieved: October 2, 2022
- Biri Profile. 2010. Municipal Planning and Development (MPDC).
- Brown algae. 2022. Britannica. <https://www.biologyonline.com/dictionary/brown-algae> Accessed: November 17, 2022
- Capoquian, Nerissa C. 2010. Inventory Assessment of Seaweeds in Gamay, Northern Samar. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Carbohydrate. Britannica. 2022. <https://www.britannica.com/science/carbohydrate> Retrieved: October 24, 2022
- Cox-Georgian, D., N. Ramadoss, C. Dona, C. Basu. 2019. Therapeutic and Medicinal Uses of Terpenes. Medicinal Plants. pp. 333-359. doi:https://doi.org/10.1007/978-3-030-31269-5_15 Accessed: October 27, 2022
- Cruz, Mark Gil M. 2014. Phytochemical Screening of the Ethanol Extracts of *Gliricidlasepium* (Jacq.) Steud. (Kakawate). Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Dhargalkar, V.K., D. Kavlekar. 2004. Seaweeds – a field manual, X.N. Verlecar, V. Rathod (Eds). National Institute of Oceanography. Dona Paula, Goa – 403 004.
- Deyab, M., T. Elkatory, & F. Ward. 2016. Qualitative and Quantitative Analysis of Phytochemical Studies on Brown Seaweed, *Dictyota dichotoma*. International Journal of Engineering Development and Research. <https://www.researchgate.net/publication/311717544> Retrieved: October 24, 2022
- De la Rosa, L.A., J.O.M. Escamilla, J.R. Garcia, & E.A. Parilla. 2019. Phenolic Compounds. In: Postharvest Physiology and Biochemistry of Fruits and Vegetables, E.M. Yahia (Eds). Woodhead Publishing. Chapter 12, pp.253-271. <https://doi.org/10.1016/B978-0-12-813278-4.00012-9>. Retrieved: February 21, 2023
- Elmegeed, Dalia F., Doaa A. Ghareeb, Muhammed Elsayed, & Muhammad El-Saadani. 2014. Phytochemical constituents and bioscreening activities of green algae (*Ulva lactuca*). International Journal of Agricultural Policy and Research: 2 (11), pp. 373-378. doi: <http://dx.doi.org/10.15739/IJAPR.009> Accessed: October 2, 2022
- Food test 3 - Emulsion (ethanol) test for Fats. 2022. Biology Notes. <https://biology-igcse.weebly.com/-food-test-3-emulsion-ethanol-test-for-fats.html> Retrieved: November 8, 2022

- Galezoga, Divina M. 2019. Seaweed Resources in the Island Town of Biri, Northern Samar, Philippines. University of Eastern Philippines, Catarman, Northern Samar.
- Gangasani, J.K., D.B. Pemmaraju, U.S.N. Murthy, A.K. Rengan, V.G.M. Naidu. 2022. Chemistry of Biomolecules. In: Herbal Biomolecules in Healthcare Applications, S.C. Mandal, A.K. Nayak, A.K. Dhara (Eds). Academic Press. Chapter 4, pp. 63-79. <https://doi.org/10.1016/B978-0-323-85852-6.00018-4>. Retrieved: October 27, 2022
- Green algae. 2022. Britannica. <https://www.britannica.com/science/green-algae> Accessed: November 17, 2022
- Guasis, John Paulo E. 2016. Species Composition of Seaweeds in Mondragon, Northern Samar. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Heinrich, Michael, Jeffrey Mah, & Vafa Amirkia. 2021. Alkaloids Used as Medicines: Structural Phyto-chemistry Meets Biodiversity – An Update and Forward Look. *Molecules*. 26(7): 1836. <https://doi.org/10.3390/molecules26071836> Retrieved: May 16, 2023
- Imran, Md. A.S., F.R. Bhuiyan, S.R. Ahmed, P. Shanzana, M.A. Moloi, S.H. Foyosal, S.B. Dabi, & M. Hasan. 2021. Phytochemical constituency profiling and antimicrobial activity screening of seaweeds extracts collected from the Bay of Bengal Coasts. *Journal of Advanced Biotechnology and Experimental Therapeutics*: 4(1), 25- 34. doi: <https://doi.org/10.5455/jabet.2021.d103> Retrieved: October 2, 2022
- Jagtap, A. S. & S. N. Meena, 2022. Seaweed farming: A perspective of sustainable agriculture and socioeconomic development. In: *Natural Resources Conservation and Advances for Sustainability*, M.K. Jhariya, R.S. Meena, A. Banerjee, & S. N Meena (Eds). Elsevier. <https://doi.org/10.1016/B978-0-12822976-7.00022-3> Accessed: November 8, 2022
- Kancherla, N., A. Dhakshinamoothi, K. Chitra, and R.B., Komaram. 2019. Preliminary Analysis of Phytoconstituents and Evaluation of Anthelmintic Property of *Cayratia auriculata* (In Vitro). *Journal of Clinical Medicine*. 14(4): 350-356 <https://doi.org/10.26574/maedica.2019.14.4.350> Retrieved: April 26, 2023
- Khan, Nida Tabassum. 2019. ANTHRAQUINONES – A Naturopathic Compound. *Journal of New Developments in Chemistry*. 2(2): 25-28 doi: <https://doi.org/10.14302/issn.2377-2549.jndc-18-2569> Accessed: October 27, 2022
- Lin, Derong, Mengshi Xiao, Jingjing Zhao, Zhuohao Li, Baoshan Xing, Xindan Li, Maozhu Kong, Liangyu Li, Qing Zhang, Yaowen Liu, Hong Chen, Wen Qin, Hejun Wu, & Saiyan Chen. 2016. An Overview of Plant Phenolic Compounds and Their Importance in Human Nutrition and Management of Type 2 Diabetes. *Molecules*. 21(10): 1374 <https://doi.org/10.3390/molecules21101374> Retrieved: May 16, 2023
- Lipid. Britannica. 2022. <https://www.britannica.com/science/lipid> Retrieved: October 24, 2022

- MacArtain, P., C.I.R. Gill, M. Brooks, R. Campbell, & I.R. Rouland. 2007. Nutritional Value of Edible Seaweeds. *International Life Sciences Institute*. doi: <https://doi.org/10.1301/nr.2007.dec.535-543> Retrieved: October 24, 2022
- Measuring moisture content: It's more complicated than you think. 2022. Meter Group https://www.metergroup.com/en/meter_food/moisturecontent Accessed: November 8, 2022
- Mengullo, Lynette G. 2001. An Inventory Assessment of Seaweeds in Gamay, Northern Samar. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Mohy El-Din, S.M. & M.D. El-Ahwany. 2016. Bioactivity and Phytochemical Constituents of Marine Red Seaweeds (*Jania rubens*, *Corallinamediterranea*, and *Pterocladia capillacea*). *Journal of Taibah University for Science*:10(4), pp. 471-484. doi: <https://doi.org/10.1016/j.jtusci.2015.06.004> Accessed: October 24, 2022
- Moisture. Cambridge Dictionary. 2022. <https://dictionary.cambridge.org/us/dictionary/english/moisture> Retrieved: October 24, 2022
- Muth, A.F. & M.V. Congdon. 2019. Seaweeds and Seagrasses. In: *Encyclopedia of Ocean Sciences* (Third Edition), J.K. Cochran, H.J. Bokuniewicz, & P.L. Yager (Eds). Academic Press. pp. 765-774. <https://doi.org/10.1016/B978-0-12409548-9.11265-5> Accessed: November 8, 2022
- Nazarudin M.F., N.H. Alias, S. Balakrishnan, W.N.I. Wan Hasnan, N.A.I. Noor Mazli, M.I. Ahmad, I.S. Md Yasin, A. Isha, M. Aliyu-Paiko. 2021. Chemical, Nutrient and Physicochemical Properties of Brown Seaweed, *Sargassum polycystum* C. Agardh (Phaeophyceae) Collected from Port Dickson, Peninsular Malaysia. *Molecules*. 26(17): 5216. doi: <https://doi.org/10.3390/molecules26175216>. Accessed: October 31, 2022
- Nofies, Sandme B. 2007. Inventory Assessment of Seaweeds in Selected Barangays of Palapag, Northern Samar. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, CatarmanNorthernSamar.
- Northern Samar Profile. 2022. Phil Atlas. https://www.philatlas.com/isayas/r08/northern_samar.html Retrieved: November 17, 2022
- Patilan, Jeneky E. 2017. Nutritional and Nutraceutical Content of *Sauropusandrogynus* (Linn.) Merr. (Chinese Malunggay) Leaf. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.
- Peñalver, R., J.M. Lorenzo, G. Ros, R. Amarowicz, M. Pateiro, & G. Nieto. 2020. Seaweeds as a Functional Ingredient for a Healthy Diet. *Mar Drugs*. 18(6): 301. <https://doi.org/10.3390/md18060301> Retrieved: November 8, 2022
- Phytochemical Screening. *American Journal of Phytomedicine and Clinical Therapeutics*. 2022.

<https://www.imedpub.com/scholarly/phytochemicalscreening-journals-articles-ppts-list.php> Retrieved: October 24, 2022

- Protein. Merriam-Webster Dictionary. 2022. <https://www.merriamwebster.com/dictionary/protein> Accessed: October 24, 2022
- Qin, Yimin. 2018. Health Benefits of Bioactive Seaweed Substances. In: Bioactive Seaweeds for Food Applications. Academic Press. Chapter 9, pp. 179-200. <https://doi.org/10.1016/B978-0-12-813312-5.00009-1> Retrieved: November 8, 2022
- Qiu, S., H. Sun, A.H. Zhang, H.Y. Xu, G.L. Yab, Y. Han, X.J. Wang. 2014. Natural alkaloids: basic aspects, biological roles, and future perspectives. Chin J Nat Med. 6: 401-6. doi: [https://doi.org/10.1016/S1875-5364\(14\)60063-7](https://doi.org/10.1016/S1875-5364(14)60063-7) Retrieved: October 27, 2022
- Roy, S., B. Nayak, M. Roy, A. Mitra, D. Halder, A. Chatterjee, S. Chakraborty, & T. Chakraborty. 2016. Antimicrobial and Phytochemical Screening of Seaweeds: Enteromorpha intestinalis and Ulva lactuca Collected from Indian Sunderbans Delta Region. IOSR Journal of Pharmacy and Biological Sciences: 11 (4), pp. 01-05. <https://www.researchgate.net/publication/305618118> Retrieved: October 2, 2022
- Red algae. 2022. Britannica. <https://www.britannica.com/science/red-algae>. Accessed: November 17, 2022
- Seaweed. Britannica. 2022. <https://www.britannica.com/science/seaweed>. Accessed: October 24, 2022
- Seenivasan, R. M. Rekha, H. Indu, S. Geetha. 2012. Antibacterial Activity and Phytochemical Analysis of Selected Seaweeds from Mandapam, Coast India. Journal of Applied Pharmaceutical Science: 2 (10), pp. 159-169 doi: <https://doi.org/10.7324/JAPS.2012.21030> Accessed: October 2, 2022
- Shaikh, J.R., M.K. Patil. 2020. Qualitative tests for preliminary phytochemical screening: An overview. International Journal of Chemical Studies. 8(2): 603-608. <https://doi.org/10.22271/chemi.2020.v8.i2i.8834> Accessed: April 14, 2023
- Souza, Cassio R.M., Wallace P. Bezerra, & Janeusa T. Souto. 2020. Marine Alkaloids with Anti-Inflammatory Activity: Current Knowledge and Future Perspectives. Mar Drugs. 18(3): 147. <https://doi.org/10.3390/md18030147> Retrieved: May 16, 2023
- Subba Rao, P.V., C. Periyasamy, K. Suresh Kumar, A. Srinivasa Rao, & D. Anantharaman. 2018. Seaweeds: Distribution, Production and Uses. In: Bioprospecting of Algae, M.N. Noor, S.K. Bhatnager, & S.K. Sinha (Eds). Chapter 6, pp. 59-78.
- Subba Rao, P.V., K. Ganesan, & K. Suresh Kumar. 2009. Seaweeds: A Survey of Research and Utilization. In: Algal Biology and Biotechnology, J.I.S. Kattar,
- D.P. Singh, & I.K. Gurpreet Kaur (Eds). International Publishing House, PVT Ltd. New Delhi, Bangalore. pp. 165-178.
- Subba Rao, P.V., V.A. Mantri, K. Ganesan, & K. Suresh Kumar. 2007. Seaweeds as a human diet: An emerging trend in the new Millenium. In: Advances in Applied Phycology, R.K. Gupta & V.D. Pandey (Eds). Daya Publishing House, New Delhi. pp. 85-96.

Surio, Randy M. 2008. Distribution and Abundance of Seaweeds in Selected Barangays of Pambujan, Northern Samar. Undergraduate Thesis. College of Science, University of Eastern Philippines, University Town, Catarman, Northern Samar.

Tests for Carbohydrates. 2022. A-Level Biology. Retrieved: <https://alevelbiology.co.uk/notes/tests/for/carbohydrates/> November 8, 2022

Vedantu. 2023. Test for Phenolic Group. <https://www.vedantu.com> Accessed: April 25, 2023

Whitehead, K., J.I. Hedges. 2005. Photodegradation and Photosensitization of Mycosporine-like amino acids. J Photochem Photobiol B: Biol. 80: 115-121.

Yang, Y., S. Laval, & B. Yu. 2021. Chemical Synthesis of Saponins. In: Advances in Carbohydrate Chemistry and Biochemistry, D.C. Baker (Eds). Academic Press. Chapter 2 (79), pp. 63-150. <https://doi.org/10.1016/bs.accb.2021.10.001>. Accessed: February 21, 2023

UNDER PEER REVIEW