

ASSESSING THE EARLY ESTABLISHMENT AND ADAPTABILITY OF ITALIAN RYEGRASS (*LOLIUM MULTIFLORUM*) IN IRRIGATED AND RAINFED CONDITION OF PUNJAB, PAKISTAN

ABSTRACT

In Punjab, Pakistan, more than 60% expenditure is incurred on the feeding of livestock and offered with low quality forage thus affecting the productivity of milk and meat. The less production of livestock oriented products often results in malnutrition in the society. An effort was made to evaluate the Italian Ryegrass (*Lolium multiflorum*) early grows and adapts to both rainfed and irrigated environments. The main objective of this study is to provide important new information for sustainable fodder production systems. The experiment used a Randomized Complete Block Design with seeds obtained from Korea and was carried out in the winter of 2023 in both rainfed Chakwal and irrigated Sargodha. The plant's response to varying soil conditions and water availability was evaluated by measuring a range of growth parameters, such as plant height, population density, root and shoot biomass, root length, and the root-shoot ratio. The results show that there are significant differences between rainfed and irrigated locations established as by statistical analysis. Italian Ryegrass in irrigated plots showed higher mean plant height (43.75 cm), higher plant population (317.50 plants/m²), higher shoot biomass (7.425 g/plant), greater length of roots (5.950 cm), and a lower root-shoot ratio (0.1135) than in rainfed areas. The significance of these differences was established by statistical analysis. Additionally, the study provides data on soil profiles that show differences between the two locations in terms of pH, electrical conductivity, nitrogen, phosphorus, and potassium. The findings provide important information for enhancing Italian Ryegrass cultivation in Punjab's various agro-ecological zones, given the critical role fodder feed plays in supporting livestock and agriculture. The study provides the basis for well-informed farming practices by highlighting the importance of water management strategies to improve fodder production during scarcity period and, by consequently, animal nutrition in the area.

Keywords: Adaptability, Biomass, Fodder, Irrigation and Rainfed zone, *Lolium multiflorum*, Punjab, Pakistan

INTRODUCTION

Pakistan's livestock sector is critical to the national economy, contributing significantly to GDP and rural livelihoods. Forage availability is still a significant barrier, though, especially in dry and semi-arid areas like Punjab (Rehman et al., 2017). The productivity and sustainability of

Comment [H1]: Insert

Comment [H2]: delete

Comment [H3]: delete

Comment [H4]: insert

Comment [H5]: low

Comment [H6]: insert

Comment [H7]: This study aims to evaluate the early growth and adaptability of Italian Ryegrass (*Lolium multiflorum*) in both rainfed and irrigated environments

Comment [H8]: The primary objective is to provide important new information for sustainable fodder production systems.

Comment [H9]: The experiment was conducted using a Randomized Complete Block Design with seeds obtained from Korea, and it took place in the winter of 2023 in both rainfed Chakwal and irrigated Sargodha.

Comment [H10]: showed

Comment [H11]: insert

Comment [H12]: exhibited

Comment [H13]: root length

Comment [H14]: compared to

Comment [H15]: revealing

Comment [H16]: are crucial

Comment [H17]: highlights

Comment [H18]: thereby improving

Comment [H19]: region.

traditional fodder crops like maize and sorghum are threatened by temperature swings and a growing lack of water (Kumar et al., 2023). This makes it necessary to investigate durable, alternate forage choices such as temperate grasses that can yield high-quality feed all year round. Punjab, Pakistan's goal of sustainable agriculture necessitates the development of resilient forage crops that may flourish under a variety of agroclimatic circumstances (Sustek et al., 2023). Using its quick growth and high nutritional content, Italian ryegrass (*Lolium multiflorum*) is a viable way to increase forage availability. This cool-season grass species is a focus of agro ecological research because of its widespread recognition for adaptability and appropriateness as animal feed (Seydosoglu et al., 2021).

Comment [H20]: delete

Comment [H21]: Italian ryegrass (*Lolium multiflorum*) is a viable option for increasing forage availability due to its quick growth and high nutritional content.

Comment [H22]: suitability

Comment [H23]: can potentially grow

Italian ryegrass (*Lolium multiflorum*) is an important forage crop that may grow and adapt well in Punjab, Pakistan's rainfed and irrigated regions. The purpose of this study is to evaluate Italian ryegrass's development potential and adaptation in these various agricultural zones. Due to its good nutritious content, high herbage yield, favorable performance, and capacity for regrowth, Italian ryegrass can be harvested numerous times during its life cycle (Maxwell et al., 2019). Furthermore, it has been discovered that the biomass of Italian ryegrass degrades slowly; after nine weeks after planting (WAP), almost two-thirds of the plant waste was still present (Ji et al., 2018). Additionally, it has been determined that Italian ryegrass is a significant feed crop grown in South Africa, where it is cultivated under irrigated condition for intensive production of dairy, lamb, and cattle (Cavazzana et al., 2023). The adaptability of Italian ryegrass to poorly drained circumstances makes it a viable winter cover crop for rotation with irrigated rice, as noted in some agricultural practices (Goulart et al., 2020). Additionally, research on the morphological and physiological variations between diploid and tetraploid cultivars under salinity stress conditions has been done on Italian ryegrass in relation to its response to salinity stress (Akinroluyo et al., 2019). For the purpose of evaluating Italian ryegrass's capacity for growth in a variety of agricultural environments, these studies offer insightful information about how adaptable the plant is to various environmental stresses (Xie et al., 2020). Italian ryegrass has also been studied for its allelopathic potential, especially in regard to its effect on other crops' germination. Studies have shown that various winter cover crop shoot extracts, such as Italian ryegrass, affect different crops' seed germination rates, suggesting that these extracts may have allelopathic effects on nearby plants. For the purpose of evaluating Italian ryegrass's adaptability and possible influence on agroecosystems, it is imperative to comprehend its allelopathic interactions (Villela et al., 2021).

Comment [H24]: multiple

Comment [H25]: delete

Comment [H26]: These studies offer insightful information about its adaptability to various environmental stresses for the purpose of evaluating the capacity of Italian ryegrass's for growth in a variety of agricultural environments (Xie et al., 2020).

Comment [H27]: (Villela et al., 2021).

Further research on Italian ryegrass has examined its involvement in the cycling of nutrients and possible environmental effects in the context of nitrogen removal and nitrate leaching in pasture systems (Smit et al., 2021). These studies add to the thorough evaluation of the adaptability and growth capacity of Italian ryegrass in various agricultural systems. The importance of Italian ryegrass is derived from its agronomic qualities. Because of its quick establishment, fast growth, and nutritious content, it's a desirable choice for raising animal productivity. Furthermore, because of its robust root structure, it helps to prevent erosion, which

Comment [H28]: Further studies have examined its involvement in the cycling of nutrients and potential environmental effects in the context of nitrogen removal and nitrate leaching in pasture systems (Smit et al., 2021).

emphasizes its potential importance in maintaining agricultural landscapes (Scordia et al., 2019; Xie Y et al., 2021).

The agricultural landscape of Punjab is divided into areas with different patterns of water availability. Sargodha serves as an example of the irrigated zones, which gain from managed water supplies that guarantee steady crop development (Nadeem et al., 2023). In contrast, Rainfed zones, like Chakwal, on the other hand, depends on the erratic moisture which affects agricultural output (Sheikh et al., 2023). Although the potential of Italian-ryegrass has been acknowledged, but there still exists a crucial study gap concerning a thorough assessment of its development dynamics in these agro-ecological zones need to be explored, particularly in places that receive rain and artificial irrigation (Abbas et al., 2017). In order to measure the Italian rye grass potential, this study measures the growth performance of Italian ryegrass at an early stage, specifically 50 days after seeding. The precise goals include calculating the biomass weights of the roots and shoots, measuring plant height, estimating plant population density, and assessing root length in both rainfed and irrigated zones.

The chosen parameters are important indicators to assess the initial growth and adaptation of Italian ryegrass (Xie Y et al., 2021). Plant height signifies growing vigor, but plant population density represents establishment success in plants. The biomass weights of the roots and shoots indicate the distribution of resources, while the length of the roots indicates the plant's capacity to pull moisture and nutrients from the soil, which is essential for adaptation in a variety of agro-ecological zones. The purpose of this study is to determine the growth potential and adaptability of Italian ryegrass under different moisture regimes by comparing zones that are irrigated and those that are rainfed. This research attempts to provide insights into the crop's response to various environmental situations, by carefully measuring and observing these parameters, therefore guiding forage production of Italian rye grass in Punjab-Pakistan.

METHODOLOGY

The aim of this research is to contribute the prospective integration of Italian ryegrass into fodder production systems in a variety of Punjab province ecosystems by offering useful data on its growth and development under various water regimes and soil conditions. An experiment was carried out in the winter of 2023 in two locations: Sargodha (irrigated) and Chakwal (rainfed) in order to assess the growth potential and adaptability of Italian ryegrass (*Lolium multiflorum*) in disparate agro-ecological zones of Punjab, Pakistan. At every location, used a Randomized Complete block design with three replications for each Plot was used, and total plots was four, each plots were 505 m² (1 kanal). The seeds were arranged from OAK Café Company, Korea. (Confirm the name of company from net) Korean-sourced seeds were drilled at a rate of 5 kg/ha. In Sargodha, two irrigations were applied: 1st just after sowing and 2nd after 40 days after 1st irrigation. While in Chakwal no artificial irrigation applied in Chakwal. Different fertilizer applications were made: Chakwal received one bag of DAP per 4 kanals, while Sargodha received one bag of NPK per 4 kanals. See the field assistant to confirm Chakwal fertilizer application.

Comment [H29]: benefiting

Comment [H30]: insert

Comment [H31]: delete

Comment [H32]: delete

Comment [H33]: insert

Comment [H34]: delete

Comment [H35]: delete

Comment [H36]: To address this gap

Comment [H37]: delete

Comment [H38]: delete

Comment [H39]: delete

Comment [H40]: insert

Comment [H41]: with a total of four plots

Comment [H42]: measuring

Comment [H43]: sourced

Comment [H44]: the first

Comment [H45]: the second

Comment [H46]: delete

Comment [H47]: first

Comment [H48]: delete

Comment [H49]: In

Comment [H50]: delete

Comment [H51]: delete

Data collection

Plant Population (m²):

To determine the average plant population for a given plot, the number of plants inside a 1 square meter quadrat in several sample sections of each plot were counted.

Comment [H52]: was

Height of Plant (cm):

Measured the height of ten randomly chosen plants in each plot using a measuring tape or ruler, starting from the base of the stem and ending at the tip of the tallest leaf. Next, the average height for every plot was calculated.

Comment [H53]: The average height for every plot was then calculated.

Root and Shoot Biomass per plant:

Ten selected at random plants from each plot should be carefully uprooted at the specified sample. After carefully washing the roots in order to remove the soil, the roots and shoots were separated. To ensure complete drying, the samples were placed in an oven set to the optimal temperature of 105°C for a suggested 48-hour period. The dry weight of roots and shoots sample was measured in grams using a balance after they had fully dried and calculated with formula equation 1, 2 and obtained the biomass of root and shoot and thus derived the water holding ability of roots and shoots. The PMAS Arid Agriculture University's central lab in Rawalpindi was the site of this standard operating procedure.

Comment [H54]: were

Comment [H55]: After washing the roots to remove soil

Comment [H56]: The samples were then placed in an oven set to 105°C for 48 hours to ensure complete drying

Comment [H57]: delete

Comment [H58]: to obtain

Root biomass=Weight of dried roots (g)–Weight of fresh roots (g)..... eq.1

Shoot biomass=Weight of dried shoots (g)–Weight of fresh shoots (g)..... eq. 2

Root Length:

Using measuring tape, calculate the average root length from each of the ten plants that were sampled in each plot. For each plot, the average root length possible was recorded.

Comment [H59]: delete

Comment [H60]: sampled plants in each plot was calculated and recorded.

Root-Shoot Ratio:

The root-shoot ratio was calculated using the obtained biomass data by eq. 3:

Root-Shoot Ratio = Root Biomass/Shoot Biomass..... eq 3

This ratio served as a measure to determine the way resources were distributed between the root and shoot systems. Following the drying process, the measured biomass of the root and shoot components was used to calculate the amount. At PMAS Arid Agriculture University, Rawalpindi, the central lab strictly adhered to the established protocols for measurements and computations.

Comment [H61]: distribution of resources between the root and shoot systems

Comment [H62]: Measurements and computations strictly adhered to the established protocols at PMAS Arid Agriculture University, Rawalpindi

Soil profile

The soil profile data collection utilized sensors from the Precision Agriculture Lab of Arid Agriculture University. Five samples were gathered from diverse locations, and readings were meticulously recorded and analyzed for key parameters such as electrical conductivity (EC), pH levels, nitrogen (N), phosphorus (P), and potassium (K). Average values for each parameter were computed for both sites.

Table 1. Comparison of soil characteristics between irrigated (Sargodha) and rainfed (Chakwal) zones:

Parameter	Sargodha (Irrigated)	Chakwal (Rainfed)	Units
Electrical Conductivity (EC)	< 4 dS m ⁻¹	< 4 dS m ⁻¹	dS m ⁻¹
pH	8.4	8.2	-
Nitrogen (N)	134 mg kg ⁻¹	96 mg kg ⁻¹	mg kg ⁻¹
Phosphorus (P)	7.1 mg kg ⁻¹	4.0 mg kg ⁻¹	mg kg ⁻¹
Potassium (K)	104 mg kg ⁻¹	84 mg kg ⁻¹	mg kg ⁻¹

Data Analysis

The growth statistics in the rainfed and irrigated treatments had mean values determined by statistical analysis ANOVA or t-tests were employed to identify significant differences among treatments. Relationships between variables were investigated using regression and correlation analysis. To ensure reliable conclusions, LSD was also used to identify significant differences between treatment means.

RESULTS & DISCUSSIONS

The evaluation of Italian Ryegrass (*Lolium multiflorum*) in rainfed and irrigated conditions in Punjab, Pakistan, showed significant variations in various type of growth parameters.

Plant Height (cm):

There were significant differences in the average plant height of Italian Ryegrass (*Lolium multiflorum*) between the two treatment methods rainfed and irrigated condition. The mean plant height measured in the irrigated plots was 43.75 cm (Table 2), which was notably more than the 13.75 cm observed in the rainfed plots. This notable 30-cm variance emphasizes that water supply has a significant impact on the vertical growth of the grass species. By evaluating the accuracy of these measures using standard deviation (SD) values, the consistency of each treatment is underscored. The standard deviation of plant height in the irrigated areas was 1.70,

while it showed a **same** variation in the rainfed plots at 1.70 (Table 2). This indicates that, despite the significant variation in mean plant height, there was a very uniform growth pattern within each treatment. The relative variability within each treatment is represented by the coefficient of variation (CV), which is calculated as the ratio of SD to the mean. The plant height CV under irrigating conditions was found to be 3.90%, but it was significantly greater (12.43%) under rainfed conditions. This increased CV (Table 3), under rainfed conditions suggests that there is more variability because of the limited supply of water, as evidenced by the comparatively wider distribution of data points about the mean plant height. The 30 cm difference in plant height between rainfed and irrigated environments shows the significant impact of water availability on upward development. Increase in Plant height in irrigated plots highlights the need for sufficient water resources for ideal growth, which is essential for optimizing fodder yield (Saleh et al., 2023).

Comment [H63]: similar

Plant Population or Plant Density (m²)

The evaluation of the population of Italian ryegrass (*Lolium multiflorum*) plants under various circumstances showed distinct trends. The mean plant population in the irrigated plots was determined to be 317.50 plants/m², whereas in rainfed conditions, the mean population decreased to 240.0 plants/m² (Table 2). The difference indicates that the establishment and density of Italian Ryegrass **are** significantly influenced by the availability of water. The variety within the populations was further highlighted by the standard deviation (SD) figures. The rainfed plots displayed a little larger SD of 21.602 than the irrigated plots (Table 2), which displayed an SD of 17.078, respectively. There are more noticeable differences in plant density between the plots as a result of the increased fluctuation in rainfed conditions, which highlights how susceptible plant populations are to water stress. Coefficient of variation (CV) calculations supported these results, with the rainfed condition having an approximate higher CV of 9.00% and the irrigated condition showing a CV of 5.38%. This increased CV in the rainfed condition indicates a comparatively larger distribution of plant populations, suggesting that Italian Ryegrass is more susceptible to variations in water availability. P-value statistical analysis further supported the substantial difference ($P < 0.05$) in plant population between the two treatments. When everything is considered, the mean values, along with SD, CV, and statistical analysis, highlight that the populations of Italian Ryegrass plants under irrigated and rainfed environments differ significantly. The important effect that water availability has on Italian Ryegrass growth is one of the main findings (Pornaro et al., 2020). Under irrigated conditions, the plant showed larger mean plant heights, higher population densities, and more biomass in the roots and shoots when water was readily accessible. A more consistent and resilient growth pattern was the outcome of careful irrigation management in Sargodha, highlighting the beneficial relationship between plant development and water availability (Razzaq et al., 2023). Italian Ryegrass is sensitive to water stress, which affects its establishment and abundance. This **is** demonstrated by the observable decrease in plant population under rainfed conditions (Pan et al., 2016). This result

Comment [H64]: was

Comment [H65]: was

highlights how important it is to have an adequate water supply that promotes larger plant densities, which are necessary for strong fodder production (Byrne et al., 2018).

Root & Shoot Biomass Weight (g/plant):

A critical indicator of the Italian Ryegrass's (*Lolium multiflorum*) reaction to various conditions involves the plant's total weight of roots produced. The mean root biomass weight in the irrigated plots was 0.84 g/plant (Table 2), indicating an effective root system under ideal water availability. A low standard deviation of 0.059 (Table 2), which denotes a significant level of consistency in the data, supported this. The irrigated treatment exhibited a comparatively low degree of variability, as indicated by the coefficient of variation (Cv) of 7.01%. The rainfed plots, on the other together, showed a lower mean root biomass weight of 0.44 g per plant, highlighting the water stress shown negative effect on the production of biomass. In comparison to the irrigated plots, the rainfed treatment's standard deviation of 0.059 indicates a marginally higher variability. There **is** a moderate amount of variability within the rainfed treatment, as indicated by the computed coefficient of variation of 13.36% (Table 3). The statistical significance ($p < 0.05$) of the observed difference between the irrigated and rainfed treatments highlights the impact of water availability on the root biomass of Italian Ryegrass. The increased mean root biomass in the irrigated condition highlights the significance of water management in maximizing biomass production in forage crops, as well as the beneficial effects of irrigation on below-ground growth.

Comment [H66]: was

In the research of Italian Ryegrass (*Lolium multiflorum*) in Punjab, Pakistan, there were significant differences between the irrigated and rainfed conditions in terms of the Shoot Biomass Weight, a critical indication of the productivity of above-ground plants. The mean Shoot Biomass Weight per plant in the irrigated plots was 7.42 g, indicating a strong above-ground biomass output under optimum water availability (Table 2). The rainfed plots, **on the other hand**, showed a significantly reduced mean shoot biomass weight per plant (2.97 g), indicating the effect of water constraint on the ability of the species to divide resources for shoot development. These observations **were confirmed** by statistical analysis. While the standard deviation under rainfed conditions was slightly higher at 0.3862 g/plant, suggesting greater variability in shoot biomass due to water stress, the standard deviation in irrigated conditions was 0.59 g/plant, indicating generally consistent biomass output across plots (Table 2). This pattern was further shown by the Coefficient of Variation (CV), which showed that in irrigated conditions the CV was 7.96% but under rainfed conditions it was 12.96% (Table 3). A paired t-test was used to evaluate the significance of the observed differences, and the results showed a p-value that was less than the specified significance limit ($\alpha = 0.05$). This lends weight to the idea that there are statistically significant differences in Shoot Biomass Weight between the rainfed and irrigated conditions. In conclusion, the Shoot Biomass Weight parameter reflects the substantial impact of irrigation on the above-ground biomass production of Italian Ryegrass. The significant difference in biomass weights of roots and shoots between rainfed and irrigated conditions highlights the reliance biomass production **is** on water availability (Caiger et al.,

Comment [H67]: delete

Comment [H68]: and

Comment [H69]: delete

2021). Increased biomass in irrigated plots is an indication that water is essential for root and shoot development below ground, which is essential for improving feed quantity and quality (Bhattarai et al., 2020). The study carried out in Punjab, Pakistan, investigated the growth of Italian ryegrass in different water conditions.

Root Length (cm)

The investigation of root length in Italian ryegrass (*Lolium multiflorum*) in Punjab, Pakistan, under rainfed and irrigated conditions, revealed significant variations. The average root length in the supplying water plots was found to be 5.95 cm (Table 2). This number, in addition to the low standard deviation of 0.4203, indicates the growth trend across the four plots under evaluation was mostly uniform and steady. This measure's stability is further supported by the coefficient of variation (CV), which has a low value of 7.06%. The lowest degree of variability implied by the low CV supports the validity of the mean as a representative indicator. The mean root length of 3.95 cm and the slightly greater standard deviation of 0.4203 were observed in the rainfed plots, in contrast. The computed coefficient of variation (CV) of 10.63% indicates a little higher level of variability in comparison to the irrigated situations (Table 3). The statistical significance of the observed difference in mean root length between irrigated and rainfed treatments is shown by a p-value below the specified significance level, such as 0.05. This lends credibility to the hypothesis that irrigation significantly affects Italian Ryegrass root length. The findings demonstrate the consistency and dependability of these measurements in addition to the quantitative difference in root length between the treatments. Experiments that were rainfed had shorter roots (3.950 cm) and more variability in root length than irrigated plots, which had an average of 5.950 cm. This shows that the availability of water has an evident impact on root length (Malcolm et al., 2015).

Comment [H70]: root length also demonstrated notable differences between the two conditions

Root Shoot Ratio

Italian ryegrass (*Lolium multiflorum*) development under various conditions in Punjab, Pakistan, was studied in relation to the Root Shoot Ratio, a crucial measure of the distribution of resources in plants. This ratio demonstrates how the biomass of the plant system divides between above-ground (shoot) and below-ground (root). The average Root Shoot Ratio under irrigation was found to be 0.11 (Table 2), with a standard deviation (SD) of 0.001 (Table 2). This suggests that resources were divided fairly evenly between roots and shoots throughout the plots. On the other hand, the mean Root Shoot Ratio in rainfed conditions was 0.1487, with a standard deviation of 0.001, showing a larger variability. It also means that, in the absence of irrigation, there may be more variation in the distribution of resources between roots and shoots, which indicates the plant's reaction to water stress. The relative variability of the Root Shoot Ratio within each treatment is indicated by the Coefficient of Variation, which is computed as the ratio of SD to mean. In comparison to the rainfed plots, where the CV was greater at 12.20%, the irrigated plots had a significantly lower CV at 9.26% (Table 3), suggesting a more uniform distribution of resources. The Root Shoot Ratio of the irrigated and rainfed conditions differed

significantly ($P < 0.05$) according to statistical analysis using the relevant procedures (e.g., t-test or ANOVA). This importance highlights that water availability affects the way that roots and shoots are distributed in Italian ryegrass. The Root Shoot Ratio also showed a substantial difference between rainfed and irrigated conditions, highlighting the impact of water on the distribution of resources within the plant (Junchao., et al., 2023) These results highlight that water influences the growth of Italian ryegrass roots and the distribution of resources (Malcolm et al., 2015; Junchao., et al., 2023).

Comment [H71]: of

Comment [H72]: distribution

Table 2. Effects of irrigated and rainfed ecology on growth parameters of Italian rye grass (*Lolium multiflorum*):

Treatment	Plant Height (cm)	Plant Population (plants/m ²)	Root Biomass Weight (g/plant)	Shoot Biomass Weight (g/plant)	Root Length (cm)	Root Shoot Ratio
Irrigated	43.75 ± 1.708 (a)	317.50 ± 17.079 (a)	0.8425 ± 0.0590 (a)	7.425 ± 0.590 (a)	5.950 ± 0.420 (a)	0.1135 ± 0.0010 (a)
Rainfed	13.75 ± 1.708 (b)	240.00 ± 21.60 (b)	0.4425 ± 0.0590 (b)	2.975 ± 0.386 (b)	3.950 ± 0.420 (b)	0.1487 ± 0.0018(b)

Table 3. Performance metrics and coefficient of variation for Italian rye grass under different conditions:

Treatment	Plant Height (cm)	Plant Population (plants/m ²)	Root Biomass Weight (g/plant)	Shoot Biomass Weight (g/plant)	Root Length (cm)	Root Shoot Ratio
Irrigated	3.90%	5.38%	7.01%	7.96%	7.06%	0.93%
Rainfed	12.43%	9.00%	13.36%	12.96%	10.63%	1.22%
Total	56.03%	16.19%	34.32%	46.58%	23.00%	14.36%

Table 4 : ANOVA

	Sum of Squares	df	Mean Square	F	Sig.
--	----------------	----	-------------	---	------

Plant Height (cm)	Between * Groups	(Combined)	1800.000	1	1800.000	617.143	.000
Treatment	Within Groups		17.500	6	2.917		
	Total		1817.500	7			
Plant Population (plants/m ²)	Between * Groups	(Combined)	12012.500	1	12012.500	31.681	.001
Treatment	Within Groups		2275.000	6	379.167		
	Total		14287.500	7			
Root Biomass Weight (g/plant)	Between * Groups	(Combined)	.320	1	.320	91.647	.000
Treatment	Within Groups		.021	6	.003		
	Total		.341	7			
Shoot Biomass Weight (g/plant)	Between * Groups	(Combined)	39.605	1	39.605	158.950	.000
Treatment	Within Groups		1.495	6	.249		
	Total		41.100	7			
Root Length (cm)	Between * Groups	(Combined)	8.000	1	8.000	45.283	.001
Treatment	Within Groups		1.060	6	.177		
	Total		9.060	7			
Root Shoot Ratio	Between * Groups	(Combined)	.002	1	.002	1122.375	.000
Treatment	Within Groups		.000	6	.000		
	Total		.002	7			

CONCLUSION

In conclusion, there were significant differences in the growth parameters of Italian Ryegrass (*Lolium multiflorum*) grown in Punjab, Pakistan, under rainfed and irrigated conditions. The grass indicated increased plant height, densities, biomass in the roots and shoots, and more even distribution of resources when it was irrigated. Rainfed conditions, on the other hand, resulted in altered distribution of resources, greater variability, and decreased growth. These results highlight the important influence that water availability has on the productivity and adaptation of Italian Ryegrass. It becomes clear that efficient water management is essential for maximizing the yields of fodder crops in areas like Punjab. For the region to support sustainable agriculture and animal feed, it is critical to understand the role that irrigation contributes in ensuring constant development and biomass production. This will assist in ensuring a steady and considerable availability of fodder.

Comment [H73]: This study aimed to examine the potential of Italian Ryegrass (*Lolium multiflorum*) as a forage crop in Punjab, Pakistan, focusing on its adaptability and growth performance in rainfed and irrigated conditions

Comment [H74]: The findings demonstrate that Italian Ryegrass exhibits significant improvements in growth parameters under irrigated conditions compared to rainfed conditions. The differences in plant height, plant population, root biomass, shoot biomass, and root length highlight the importance of adequate water supply for the successful cultivation of Italian Ryegrass in this region.

Comment [H75]: In irrigated conditions, Italian Ryegrass achieved greater biomass production and better establishment, suggesting that it is a viable forage crop option for areas with sufficient water availability. However, in rainfed conditions, the growth performance was notably lower, indicating the challenges faced by the crop under limited water availability. The consistent root-shoot ratio across both conditions suggests that while the overall biomass is influenced by water supply, the allocation of resources between roots and shoots remains relatively stable. The crop's ability to thrive under irrigated conditions presents an opportunity for enhancing forage availability and quality in areas with reliable water supply. However, for rainfed regions, further research and management strategies are needed to improve the performance of Italian Ryegrass under water-limited conditions. This study lays the groundwork for future investigations into optimizing the cultivation practices and expanding the use of Italian Ryegrass in diverse agro-ecological zones of Punjab.

REFERENCES

- Abbas, T., Nadeem, M. A., Tanveer, A., Matloob, A., Farooq, N., Burgos, N. A., & Chauhan, B. S. (2017). Confirmation of resistance in littleseed canarygrass (*Phalaris minor* retz) to accase inhibitors in central Punjab-Pakistan and alternative herbicides for its management. *Pak J Bot*, 49(4), 1501-1507.
- Akinroluyo, O. K., URBANAVIČIŪTĖ, I., JAŠKŪNĖ, K., KEMEŠYTĖ, V., & STATKEVIČIŪTĖ, G. (2019). Differences in salt tolerance between diploid and autotetraploid lines of *Lolium multiflorum* at the germination and vegetative stages. *Zemdirbyste-Agriculture*, 106(4).
- Bektas, H., Hohn, C. E., Lukaszewski, A. J., & Waines, J. G. (2023). On the Possible Trade-Off between Shoot and Root Biomass in Wheat. *Plants*, 12(13), 2513.
- Bhattarai, B., Singh, S., West, C. P., Ritchie, G. L., & Trostle, C. L. (2020). Water depletion pattern and water use efficiency of forage sorghum, pearl millet, and corn under water limiting condition. *Agric Water Manage*, 238, 106206.
- Bodner, G., Nakhforoosh, A., & Kaul, H. P. (2015). Management of crop water under drought: a review. *Agron Sustain Dev*, 35, 401-42.
- Byrne, N., Gilliland, T. J., Delaby, L., Cummins, D., & O'Donovan, M. (2018). Understanding factors associated with the grazing efficiency of perennial ryegrass varieties. *Ear J Agron*, 101, 101-108.
- Caiger-Watson, B. M. (2021). Alternative pasture species to perennial ryegrass/white clover suitable for use under irrigation in the Midlands of Tasmania (Doctoral dissertation, University Of Tasmania).
- Cavazzana, J. F., Paris, W., Molinete, M. L., Auache Filho, A. A., Denardin Costa, O. A., de Paula, A. L., ... et al. (2023). Beef steer performance on African Bermuda grass pasture overseeded with black oat and annual ryegrass: effects of added irrigation and temperate legumes. *Agronomy for Sustainable Development*, 43(1), 9.
- Goulart, R. Z., Reichert, J. M., & Rodrigues, M. F. (2020). Cropping poorly-drained lowland soils: Alternatives to rice monoculture, their challenges and management strategies. *Agricultural Systems*, 177, 102715.
- Ji, H. C., Whang, T. Y., Lee, K. W., Kim, W. H., Woo, J. H., Hong, K. H., & Choe, K. W. (2018). Growth characteristics and productivity of Italian ryegrass (*Lolium multiflorum* Lam.) new variety, 'Green Call'. *Journal of the Korean Society of Grassland and Forage Science*, 38(4), 247-252.

- Junchao, J., Chaodong, L., & Zhangyue, C. (2023). Soil Texture Rather Than Water Potential Determines the Root: Shoot Ratio in Ryegrass and Alfalfa. *J Soil Sci Plant Nutr*, 23(1), 1297-1305.
- Kumar, S., Singh, P., Devi, U., Yathish, K. R., Saujanya, P. L., Kumar, R., & Mahanta, S. K. (2023). An overview of the current fodder scenario and the potential for improving fodder productivity through genetic interventions in India. *Animal Nutrition and Feed Technology*, 23(3), 631-644.
- Malcolm, B. J., Moir, J. L., Cameron, K. C., Di, H. J., & Edwards, G. R. (2015). Influence of plant growth and root architecture of Italian ryegrass (*Lolium multiflorum*) and tall fescue (*Festuca arundinacea*) on N recovery during winter. *Grass Forage Sci*, 70(4), 600-610.
- Maxwell, T. M., McLenaghan, R. D., Edwards, G. R., Di, H. J., & Cameron, K. C. (2019). Italian ryegrass swards reduce N leaching via greater N uptake and lower drainage over perennial ryegrass cultivars varying in cool season growth rates. *New Zealand Journal of Agricultural Research*, 62(1), 69-82.
- Nadeem, A. A., Zha, Y., Shi, L., Zafar, Z., Ali, S., Zhang, Y., ... et al. (2023). SAFER-ET based assessment of irrigation patterns and impacts on groundwater use in the central Punjab, Pakistan. *Agric Water Manage*, 289, 108545.
- Nairizi, S. (2017). *Drought and Water Scarcity*. International Commission on Irrigation and Drainage: New Delhi, India.
- Pan, L., Zhang, X., Wang, J., Ma, X., Zhou, M., Huang, L., ... et al. (2016). Transcriptional profiles of drought-related genes in modulating metabolic processes and antioxidant defenses in *Lolium multiflorum*. *Front Plant Sci*, 7, 519.
- Pornaro, C., Serena, M., Macolino, S., & Leinauer, B. (2020). Drought stress response of turf-type perennial ryegrass genotypes in a Mediterranean environment. *Agronomy*, 10(11), 1810.
- Razzaq, A., Liu, H., Xiao, M., Mehmood, K., Shahzad, M. A., & Zhou, Y. (2023). Analyzing past and future trends in Pakistan's groundwater irrigation development: Implications for environmental sustainability and food security. *Environ Sci Pollut Res*, 30(12), 35413-35429.
- Rehman, A., Jingdong, L., Chandio, A. A., & Hussain, I. (2017). Livestock production and population census in Pakistan: Determining their relationship with agricultural GDP using econometric analysis. *Information Processing in Agriculture*, 4(2), 168-177.

- Rogers, M. E., Lawson, A. R., Chandra, S., & Kelly, K. B. (2020). Management options for improved survival of perennial ryegrass (*Lolium perenne* L.) under restricted irrigation during summer. *Crop Pasture Sci*, 101, 30.
- Saleh, F. F., & Al-Amery, A. H. (2023). Response of Ryegrass to Nitrogen Fertilizer under Two Irrigation Systems. *IOP Conf Ser Earth Environ Sci*, 1259(1), 012118.
- Scordia, D., & Cosentino, S. L. (2019). Perennial energy grasses: Resilient crops in a changing European agriculture. *Agriculture*, 9(8), 169.
- SEYDOŞOĞLU, S., & KÖKTEN, K. (2021). Nitrogen Requirement of Italian Ryegrass (*Lolium multiflorum*): A Review. *Turkish Journal of Range and Forage Science*, 2(1), 26-30.
- Sheikh, A., Hassan, J., Ijaz, S. S., Zaman, A., Alam, T., Ali, S., ... et al. (2023). Exploring Optimum Management Practices in Rainfed Areas to Reduce Soil erosion and Nutrient Losses. *Malays J Soil Sci*, 27, 164-178.
- Smit, H. P., Reinsch, T., Kluß, C., Loges, R., & Taube, F. (2021). Very Low Nitrogen Leaching in Grazed Ley-Arable-Systems in Northwest Europe. *Agronomy*, 11(11), 2155.
- Sustek-Sánchez, F., Rognli, O. A., Rostoks, N., Sömera, M., Jaškūnē, K., Kovi, M. R., ... & Sarmiento, C. (2023). Improving abiotic stress tolerance of forage grasses—prospects of using genome editing. *Frontiers in Plant Science*, 14, 1127532.
- Valdés-Rodríguez, O. A., Giadrossich, F., Pérez-Vázquez, A., & Moreno-Seceña, J. C. (2018). Above-and below-ground biomass and allometry of *Moringa oleifera* and *Ricinus communis* grown in a compacted clayey soil. *Flora*, 241, 35-45.
- Villela, A. L., Martinelli, R., Zenatti, T. F., Rufino-Jr, L. R., Monquero, P. A., Marluci daConceicao, P., ... & de Azevedo, F. A. (2021). Potential of two cover crops, signal grass and ruzi grass: suggested allelopathic effect on some important weeds. *Aust J Crop Sci*, 15(2), 260-270.
- Xie, F., Datta, R., & Qin, D. (2020). Plant growth and morphophysiological modifications in perennial ryegrass under environmental stress. In *Abiotic stress in plants* (p. 353). IntechOpen.
- Xie, Y., Liu, X., Ameer, M., Yu, H., Huang, Y., Li, X., ... et al. (2021). Evaluation of salt tolerance in Italian ryegrass at different developmental stages. *Agronomy*, 11(8), 1487.

UNDER PEER REVIEW

