

# Exploring the impact of *Spirulina platensis* supplementation in giant gourami (*Osphronemus goramy*) diet: on growth performance, hematological parameters, serum biochemical parameters, digestive enzyme activities, carcass composition and antioxidant ~~défense~~ defence mechanisms in giant gourami (*osphronemus goramy*)

## ABSTRACT

A 90-day trial investigated the effects of dietary *Spirulina platensis* (SP) supplementation on growth performance, haematology, serum biochemistry, digestive enzymes, and hepatic antioxidants in giant gourami (*Osphronemus goramy*). Fish ( $6.96 \pm 0.011$  g) were distributed into six treatments with 3 replicates each (15 fish per replicate) across 18, 60 L aquaria. Six isonitrogenous diets with 0% (control), 1% (SP1), 2% (SP2), 3% (SP3), 4% (SP4), and 5% (SP5) SP were provided *ad libitum* twice daily. Results showed dose-dependent improvements in final weight, weight gain, weight gain rate, protein efficiency ratio, and specific growth rate, with the highest values at 5% SP ( $p < 0.05$ ). Feed conversion ratio (FCR) decreased with higher SP, indicating enhanced efficiency. Body indices increased significantly with higher SP ( $p < 0.05$ ). Haematological parameters, including WBC (white blood cells), HB, HCT, monocyte %, and neutrophil %, improved with increasing SP significantly ( $p < 0.05$ ). Serum biochemistry showed markedly reduced cholesterol and triglycerides concentration, and increased total protein, albumin, and globulin levels with higher SP ( $p < 0.05$ ). Intestinal amylase and lipase activities increased significantly with SP ( $p < 0.05$ ), while protease remained consistent. Hepatic antioxidants revealed decreased MDA and increased SOD with higher SP ( $p < 0.05$ ). Carcass analysis showed increased protein, fibre, and fat content, and decreased ash and carbohydrates levels with a higher SP. These findings suggest that 5% SP enhances growth, biochemical parameters, haematology, digestive, and antioxidant mechanisms in *O. goramy*, supporting its potential as a dietary supplement in aquaculture.

Keywords: *Spirulina*, Feed Additive, Giant Gourami, Growth, Haematology, Immune Response, Digestive Enzyme

## 1. INTRODUCTION

In this modern era, the decline in the production of capture fisheries contrasts with the escalating demand for fish products for human consumption. Consequently, the Food and Agriculture Organization anticipates that aquaculture activities will contribute 60% of future fisheries production, a trend projected to persist and increase (FAO, 2018). The expansion of aquaculture to include a diverse range of fish species has amplified the requirement for commercial feed (2-5)2. Approximately 70% of animal production in aquaculture relies on commercial feeds rich in protein (2,6)2.

The giant gourami, *Osphronemus goramy*, stands out as a favoured species for both pisciculture (7) and the ornamental industry, primarily owing to its impressive size, reaching nearly 2 feet in length, its adaptability to freshwater as well as brackish water, and its suitability for live fish trade (8). This fish exhibits omnivorous behaviour with a strong inclination towards vegetarianism, demonstrating relative robustness and adaptability to adverse environmental conditions such as aerial respiration, which renders it highly appealing for low-input aquaculture (9). While the production of giant gourami fry relies on the natural spawning of captive broodfish held in ponds, efforts to standardize commercial-scale propagation of the species are still underway (10-12). It is noteworthy that the growth rate of gourami is relatively slow, requiring 18 months to attain a weight of 500 grams (13). Native to Southeast Asia, particularly from Indonesia, the Malay Peninsula, Thailand, and the

Comment [P1]: In italic form (*ad libitum*)

Comment [P2]: When you have inserted  $p < 0.05$  at the end of the sentence, it is not necessary to mention it in the sentence.

Comment [P3]: It is necessary to add the whole parts in replacement of Acronyms at the first time.

Comment [P4]: The same comment as the previous one (comment 3).

Comment [P5]: The same comment as the previous one (comment 3).

Formatted: Font: Italic

Comment [P6]: The format of inserted references should be similar.

Comment [P7]: It is not clear!

Mekong basin, the giant gourami *Osphronemus goramy* Lacépède, 1801, has been introduced to numerous other countries including India, Pakistan, Sri Lanka, Mauritius, Seychelles, Madagascar, Uganda, Philippines, Australia, New Zealand, Papua New Guinea, Japan, Hawaii, Colombia, France, Italy, among others, for aquaculture purposes (14,15). In India, the giant gourami was first introduced from Mauritius and Java (Indonesia) to Chennai during 1865–66 and subsequently distributed to various regions of the country (8). The cultivation of this species for food is particularly well-established in the state of Kerala, situated on the south-west coast. Furthermore, the National Fisheries Development Board, Government of India, is actively promoting this fish as one of the potential candidate species for both aquaculture and the ornamental fish trade in India (NFDB, 2016).

Comment [P8]: Refer to Comment 6!

*Spirulina platensis* is a type of blue-green algae characterized by its filamentous, spiral shape and remarkable nutritional value. Its cell wall possesses a soft texture due to a coating of complex sugars and proteins, facilitating easy digestion. This alga boasts an impressive nutrient composition, with approximately 62% protein content, as well as vitamins and antioxidants such as  $\beta$ -carotene and xanthophyll (17). (18) highlighted the presence of Gamma Linolenic Acid (GLA) and iron in *S. platensis*, alongside chlorophyll, glycogen, minerals (19), and phycocyanin and porphirin (20). Furthermore, it is rich in essential amino acids, vitamins B12 and  $\beta$ -carotene, phosphorus, iron, and calcium (21). *Spirulina platensis* is also known for its bioactive compounds with antioxidant properties, making it a valuable supplement for enhancing food (22) and serving as a significant source of micronutrients due to its high protein content (23).

*Spirulina*, renowned for its phycocyanin content (24), particularly C-phycocyanin, demonstrates antioxidant properties and may serve as a viable alternative to heparin due to its anticoagulant effects (25). Phycocyanin has been utilized for its antioxidative, antifungal and antitumor characteristics (26,27), as well as its immunostimulatory and hepatoprotective benefits (28). Multiple studies on Nile tilapia have indicated that incorporating spirulina into their diet significantly enhances growth rates, immune response, and disease resistance (29–32). Li et al. (32) have demonstrated that phycocyanin augments the activities of SOD and GPx enzymes in pancreatic beta cells. The inclusion of *Spirulina platensis* in diets can modulate the immune system (33) facilitate large-scale fish farming (34), and serve as a pivotal component for animal feed (35). Moreover, supplementing fish feed with *S. platensis* has been shown to enhance growth and survival rates (36,37) and can serve as a primary ingredient for feed production (38).

Comment [P9]: Why the references was inserted at the beginning of the sentences?!

The rise in giant gourami production can be attributed to advancements in production inputs such as feed, seed, disease management, and optimal culture techniques within giant gourami aquaculture. However, a significant challenge in culturing this fish lies in the inefficiency of production systems, largely due to their slow growth rate (39,40). In terms of feed technology development, numerous studies have been undertaken, particularly focusing on identifying alternative protein sources capable of promoting optimal growth performance in giant gourami (41,42). The utilization of chemicals for controlling bacterial or parasitic populations presents drawbacks such as bioaccumulation, pollution, and potential human health risks. Furthermore, the excessive reliance on antibiotics in aquaculture has led to issues of resistance, immunosuppression, and disruption of beneficial bacterial populations.

The utilization of *Spirulina*, known for its nutritional value and bioactive properties, offers a promising avenue for enhancing fish growth performance, immune response, and overall health, thereby contributing to the advancement of sustainable aquaculture practices. By investigating the effects of *Spirulina platensis* supplementation on the growth and health parameters of *Osphronemus goramy*, a commercially important fish species, this study aims to identify potential strategies for improving aquaculture efficiency and reducing reliance on conventional feed additives and antibiotics.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Diets

Six different diets were prepared with varying levels of *Spirulina platensis* (SP) ranging from 0% (C), 1% (SP1), 2% (SP2), 3% (SP3), 4% (SP4) and 5% (SP5). Detailed information regarding the formulation of the feeds and their proximate compositions presented in Table 1. Each diet was carefully prepared by weighing the ingredients according to the specified proportions and thoroughly mixing them for a period of 10 minutes. Subsequently, fish oil and plant oil were incorporated into the mixture, which was then passed through a 425  $\mu$ m sieve. Further mixing was carried out for

components containing fish oil for an additional 10 minutes. Afterward, distilled water was added and stirred for 5 minutes before being cooked in an autoclave. Pellets of 2.0 mm and 2.5 mm diameter were produced using a pelletizer through wet-extrusion after autoclave, followed by air-drying. The resulting pellets were then packaged in plastic bags and stored at room temperature.

**Table 1. Feed ingredients and the proximate chemical composition (% of dry weight) of the experimental diets.**

| Treatment                          | T0       | T1       | T2      | T3       | T4       | T5       |
|------------------------------------|----------|----------|---------|----------|----------|----------|
| <i>Spirulina p.</i>                | 0        | 1        | 2       | 3        | 4        | 5        |
| Soybean meal                       | 21       | 21       | 21      | 21       | 21       | 21       |
| Fish meal                          | 10.5     | 10.5     | 10.5    | 10.5     | 10.5     | 10.5     |
| Poultry by-product meal            | 16.5     | 16.5     | 16.5    | 16.5     | 16.5     | 16.5     |
| Rice bran                          | 21.3     | 20.3     | 19.3    | 19.3     | 18.3     | 18.3     |
| Wheat flour                        | 17.7     | 17.7     | 17.7    | 17.7     | 17.7     | 17.7     |
| Tapioca powder                     | 5        | 5        | 5       | 4        | 4        | 3        |
| Plant oil                          | 2        | 2        | 2       | 2        | 2        | 2        |
| Fish oil                           | 3        | 3        | 3       | 3        | 3        | 3        |
| Vitamin mineral mix <sup>1</sup>   | 3        | 3        | 3       | 3        | 3        | 3        |
| Total                              | 100      | 100      | 100     | 100      | 100      | 100      |
| Moisture                           | 8.74     | 8.84     | 8.86    | 8.82     | 8.76     | 8.88     |
| Dry matter                         | 91.26    | 91.16    | 91.14   | 91.18    | 91.24    | 91.12    |
| Protein                            | 30.399   | 30.289   | 30.179  | 30.179   | 30.069   | 30.069   |
| Ether extraction                   | 5.24     | 5.33     | 5.27    | 5.38     | 5.28     | 5.3      |
| Ash                                | 6.34     | 6.25     | 6.25    | 6.38     | 6.38     | 6.4      |
| Fiber                              | 4.2      | 4.26     | 4.24    | 4.23     | 4.36     | 4.3      |
| CHO                                | 43.85065 | 43.66796 | 43.7554 | 43.29211 | 43.28394 | 43.00126 |
| NfeNFE** (%) <sup>2</sup>          | 44.98    | 44.93    | 44.83   | 44.74    | 44.52    | 44.46    |
| Gross energy (kkal/g) <sup>3</sup> | 405.26   | 404.8    | 403.95  | 403.08   | 402.61   | 401.58   |

<sup>1</sup>Composition of vitamin mineral mixture of 1 kg: Vitamin A - 50,00,000 IU; Vitamin D3 - 10,00,000 IU; Vitamin B2 - 2.0 g; Vitamin E - 750 units; Vitamin K - 1.0 g; Calcium pantothenate 2.5 g; Nicotinamide - 10.0 g; Vitamin B12 - 6.0 g; Choline Chloride - 150.0 g; Calcium - 750.0 g; Manganese - 27.5 g; Iodine - 1.0 g; Ion - 7.5 g; Zinc - 15.0 g; Copper - 2.0 g; Cobalt - 0.45 g.

<sup>2</sup>NFE: Nitrogen free extract calculated using the following equation: NFE = 100 - (crude protein + ether extract + crude fiber + ash).

<sup>3</sup>GE: Gross energy calculated on the basis of 23.6, 39.4 and 17.2 k joule gross energy g-1 protein, ether extract and NFE, respectively.

P/E ratio: Protein energy ratio (mg crude protein kJ-1 gross energy) = CP/GE × 1000.

## 2.2 Experimental fish

A collection of 270 superficially robust giant gourami (*Osphronemus goramy*), averaging 6.96 ± 0.011g in weight, was procured from a seed farm. These fish were subjected to a one-week acclimatization period in laboratory conditions before being randomly distributed into six dietary groups. Each group of juvenile gouramis was accommodated in aquariums 60L, with a stocking density of 15 fish per aquarium. To maintain optimal conditions, each aquarium was equipped with a heater set to maintain a temperature of 30°C and aeration equipment. Before introducing the juveniles, the aquariums underwent sterilization using 30 ppm chlorine. Throughout the one-week acclimatization period, the juveniles were provided with a commercial diet containing 30% protein ad libitum. Body weight measurements were taken individually at the commencement and conclusion of the maintenance period for each aquarium. To sustain water quality, 25% of the total aquarium water volume was replaced before the morning feeding. Over the 90-days experimental period, fish were weighed every 15 days to adjust the daily feed ration, following the recommendation of the National Research Council (43), which was administered twice daily at 9:00 am and 5:00 pm.

## 2.3 Sample collection

At the end of 90-day experiment period, all fish underwent a 24-hour fasting period before being subjected to the final sampling. Fish samples were randomly collected from each treatment group and anesthetized using Tricainemethanesulfonate. Blood samples were then obtained from the caudal blood vessel using a 1 ml syringe (MDI Europa GmbH, Germany). After blood collection, the blood samples were immediately divided into two equal parts. One part was transferred to a tube containing an anti-coagulant (heparin) for hematological analysis, while the other part was transferred to non-heparinized tubes for biochemical and other analyses. The samples were then centrifuged at 4°C with 3000 rpm for 10 minutes and stored at -80 °C until further use, following the protocol

outlined by Adel et al(44). Additionally, carcass composition analysis was conducted on three fish samples, which were stored at  $-20^{\circ}\text{C}$ .

## 2.4 Growth performance

The daily feed intake (FI) of the fish was monitored consistently throughout the entire experiment. Before being weighed, the fish were carefully dried using fresh filter paper to eliminate any excess moisture. They were then weighed using a digital balance. The parameters were computed using the following formulas:

- 1) Weight gain rate (WGR, %) =  $100 \times (\text{Final body weight} - \text{initial body weight}) / \text{Initial body weight}$  (45)
- 2) WG (g fish-1) = Final body weight - Initial weight of the fish (g) (45)
- 3) Specific growth rate (SGR, % day<sup>-1</sup>) =  $100 \times [(\ln \text{ final individual weight} - \ln \text{ initial individual weight}) / \text{number of days}]$
- 4) ADG (g fish-1 day-1) = (Final body weight - Initial weight of the fish (g)) / Day
- 5) Feed conversion ratio (FCR) = Total diet fed / total wet weight gain (45)
- 6) Feed efficiency ratio (FER) =  $1 / \text{FCR}$
- 7) Survival rate (SR, %) =  $100 \times (\text{Final fish number} / \text{Initial fish number})$ .
- 8) Protein efficiency ratio (PER) =  $(\text{Final biomass (g)} - \text{initial biomass (g)}) / \text{Protein intake (46)}$

Immediately, three fish from each tank were dissected to determine the Viscerosomatic index (GSI%), Hepatosomatic index (HSI%), Visceral fat-somatic indexes (VFSI%), and Bilesomatic index (BSI) utilizing the prescribed formulas:

GSI =  $100 [\text{viscera weight (g)} / \text{whole body weight (g)}]$

HSI =  $100 [\text{liver weight (g)} / \text{whole body weight (g)}]$

VFSI =  $100 [\text{visceral fat weight (g)} / \text{whole body weight (g)}]$

BSI =  $100 [\text{Bile weight (g)} / \text{weight of liver (g)}]$

## 2.5 Proximate analysis

The fish body samples collected before infection, comprising three fish from each replicate (totaling nine from each group), underwent chemical analysis for moisture, protein, fat, and ash using the standard method recommended by the Association of Official Analytical Chemists (47). Approximately five fish per treatment were included in the analysis. Dry matter (DM) content was determined by drying the samples in an oven at  $105^{\circ}\text{C}$  for 24 hours. Crude protein was quantified using the micro-Kjeldhal method, with nitrogen content multiplied by 6.25 (utilizing a Kjeltchautoanalyzer, Model VELP Scientifica, UDK 127). Crude lipids were assessed using the Soxhlet method with ether extraction (Model VELP Scientifica, SER 148), while ash content was determined by burning samples at  $550^{\circ}\text{C}$  for 16 hours. Crude fiber (CF) analysis was conducted after digestion with 5% sulfuric acid and 5% sodium hydroxide for 15 minutes; the residues were subsequently dried and ashed. The nitrogen-free extract (NFE) was calculated using the formula:  $\text{NFE} = 100 - (\text{CP} + \text{EE} + \text{CF} + \text{ash})$ .

## 2.6 Biochemical Measurements

Superoxide dismutase (SOD Umg/prot) and malondialdehyde (MDA  $\mu\text{g/g}$ ) levels were determined following the methods outlined by Nishikimi et al(48) and Mihara & Uchiyama (49), respectively. The concentration of total protein (g dL<sup>-1</sup>) in plasma samples from the various experimental groups was measured using the Biuret method as described by AG(50). Albumin concentration (g dL<sup>-1</sup>) was assessed using the bromocresol green method (51), while globulin concentration (g dL<sup>-1</sup>) was computed by subtracting albumin from total protein. The Albumin to Globulin ratio (A/G ratio) was obtained by dividing albumin by globulin. Total cholesterol (TC) and triglycerides (TG) levels were determined using colorimetric diagnostic kits provided by spectrum-bioscience (Egyptian Company for Biotechnology, Cairo, Egypt), following the protocols of Allain et al(52) and McGowan et al. (53), respectively.

## 2.7 Hematological parameters analysis

Red blood cell (RBC,  $10^6 \text{ mm}^3$ ) and white blood cell (WBC,  $10^3 \text{ mm}^3$ ) counts, hemoglobin (Hb, g dL<sup>-1</sup>), and hematocrit (Ht %) were measured according to the method detailed by Li et al. (54).






Fig 1. Body weight and weight gain of *Osphronemus goramy* during the 90-day experiment





Fig12: FCR, FER, PER and DGR of *Osphronemus goramy* during the 90-day experiment



**Fig 23:** Specific growth rate (SGR) of *Osphronemus goramy* during the 90-day experiment

### 3.2. Body indices

The body indices of fish fed with varying spirulina concentration diets are listed in Table 3. The hepatosomatic index (HSI) exhibited a significant increase from T0 (0% SP) to T5 (5% SP) over the experimental period ( $p < 0.05$ ). Similarly, the viscerosomatic index (GSI%) showed a significant upward trend from T0 (0% SP) to T5 (5% SP), indicating an increase in visceral mass relative to body weight ( $p < 0.05$ ). The visceral fat-somatic indexes (VFSI%) displayed a significant rise from T0 to T5 (5% SP), signifying an augmentation in visceral fat deposition over time ( $p < 0.05$ ). The bile somatic index (BSI%) demonstrated a statistically significant increase from T0 to T5 (5% SP), indicating an elevation in bile accumulation relative to body weight ( $p < 0.05$ ).

**Table 3.** Mean (SE) value condition factor and body indices of *Osphronemus goramy* during the 90-day experiment

|                                      | T0 (0% SP)               | T1 (1% SP)               | T2 (2% SP)                | T3 (3% SP)                | T4 (4% SP)                | T5 (5% SP)                |
|--------------------------------------|--------------------------|--------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| Hepatosomatic index (HSI) (%)        | 0.926±0.012 <sup>c</sup> | 0.956±0.008 <sup>d</sup> | 0.976±0.008 <sup>d</sup>  | 0.993±0.008 <sup>c</sup>  | 1.1±0.011 <sup>b</sup>    | 1.29±0.005 <sup>a</sup>   |
| Viscerosomatic index (GSI%)          | 2.706±0.029 <sup>f</sup> | 2.78±0.017 <sup>e</sup>  | 2.853±0.014 <sup>d</sup>  | 2.983±0.014 <sup>c</sup>  | 3.216±0.02 <sup>b</sup>   | 3.493±0.02 <sup>a</sup>   |
| Visceral fat-somatic indexes (VFSI%) | 2.09±0.005 <sup>f</sup>  | 2.133±0.008 <sup>e</sup> | 2.17±0.005 <sup>d</sup>   | 2.21±0.005 <sup>c</sup>   | 2.29±0.005 <sup>b</sup>   | 2.41±0.005 <sup>a</sup>   |
| Bilesomatic (BSI%)                   | 9.916±0.014 <sup>f</sup> | 10±0.025 <sup>e</sup>    | 10.206±0.008 <sup>d</sup> | 10.296±0.014 <sup>c</sup> | 10.343±0.008 <sup>b</sup> | 10.443±0.008 <sup>a</sup> |

### 3.3. Hematological parameters

As indicated in Table 4, no significant differences were observed in RBC levels among all treatments ( $P > 0.05$ ). However, WBC, Hb, HCT, Monocyte, and Neutrophil levels exhibited a significant increase from T0 to T5 (5% SP) as spirulina concentration increased ( $p < 0.05$ ). Lymphocyte levels demonstrated significant fluctuations across different concentrations of spirulina, with the highest level observed at T5 (5% SP) and the lowest in T0 ( $p < 0.05$ ).

**Table 4.** Effects of dietary *S. platensis* on hematological parameters in *Osphronemus goramy*.

|                                  | T0 (0% SP)                | T1 (1% SP)                | T2 (2% SP)                | T3 (3% SP)                | T4 (4% SP)                | T5 (5% SP)                |
|----------------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| RBC ( $\text{mm}^3 \cdot 10^6$ ) | 3.216±1.608 <sup>a</sup>  | 4.933±0.008 <sup>a</sup>  | 4.996±0.008 <sup>a</sup>  | 5.123±0.012 <sup>a</sup>  | 5.263±0.008 <sup>a</sup>  | 5.413±0.008 <sup>a</sup>  |
| WBC ( $\text{mm}^3 \cdot 10^4$ ) | 8.886±0.02 <sup>f</sup>   | 9.316±0.02 <sup>e</sup>   | 9.686±0.012 <sup>d</sup>  | 10.516±0.02 <sup>c</sup>  | 11.203±0.017 <sup>b</sup> | 12.303±0.024 <sup>a</sup> |
| (Hb, g dL <sup>-1</sup> )        | 5.526±0.012 <sup>c</sup>  | 5.67±0.011 <sup>d</sup>   | 5.793±0.008 <sup>c</sup>  | 5.766±0.014 <sup>c</sup>  | 5.906±0.008 <sup>b</sup>  | 6±0.011 <sup>a</sup>      |
| HCT (%)                          | 33.513±0.008 <sup>f</sup> | 33.813±0.008 <sup>e</sup> | 34.1±0.005 <sup>d</sup>   | 34.423±0.008 <sup>c</sup> | 34.846±0.008 <sup>b</sup> | 35.216±0.008 <sup>a</sup> |
| Monocyte (%)                     | 2.51±0.011 <sup>f</sup>   | 2.776±0.017 <sup>e</sup>  | 2.916±0.014 <sup>d</sup>  | 3.33±0.015 <sup>c</sup>   | 3.79±0.015 <sup>b</sup>   | 4.21±0.011 <sup>a</sup>   |
| Lymphocyte (%)                   | 76.233±0.12 <sup>a</sup>  | 73.066±0.12 <sup>d</sup>  | 75.166±0.088 <sup>b</sup> | 72.733±0.088 <sup>d</sup> | 74.333±0.12 <sup>c</sup>  | 70.733±0.12 <sup>e</sup>  |

|                |                           |                           |                           |                           |                          |                           |
|----------------|---------------------------|---------------------------|---------------------------|---------------------------|--------------------------|---------------------------|
| Neutrophil (%) | 10.116±0.012 <sup>a</sup> | 11.423±0.014 <sup>c</sup> | 12.663±0.008 <sup>d</sup> | 12.983±0.017 <sup>e</sup> | 13.65±0.015 <sup>b</sup> | 14.306±0.008 <sup>a</sup> |
|----------------|---------------------------|---------------------------|---------------------------|---------------------------|--------------------------|---------------------------|

### 3.4. Serum biochemical parameters

The serum biochemical parameters of fish fed with Spirulina are showed in Table 5. Fish fed the control diet exhibited the highest levels of cholesterol and triglyceride, with these levels decreasing as the dietary *S. platensis* levels increased ( $P<0.05$ ). Total protein (g dL<sup>-1</sup>), albumin (g/dL), and globulin (g/dL) levels increased with the increase in spirulina concentration. The significantly highest levels of protein (g dL<sup>-1</sup>), albumin (g/dL), and globulin (g/dL) were observed in the T5 (5% SP) diet group ( $P<0.05$ ). The A:G ratio (g/dL) was significantly highest in the T0 (0% SP) diet group and lowest in the T1 (1% SP) diet group ( $P<0.05$ ).

**Table5. The effects of dietary *Spirulina platensis* phycocyanin (SPC) on the blood biochemical parameters of *Osphronemus goramy*.**

|                                     | T0 (0% SP)                 | T1(1% SP)                | T2(2% SP)                 | T3(3% SP)                 | T4(4% SP)                  | T5(5% SP)                  |
|-------------------------------------|----------------------------|--------------------------|---------------------------|---------------------------|----------------------------|----------------------------|
| Total protein (g dL <sup>-1</sup> ) | 3.53±0.015 <sup>f</sup>    | 3.94±0.017 <sup>e</sup>  | 4.186±0.012 <sup>d</sup>  | 4.36±0.011 <sup>c</sup>   | 4.663±0.008 <sup>b</sup>   | 4.833±0.014 <sup>a</sup>   |
| Albumin (g/dL)                      | 1.843±0.012 <sup>f</sup>   | 1.91±0.011 <sup>e</sup>  | 1.97±0.005 <sup>d</sup>   | 2.093±0.008 <sup>c</sup>  | 2.153±0.008 <sup>b</sup>   | 2.27±0.011 <sup>a</sup>    |
| Globulin(g/dL)                      | 1.823±0.014 <sup>f</sup>   | 2.033±0.02 <sup>e</sup>  | 2.093±0.008 <sup>d</sup>  | 2.176±0.008 <sup>c</sup>  | 2.256±0.014 <sup>b</sup>   | 2.33±0.015 <sup>a</sup>    |
| A:G ratio (g/dL)                    | 1.011±0.003 <sup>a</sup>   | 0.939±0.015 <sup>c</sup> | 0.941±0.006 <sup>c</sup>  | 0.961±0.001 <sup>bc</sup> | 0.954±0.005 <sup>bc</sup>  | 0.974±0.007 <sup>c</sup>   |
| Triglyceride (Mg/dl)                | 315.366±0.384 <sup>d</sup> | 300.3±0.115 <sup>b</sup> | 293.633±0.12 <sup>c</sup> | 289.7±0.115 <sup>d</sup>  | 283.733±0.12 <sup>c</sup>  | 280.066±0.12 <sup>f</sup>  |
| Cholesterol (mg/dl)                 | 191.9±0.115 <sup>a</sup>   | 187.9±0.057 <sup>b</sup> | 181.1±0.115 <sup>c</sup>  | 178±0.057 <sup>d</sup>    | 172.933±0.088 <sup>e</sup> | 168.566±0.145 <sup>f</sup> |

### 3.5. Intestinal digestive enzymes and Hepatic antioxidant activities

Table 6 illustrates the digestive enzymes activities and antioxidant activities of fish fed with Spirulina. Both Amylase and Lipase activities experienced a noteworthy increase as the concentration of spirulina increased from 0% to 5% ( $p<0.05$ ). These enzymatic activities demonstrated a gradual elevation with the increasing spirulina concentration in the diet. However, there was no significant variation observed in Protease activity across different concentrations of spirulina ( $p>0.05$ ). MDA levels showed a significant decrease from T0 (0% SP) to T5 (5% SP) with the increasing concentration of spirulina ( $p<0.05$ ). Conversely, SOD levels exhibited a significant increase from T0 (0% SP) to T5 (5% SP) with the increasing spirulina concentration ( $p<0.05$ ).

**Table6. Intestinal digestive enzymes activities and Hepatic antioxidant activities of *Osphronemus goramy*.**

|                           | T0 (0% SP)               | T1(1% SP)                | T2(2% SP)                | T3(3% SP)                | T4(4% SP)                | T5(5% SP)                |
|---------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Amylase (U/mg substrate)  | 2.11±0.011 <sup>f</sup>  | 2.326±0.014 <sup>e</sup> | 2.25±0.011 <sup>d</sup>  | 2.456±0.008 <sup>c</sup> | 2.656±0.02 <sup>b</sup>  | 2.873±0.012 <sup>a</sup> |
| Lipase (U/mg substrate)   | 6.803±0.017 <sup>f</sup> | 7.346±0.014 <sup>e</sup> | 7.126±0.017 <sup>d</sup> | 7.833±0.02 <sup>c</sup>  | 7.553±0.017 <sup>b</sup> | 7.973±0.008 <sup>a</sup> |
| Protease (U/mg substrate) | 0.019±0 <sup>e</sup>     | 0.022±0 <sup>b</sup>     | 0.019±0 <sup>f</sup>     | 0.02±0 <sup>c</sup>      | 0.023±0 <sup>b</sup>     | 0.025±0 <sup>a</sup>     |
| MDA (µg/g)                | 2.02±0.017 <sup>a</sup>  | 1.96±0.011 <sup>b</sup>  | 1.906±0.008 <sup>c</sup> | 1.836±0.014 <sup>d</sup> | 1.496±0.017 <sup>e</sup> | 1.356±0.012 <sup>f</sup> |
| SOD (Umg/prot)            | 6.406±0.008 <sup>f</sup> | 7.25±0.011 <sup>e</sup>  | 7.73±0.011 <sup>d</sup>  | 8.03±0.011 <sup>c</sup>  | 8.36±0.011 <sup>b</sup>  | 8.576±0.008 <sup>a</sup> |

### 3.6. Carcas composition

Protein content exhibited a significant increase ( $p < 0.05$ ) as spirulina content increased, from 17.07% ± 0.03 at T0 to 18.6% ± 0.01 at T5 (Table 7). Fat content showed a similar trend, with significant differences observed between each treatment ( $p < 0.05$ ). Fiber content also increased significantly with higher SP content, ranging from 0.96% ± 0 at T0 to 1.2% ± 0.01 at T5. ASH (Ash content) decreased significantly ( $p<0.05$ ) with increasing SP content, with values ranging from 5.13% ± 0.01 at T0 to 4.23% ± 0.01 at T5. Carbohydrate (CHO) content displayed a significant decrease ( $p<0.05$ ) as SP content increased, with values ranging from 1.98% ± 0.01 at T0 to 1.21% ± 0 at T5. Dry matter content showed significant variability across SP levels, and Moisture content exhibited an inverse relationship with spirulina content. Overall, the results indicate that increasing SP content in the body composition led to significant changes in nutritional composition, particularly in protein, fat, fibre, ash, and carbohydrate contents.

**Table7. Body composition of *Osphronemus goramy*.**

|                  | T0 (0% SP)              | T1 (1% SP)              | T2 (2% SP)               | T3 (3% SP)              | T4 (4% SP)              | T5 (5% SP)              |
|------------------|-------------------------|-------------------------|--------------------------|-------------------------|-------------------------|-------------------------|
| Crude Protein%   | 17.07±0.03 <sup>a</sup> | 17.36±0.01 <sup>c</sup> | 17.51±0.01 <sup>d</sup>  | 17.79±0.01 <sup>c</sup> | 18.19±0 <sup>b</sup>    | 18.6±0.01 <sup>a</sup>  |
| Crude Fat        | 4.2±0.01 <sup>d</sup>   | 4.29±0.01 <sup>c</sup>  | 4.28±0.01 <sup>c</sup>   | 4.51±0.02 <sup>b</sup>  | 4.53±0.02 <sup>b</sup>  | 4.61±0.01 <sup>a</sup>  |
| Crude Fiber      | 0.96±0 <sup>d</sup>     | 0.98±0.01 <sup>d</sup>  | 1.04±0 <sup>c</sup>      | 1.08±0.01 <sup>b</sup>  | 1.11±0.01 <sup>b</sup>  | 1.2±0.01 <sup>a</sup>   |
| ASH              | 5.13±0.01 <sup>f</sup>  | 4.91±0.01 <sup>c</sup>  | 4.81±0.01 <sup>d</sup>   | 4.51±0.01 <sup>c</sup>  | 4.3±0.01 <sup>b</sup>   | 4.23±0.01 <sup>a</sup>  |
| CHO              | 1.98±0.01 <sup>a</sup>  | 1.96±0 <sup>a</sup>     | 1.95±0.01 <sup>a</sup>   | 1.97±0 <sup>a</sup>     | 1.52±0 <sup>b</sup>     | 1.21±0 <sup>c</sup>     |
| Dry matter       | 29.35±0.06 <sup>d</sup> | 29.52±0.02 <sup>c</sup> | 29.61±0.04 <sup>c</sup>  | 29.87±0.03 <sup>a</sup> | 29.66±0.02 <sup>b</sup> | 29.87±0.05 <sup>a</sup> |
| Moisture content | 70.64±0.06 <sup>a</sup> | 70.48±0.02 <sup>b</sup> | 70.39±0.04 <sup>bc</sup> | 70.12±0.03 <sup>d</sup> | 70.33±0.02 <sup>c</sup> | 70.13±0.05 <sup>d</sup> |

### 3.7. Water quality

The water quality of the concrete pond where giant gourami juveniles were reared was assessed. Parameters including water temperature, dissolved oxygen (DO), total alkalinity, hardness, pH, and nitrates were measured and found to be within the typical range recommended by WHO/FAO standards, as outlined in Table 8.

**Table8. Water quality parameter observed during 90 days of experiment**

|   | T0 (0% SP)               | T1 (1% SP)               | T2 (2% SP)               | T3 (3% SP)               | T4 (4% SP)               | T5 (5% SP)               |
|---|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
| Temperature (°C)                              | 28.56±0.088 <sup>c</sup> | 28.93±0.088 <sup>d</sup> | 28.6±0.057 <sup>c</sup>  | 30.1±0.057 <sup>b</sup>  | 30.7±0.057 <sup>a</sup>  | 29.3±0.057 <sup>c</sup>  |
| PH  | 7.833±0.008 <sup>a</sup> | 7.396±0.012 <sup>f</sup> | 7.496±0.008 <sup>c</sup> | 7.533±0.008 <sup>d</sup> | 7.716±0.008 <sup>c</sup> | 7.793±0.008 <sup>b</sup> |
| DO (mg/L)                                     | 5.813±0.012 <sup>c</sup> | 6.096±0.012 <sup>b</sup> | 6.24±0.011 <sup>a</sup>  | 5.963±0.012 <sup>c</sup> | 5.863±0.008 <sup>d</sup> | 6.266±0.012 <sup>a</sup> |
| Total Alkalinity (mg/L as CaCO <sub>3</sub> ) | 55.33±0.088 <sup>d</sup> | 56.46±0.088 <sup>c</sup> | 59.53±0.088 <sup>b</sup> | 54.73±0.088 <sup>e</sup> | 60.06±0.088 <sup>a</sup> | 59.33±0.088 <sup>b</sup> |
| Hardness (mg/L as CaCO <sub>3</sub> )         | 66.56±0.088 <sup>c</sup> | 68.56±0.088 <sup>d</sup> | 69.23±0.088 <sup>c</sup> | 65.56±0.088 <sup>f</sup> | 69.86±0.088 <sup>b</sup> | 70.23±0.088 <sup>a</sup> |
| NO <sub>3</sub> (mg/L)                        | 0.054±0.001 <sup>a</sup> | 0.031±0.001 <sup>c</sup> | 0.046±0.001 <sup>b</sup> | 0.035±0.001 <sup>c</sup> | 0.047±0.001 <sup>b</sup> | 0.057±0 <sup>a</sup>     |
| NO <sub>2</sub> (mg/L)                        | 0.027±0.002 <sup>c</sup> | 0.069±0.001 <sup>b</sup> | 0.021±0.001 <sup>f</sup> | 0.063±0.001 <sup>c</sup> | 0.041±0.001 <sup>d</sup> | 0.079±0 <sup>a</sup>     |
| TAN (mg/L)                                    | 0.606±0.008 <sup>c</sup> | 0.773±0.008 <sup>d</sup> | 0.48±0.005 <sup>f</sup>  | 1.506±0.008 <sup>b</sup> | 1.806±0.008 <sup>a</sup> | 0.896±0.008 <sup>c</sup> |

## 4. DISCUSSION

### 4.1. Growth performance

During feeding trial, we observed a significant increase in the growth performance of *Osphronemus goramy* when the Spirulina concentration in the feed increased from 1% to 5% over a span of 90 days. The inclusion of Spirulina in aquafeed led to an increase in hunger and notably elevated feed intake, resulting in improvements in parameters such as Final weight (FW), weight gain (WG), weight gain% (WG%), feed efficiency ratio (FER), average daily growth (ADG), and specific growth rate (SGR). These findings are consistent with previous research (29,36,59,60). Studies by Hassanpour et al. (61) and Zeinab et al. (62) found that the supplementation of Spirulina in the diets of rainbow trout and Nile Tilapia, respectively, resulted in higher body weights compared to control groups. Similarly, research by Jana et al. (Jana et al., 2014), Simanjuntak et al. (13), and Abu-Elala et al. (63) demonstrated significant growth enhancement in fish fed with Spirulina, without any adverse effects (64) and with potential as a feed material (38). Additionally, the inclusion of Spirulina in aquafeed improved survival, growth rate, and feed utilization in *Macrobrachium rosenbergii* prawn culture at concentration level from 5% to 20%, as observed in similar studies on *Litopenaeus vannamei* (65). Overall, Spirulina shows promise as a natural supplement for enhancing growth and health in aquaculture.

*Spirulina platensis* enhances growth in fish due to its highly digestible protein content, abundant essential fatty acids, vitamins (such as vitamin B12), and minerals (66,67). Its cellular structure lacks cellulose, making it easily digestible by fish digestive enzymes, aided by the presence of mucopolymers (64). Additionally, Spirulina promotes the growth of beneficial intestinal flora in fish and aids in the breakdown of indigestible feed components (36,64). Parameters like protein

Comment [P10]: Refer to comment 6.

efficiency ratio (PER), survival rate (SR%), and ~~feed conversion ratio (FCR)~~ significantly improve with increased Spirulina concentration.

#### 4.2. Body indices

At the end of experiment result was found that the body indices (HIS%, VFSI%, GSI% and BSI%) was increased with increase in concentration of spirulina in diet from (0% to 5%). Significant highest Hepatosomatic index (HSI) (%), Viscerosomatic index (GSI%), Visceral fat-somatic indexes (VFSI%) and Bilesomatic (BSI%) observed in T5 (5% SP) diet ( $p < 0.05$ ). A higher hepatosomatic index (HSI) reflects increased glycogen levels and favorable environmental conditions. This relationship was observed in separate studies conducted by Velasquez et al. (68) on Nile tilapia and Tulli et al. (69) on European sea bass juveniles (*Dicentrarchus labrax*), where diets containing 20% *Tetraselmis suecica* microalgae resulted in lower HSI compared to control diets without microalgae. This connection between the level of microalgae inclusion and improved lipid mobilization efficiency in fish liver has also been noted by (70) and (71). The viscerosomatic index (VSI) serves as an indicator of fish digestive capacity. Consistent findings were reported by (68) in Nile tilapia, where the highest VSI was observed in fish fed diets with 0% to 30% microalgae inclusion, showing a downward trend as the level of Arthrospira inclusion increased. Nevertheless, it's essential to recognize that VSI represents just one aspect of fish digestive capacity, as other factors likely play a role.

#### 4.3. Hematological parameters

In our study, we observed a dose-dependent increase in blood parameters, including, WBC, HCT, Hb, Monocyte (%) and Neutrophil (%), in fish fed with 5% *Spirulina platensis* compared to the control group. But there is no significant difference was found in the RBC count among fish fed different concentrations of Spirulina ( $p > 0.05$ ) and Lymphocyte (%), count is significantly decrease with increase in concentration of spirulina ( $p < 0.05$ ). the RBC count Similar findings were reported by (72) and (34). Promya and Chitmant(73) demonstrated higher WBC and RBC counts in African sharp tooth catfish fed diets containing 5% *A. platensis* + basal. (44) found increased RBC and Hb levels in Great Sturgeon fed with *S. platensis*. Spirulina's high iron content significantly affects erythropoiesis, as seen in studies on anaemic rats (74). (75) observed the highest levels of RBC, WBC, and Hb in Nile tilapia with 10% Spirulina supplementation. Similar increases in WBC were reported in large yellow croaker and carp with Spirulina supplementation(54,74). This finding was also supported by (76) and (77). This is attributed to polypeptide-phycoerythrin, enhancing WBC production (78). (62) noted increased hemoglobin levels in Nile tilapia fed Spirulina, while (36) found 10% *S. platensis* supplementation increased hematocrit values. However, carp RBCs were unaffected by 3.0–5.0 g/kg Spirulina (79).

#### 4.4. Blood biochemical parameters

In fish, the levels of total protein, albumin, and globulin in serum provide valuable diagnostic insights into overall nutritional status, vascular system integrity, and liver function (80). The current study observed that supplementation with 5% spirulina resulted in higher levels of total protein, albumin, and globulin compared to the control group. Conversely, previous research by (13) found that diets containing 2.0–6.0% Spirulina significantly increased serum levels of total protein, globulin, and the A/G ratio. Similar findings were reported by (81), where feeding with 10.0% Spirulina significantly elevated serum albumin levels, as also observed by (29). Additionally, (62) noted that increased levels of *S. platensis* correlated with higher serum total protein and globulin, although there was no significant difference in albumin levels ( $p > 0.05$ ). The rise in serum albumin levels could be attributed to Spirulina's various active compounds, including carotenoids, polysaccharides, vitamins, minerals, and linoleic acid, which act as immunostimulants, enhancing the immune system (82). Spirulina's potent antioxidant properties contribute to its ability to improve immune capacity and health status in *Osphronemus gourami*, similar to findings in *O. niloticus*(22,63). Furthermore, (83) demonstrated that different levels of *S. platensis* improved total serum protein, albumin, globulin, and biochemical responses in Mugilcephalus. Including *S. platensis* and *Sargassumilicifolium* extract in diets for Husohuso and *O. niloticus* also increased serum total protein and globulin levels (62,84). The serum albumin level plays a crucial role in regulating metabolism, maintaining osmotic pressure in the blood, and facilitating the transport of both exogenous and endogenous metabolites (85). Globulin serves as a prominent carrier/transporter in living organisms Simanjuntak et al.(13).

In the present result cholesterol and triglyceride level decrease with increase in spirulina concentration. Similarly, (86) discovered that incorporating 1.0 g/kg Spirulina into tilapia diets did not impact the serum levels of total cholesterol and triglycerides. However, diets containing 10.0% Spirulina significantly raised the serum levels of cholesterol and triglycerides(29). *Spirulina*

*platensis* contains  $\alpha$ -linolenic acid, linoleic acid, and  $\beta$ -carotene, which have the potential to lower cholesterol and triglyceride levels and prevent hypercholesterolemia and atherosclerosis (87,88).

#### 4.5. Digestive enzyme

Algae are abundant in both macro and micro elements, which have the potential to stimulate digestive enzyme activity (89). Low concentrations of microalgae in the gut may trigger the secretion of endogenous digestive enzymes and promote fish growth (90,91). (92) noted that *Isochrysis galbana* algae enhanced pancreatic and intestinal enzyme production in sea bass (*Dicentrarchus labrax*) larvae. While fish typically struggle to digest plant matter due to the cellulose cell wall, *S. platensis* is known to have a thin cell wall composed primarily of 80% pectin and 20% cellulose, facilitating improved digestion and assimilation (93). Consequently, *S. platensis* is easily digested and may enhance the assimilation of other nutrients, including antioxidants.

In the current experiment, the activity of amylase (U/mg substrate), lipase (U/mg substrate), and protease (U/mg substrate) significantly increased with the concentration of spirulina in the diet compared to the control group ( $p > 0.05$ ). Similarly, Biabani Asrami et al. (27) observed that dietary supplementation of phycocyanin extracted from Spirulina enhanced the activities of amylase, protease, and lipase in Guppy fish (*Poecilia reticulata*). Additionally, Teimouri et al. (94) found that *Spirulina platensis* enhanced digestive enzyme activity, thereby improving nutrient assimilation. Therefore, increasing the inclusion of spirulina in the diet improved the digestive enzyme activity in *Osphronemus gourami*.

#### 4.6. Antioxidant activity

Superoxide dismutase (SOD) play crucial roles in scavenging free radicals and protecting essential cellular macromolecules and organelles from oxidative damage (95). Previous studies have indicated that dietary supplementation of spirulina can reduce antioxidant activities in Nile tilapia (*Oreochromis niloticus*) (96) and goldfish (*Carassius auratus*) (97). Additionally, Li et al. (2009) demonstrated that phycocyanin enhances the activities of SOD enzymes in pancreatic beta cells. Tayag et al. (98) and Lin et al. (99) reported a direct increase in SOD enzyme activity with spirulina extract in white shrimp. Abdelkhalek et al. (100) illustrated that spirulina supplementation enhances SOD enzyme activity in *Oreochromis niloticus*. Furthermore, Yu et al. (75) observed that rainbow trout fed with 10% spirulina exhibited the highest SOD activity.

MDA serves as the end product of lipid peroxidation, leading to cellular toxicity and accelerating damage to cells and tissues (101–103). In the current study, MDA enzyme activity decreased with an increase in spirulina concentration, consistent with previous findings in Nile tilapia (*Oreochromis niloticus*) (96), and Carp (*Cyprinus carpio*) (104). Similarly, Abdelkhalek et al. (100) reported that dietary supplementation with *S. platensis* reduced MDA levels in the liver, kidney, and gills of Nile tilapia.

#### 4.7. Body composition

In this study, findings revealed that protein deposition in the body increased with higher spirulina content, notably peaking at  $18.6 \pm 0.01a$  in the T5 group (5% SP) ( $p > 0.05$ ). Likewise, lipid ( $4.61 \pm 0.01a$ ) and fibre ( $1.2 \pm 0.01a$ ) content rose with escalating spirulina concentration, while carbohydrate (CHO) ( $1.21 \pm 0.0c$ ) and ash (ASH) ( $4.23 \pm 0.01a$ ) levels declined in *Osphronemus goramy* bodies with increasing spirulina content. Overall, the body composition exhibited optimal results in the high 5% (spirulina) concentration diet, indicating efficient utilization of spirulina-infused feed by *Osphronemus goramy* bodies. These results are consistent with past research; Moreira et al. (105) proposed that spirulina LEB-18 is a valuable dietary addition for addressing nutritional deficiencies due to its rich mineral and nutritional content. Changes in body fat and protein content are linked to variations in synthesis and deposition levels within muscle tissues (29,106). (107) noted significant enhancements in the biochemical composition of *Macrobrachium rosenbergii* PL bodies when fed Spirulina-enriched Artemia nauplii. Feeding larvae of *Cyprinus carpio* with 20% Spirulina led to considerable improvements in body composition, particularly in protein composition and body fat (59).

#### 4.8. Water quality

In The present study found that water quality parameters, including temperature ( $^{\circ}\text{C}$ ), dissolved oxygen (mg/L), total alkalinity (mg/L as  $\text{CaCO}_3$ ), hardness (mg/L as  $\text{CaCO}_3$ ), Ph and nitrates (mg/L), were all within acceptable limits according to WHO/FAO standards (Table 9).

**Table9. standard water quality parameter range according to WHO/FAO limits**

| Water quality parameters                      | WHO/FAO limits | References |
|---|----------------|------------|
| Water temperatures (°C)                       | 25–33          | (108)      |
| Dissolved oxygen (mg/L)                       | 3-5            | (109)      |
| Total alkalinity (mg/L as CaCO <sub>3</sub> ) | 120            | (110)      |
| Hardness (mg/L as CaCO <sub>3</sub> )         | 168            | (110)      |
| pH  | 6.5–9.0        | (110)      |
| Nitrates (mg/L)                               | 0.2–219        | (110)      |

## 5. CONCLUSIONS

In conclusion, our experiment showed that *Osphronemus gourami* fed a diet supplemented with *Spirulina platensis* at 5% for 90 days improved fish growth performance (FW, WG, WG%, PER, FCR, FER, ADG, SGR, and SR%), increased feed and protein utilization, improve the body deposition (Protein, fat, ASH, CHO, fiber), elevated body indices (HIS%, VFSI%, GSI%, and BSI%), enhanced hematological parameters (RBC, WBC, HCT, Hb, Monocyte %, and Neutrophil %), improved biochemical responses (Total serum protein, Albumin, and Globulin), decreased levels of cholesterol and triglyceride, and increased digestive (Amylase, Lipase, and Protease) and antioxidant (SOD & MDA) enzyme activity in fish and improve the immunity.

## REFERENCES

1. FAO. (Food and Agriculture Organization). 2008 The state of world fisheries and aquaculture: Opportunities and challenges. Food and Agriculture Organization of the United Nations. 2018;209.
2. Hua K, Cobcroft JM, Cole A, Condon K, Jerry DR, Mangott A, et al. The future of aquatic protein: implications for protein sources in aquaculture diets. *One Earth*. 2019;1(3):316–29.
3. Gómez B, Munekata PE, Zhu Z, Barba FJ, Toldrá F, Putnik P, et al. Challenges and opportunities regarding the use of alternative protein sources: Aquaculture and insects. *Adv Food Nutr Res*. 2019;89:259–95.
4. Zhang Y, Lu R, Qin C, Nie G. Precision nutritional regulation and aquaculture. *Aquac Rep*. 2020;18:100496.
5. Nagappan S, Das P, AbdulQuadir M, Thaher M, Khan S, Mahata C, et al. Potential of microalgae as a sustainable feed ingredient for aquaculture. *J Biotechnol*. 2021;341:1–20.
6. Dawood MA, Koshio S. Application of fermentation strategy in aquafeed for sustainable aquaculture. *Rev Aquac*. 2020;12(2):987–1002.
7. Azrita HS, Syandri H. Morphological character among five strains of giant gourami, *Osphronemus gourami* Lacepede, 1801 (Actinopterygii: Perciformes: Osphronemidae) using a truss morphometric system. *Int J Fish Aquat Stud* [Internet]. 2015;2(6):344–50. Available from: <https://www.fisheriesjournal.com/vol2issue6/Pdf/2-6-58.1.pdf>
8. Sundara RB. Notes on the freshwater fishes of Madras. *Records Indian Museum* 7; 1916.
9. Caruso D, Arifin ZO, Subagja J, Slembrouck J, New M. *Osphronemus goramy*. Cultured Aquatic Species Information Programmed. Rome: FAO Fisheries and Aquaculture Department; 2019.

10. Amornsakun T, Kullai S, Hassan A. Feeding behavior of giant gourami, *Osphronemus goramy* (Lacepede) larvae. *Songklanakarin Journal of Science and Technology* [Internet]. 2014;36(3):261–4. Available from: <https://www.thaiscience.info/Journals/Article/SONG/10968306.pdf>
11. Arifin OZ, Imron I, Asependi A, Hendri A, Muslim N, Yani A. Hibridisasi intraspesifik antara dua populasi ikan gurami Galunggung (*Osphronemus goramy*, Lacepede, 1801). *Jurnal Riset Akuakultur*. 2018;12(4):315–23.
12. Prakoso VA, Pouil S, Prabowo MNI, Sundari S, Arifin OZ, Subagja J, et al. Effects of temperature on the zootechnical performances and physiology of giant gourami (*Osphronemus goramy*) larvae. *Aquaculture*. 2019;510:160–8.
13. Simanjuntak SBI, Indarmawan I, Wibowo ES. Impact of fed containing different levels of diets supplementation *Spirulina platensis* on growth, haematological, body composition and biochemical parameters, of Gurami (*Osphronemus goramy*). *Turk J Fish Aquat Sci*. 2018;18(5):681–90.
14. FAO. Cultured aquatic species information programme *Osphronemus goramy* (Lacépède, 1801). In: FAO Fisheries and Aquaculture Department [Internet]. 2020. Available from: <http://www.FAO.org/fishery/culturedspecies/Osphr>
15. RI W. International introduction of Inland Aauatic species. Rome, Fao Fisheries Technical Papers. 1988;(294):318.
16. NFDB. *Matsya Bharat*. National Fisheries Development Board, Government of India. 2016;8.
17. Kumari DJ, Babitha B, Jaffar S, Prasad MG, Ibrahim MD, Khan MS. Potential health benefits of *Spirulina platensis*. *Int J Adv Pharm Sci* [Internet]. 2011;2:417–22. Available from: [https://www.researchgate.net/profile/Mohammad-Ibrahim-38/publication/215961590\\_Potential\\_Health\\_Benefits\\_of\\_Spirulina\\_platensis/links/06ab3257153eddb43a4d41ff/Potential-Health-Benefits-of-Spirulina-platensis.pdf](https://www.researchgate.net/profile/Mohammad-Ibrahim-38/publication/215961590_Potential_Health_Benefits_of_Spirulina_platensis/links/06ab3257153eddb43a4d41ff/Potential-Health-Benefits-of-Spirulina-platensis.pdf)
18. Henrikson R. *Spirulina: Health discoveries from the source of life*. 2000.
19. Allen D. Nutrient comparison of *Spirulina* with carrot. Denver, Colorado. [webmaster@microhydron.com](mailto:webmaster@microhydron.com). 2000.
20. Belay A. The potential application of *Spirulina* (*Arthrospira*) as a nutritional and therapeutic supplement in health management. *Journal of the american nutraceutical association (jana)* [Internet]. 2002;5(-). Sid:26–48. Available from: <https://sid.ir/paper/546615/en>
21. Gutiérrez-Salmeán G, Fabila-Castillo L, Chamorro-Cevallos G. Aspectos nutricionales y toxicológicos de *Spirulina* (*arthrospira*). *Nutr Hosp*. 2015;32(1):34–40.
22. Bermejo P, Piñero E, Villar ÁM. Iron-chelating ability and antioxidant properties of phycocyanin isolated from a protean extract of *Spirulina platensis*. *Food Chem* [Internet]. 2008;110(2):436–45. Available from: <http://dx.doi>
23. Phang SM, Miah MS, Yeoh BG, Hashim MA. *Spirulina* cultivation in digested sago starch factory wastewater. *J Appl Phycol*. 2000;12:395–400.

24. Jaime-Ceballos BJ, Hernández-Llamas A, Garcia-Galano T, Villarreal H. Substitution of *Chaetoceros muelleri* by *Spirulina platensis* meal in diets for *Litopenaeus schmitti* larvae. *Aquaculture*. 2006;260(1–4):215–20.
25. Kamble SP, Gaikar RB, Padalia RB, Shinde KD. Extraction and purification of C-phycocyanin from dry *Spirulina* powder and evaluating its antioxidant, anticoagulation and prevention of DNA damage activity. *J Appl Pharm Sci*. 2013;3(8):149–53.
26. Bhat VB, Madyastha KM. Scavenging of peroxynitrite by phycocyanin and phycocyanobilin from *Spirulina platensis*: protection against oxidative damage to DNA. *Biochem Biophys Res Commun*. 2001;285(2):262–6.
27. Biabani Asrami M, Sudagar M, Shahraki N, Vahdat S. Effect of extracted phycocyanin from *Spirulina platensis* on growth parameters, colorations, digestive enzymes and body chemical compositions of Guppy fish (*Poecilia reticulata*). *Survey in Fisheries Sciences* [Internet]. 2019;6(1):1–8. Available from: [https://web.archive.org/web/20200715221920id\\_/http://sifisheriessciences.com/files/site1/user\\_files\\_d60aad/fallahi-A-10-444-56-b3344d0.pdf](https://web.archive.org/web/20200715221920id_/http://sifisheriessciences.com/files/site1/user_files_d60aad/fallahi-A-10-444-56-b3344d0.pdf)
28. Fernández-Rojas B, Medina-Campos ON, Hernández-Pando R, Negrette-Guzmán M, Huerta-Yepez S, Pedraza-Chaverri J. C-phycocyanin prevents cisplatin-induced nephrotoxicity through inhibition of oxidative stress. *Food Funct*. 2014;5(3):480–90.
29. Abdel-Tawab M, Ahmed H. Live *Spirulina* (*Arthrospiraplatensis*) as a growth and immunity promoter for Nile tilapia, *Oreochromis niloticus* (L.), challenged with pathogenic *Aeromonas hydrophila*. *Aquac Res* [Internet]. 2009;40(9):1037–46. Available from: <http://dx.doi>
30. Awad LZ, El-Mahallawy HS, Abdelnaeim NS, Mahmoud MM, Dessouki AA, ElBanna NI. Role of dietary *Spirulina platensis* and betaine supplementation on growth, hematological, serum biochemical parameters, antioxidant status, immune responses, and disease resistance in Nile tilapia. *Fish Shellfish Immunol*. 2022;126:122–30.
31. Mahmoud MM, El-Lamie MM, Kilany OE, Dessouki AA. *Spirulina* (*Arthrospiraplatensis*) supplementation improves growth performance, feed utilization, immune response, and relieves oxidative stress in Nile tilapia (*Oreochromis niloticus*) challenged with *Pseudomonas fluorescens*. *Fish Shellfish Immunol*. 2018;72:291–300.
32. Li XL, Xu G, Chen T, Wong YS, Zhao HL, Fan RR, et al. Phycocyanin protects INS-1E pancreatic beta cells against human islet amyloid polypeptide-induced apoptosis through attenuating oxidative stress and modulating JNK and p38 mitogen-activated protein kinase pathways. *Int J Biochem Cell Biol*. 2009;41(7):1526–35.
33. Rabadiya B, Patel P. *Spirulina*: Potential clinical therapeutic application. *J of Pharmacy Research*. 2010;3(8):1726–32.
34. El-Sheekh M, El-Shourbagy I, Shalaby S, Hosny S. Effect of feeding *Arthrospiraplatensis* (*Spirulina*) on growth and carcass composition of hybrid red tilapia (*Oreochromis niloticus* x *Oreochromis mossambicus*). *Turk J Fish Aquat Sci*. 2014;14(2):471–8.

35. Kim SS, Rahimnejad S, Kim KW, Lee KJ. Partial replacement of fish meal with *Spirulina pacifica* in diets for parrot fish (*Oplegnathus fasciatus*). *Turk J Fish Aquat Sci.* 2013;13(2).
36. Ibrahim MD, Mohamed MF, Ibrahim MA. The role of *Spirulina platensis* (*Arthrospiraplatensis*) in growth and immunity of Nile tilapia (*Oreochromis niloticus*) and its resistance to bacterial infection. 5(6). 2013;109–17.
37. Jana A, Saroch JD, Borana K. Effect of *Spirulina* as a feed supplement on survival and growth of *Pangasius sutchi*. *Int J Fish Aquat Stud [Internet]*. 2014;1(5):77–9. Available from: <https://www.fisheriesjournal.com/vol1issue5/pdf/98.1.pdf>
38. Simanjuntak SBI, Soedibya PHT, Wibowo ES. Growth performance of gurami (*Osphronemus gouramy* Lac.) the given phytoplankton *Spirulina platensis* and *Chlorella vulgaris*. 2014.
39. Aryani N, Mardiah A, Syandri H. Influence of feeding rate on the growth, feed efficiency and carcass composition of the Giant gourami (*Osphronemus goramy*). *Pak J Zool [Internet]*. 2017;49(5). Available from: [https://www.researchgate.net/profile/Ainul-Mardiah-3/publication/319604781\\_Influence\\_of\\_Feeding\\_Rate\\_on\\_the\\_Growth\\_Feed\\_Efficiency\\_and\\_Carcass\\_Composition\\_of\\_the\\_Giant\\_Gourami\\_Osphronemus\\_goramy/links/5ab5caf745851515f59a8124/Influence-of-Feeding-Rate-on-the-Growth-Feed-Efficiency-and-Carcass-Composition-of-the-Giant-Gourami-Osphronemus-goramy.pdf](https://www.researchgate.net/profile/Ainul-Mardiah-3/publication/319604781_Influence_of_Feeding_Rate_on_the_Growth_Feed_Efficiency_and_Carcass_Composition_of_the_Giant_Gourami_Osphronemus_goramy/links/5ab5caf745851515f59a8124/Influence-of-Feeding-Rate-on-the-Growth-Feed-Efficiency-and-Carcass-Composition-of-the-Giant-Gourami-Osphronemus-goramy.pdf)
40. Budi DS, Alimuddin A, Suprayudi MA. Growth response and feed utilization of giant gourami (*Osphronemus goramy*) juvenile feeding different protein levels of the diets supplemented with recombinant growth hormone. *Hayati.* 2015;22(1):12–9.
41. Nofyan E. Pengaruh Pemberian Pakan Dari Sumber Nabati Dan Hewani Terhadap Berbagai Aspek Fisiologi Ikan Gurami (*Osphronemus gouramy* L.) [the Effect of Plant and Animal Food on Various Physiological Aspects of Giant Gourami (*Osphronemus gouramy* L.)]. *J Iktiologi Indones [Internet]*. 2005;5(1):19–23. Available from: <https://doi.org/10.32491/jii.v5i1.297>
42. Yandes Z, Affandi R. Pengaruh pemberian selulos dalam pakan terhadap kondisi biologis benih ikan gurami (*Osphronemus gouramy* Lac) [Effect of cellulose in dietary on the biological condition of giant gourami fry (*Osphronemus goramy* Lac)]. *J Iktiologi Indones [Internet]*. 2003;3(1):27–33. Available from: <https://doi.org/10.32491/jii.v3i1.269>
43. Nutrient N. requirements of fish [Internet]. Washington DC: National Academy Press; 1993. Available from: [https://books.google.co.in/books?hl=en&lr=&id=M2aBtOUhRagC&oi=fnd&pg=PA1&dq=NRC+Nutrient+requirements+of+fish.++\(1993\).+Washington+DC:+National+Academy+Press.&ots=ZXhiJPPiso&sig=ZyWycMmB4zMySw5YLTh1YqYXfgE&redir\\_esc=y#v=onepage&q=NRC](https://books.google.co.in/books?hl=en&lr=&id=M2aBtOUhRagC&oi=fnd&pg=PA1&dq=NRC+Nutrient+requirements+of+fish.++(1993).+Washington+DC:+National+Academy+Press.&ots=ZXhiJPPiso&sig=ZyWycMmB4zMySw5YLTh1YqYXfgE&redir_esc=y#v=onepage&q=NRC)
44. Adel M, Yeganeh S, Dadar M, Sakai M, Dawood MA. Effects of dietary *Spirulina platensis* on growth performance, humoral and mucosal immune responses and disease resistance in juvenile great sturgeon (*Huso huso* Linnaeus, 1754). *Fish Shellfish Immunol.* 2016;56:436–44.

45. Castell JD, Tiews K. Report of the EIFAC, IUNS and ICES Working Group on Standardization of Methodology in Fish Nutrition Research, Hamburg, Federal Republic of Germany, 21-23 March 1979 (No [Internet]. 1980. Available from: <https://agris.fao.org/search/en/providers/122621/records/64774bfaf2e6fe92b36378d3>
46. Stuart JS, Hung SS. Growth of juvenile white sturgeon (*Acipenser transmontanus*) fed different proteins. *Aquaculture*. 1989;76(3-4):303-16.
47. AOAC. Official methods of analysis. In: W Horwitz (Ed). 13th ed. Washington, DC: Academic Press; 2000.
48. Nishikimi M, Rao NA, Yagi K. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. *Biochem Biophys Res Commun*. 1972;46(2):849-54.
49. Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem*. 1978;86(1):271-8.
50. Ag G. Determination of serum proteins by means of the biuret reactions. *J Biol Chem*, [Internet]. 1949;177:51. Available from: [https://cir.nii.ac.jp/crid/1573387450479762048# Citations\\_container](https://cir.nii.ac.jp/crid/1573387450479762048# Citations_container)
51. Doumas BT, Watson WA, Biggs HG. Albumin standards and the measurement of serum albumin with bromocresol green. *Clin Chim Acta*. 1971;258(1):21-30.
52. Allain CC, Poon LS, Chan CS, Richmond WFPC, Fu PC. Enzymatic determination of total serum cholesterol. *Clin Chem*. 1974;20(4):470-5.
53. McGowan MW, Artiss JD, Strandbergh DR, Zak B. A peroxidase-coupled method for the colorimetric determination of serum triglycerides. *Clin Chem*. 1983;29(3):538-42.
54. Li M, Wu W, Zhou P, Xie F, Zhou Q, Mai K. Comparison effect of dietary astaxanthin and *Haematococcus pluvialis* on growth performance, antioxidant status and immune response of large yellow croaker *Pseudosciaenacrocea*. *Aquaculture*. 2014;434:227-32.
55. Bergmeyer HU, Grassi M. Methods of enzymatic analysis [Internet]. 1983. Available from: [https://books.google.co.in/books?id=GDd2zYuLpRwC&dq=Bergmeyer,+H.U.,+Grassi,+M.+\(1983\).+Methods+of+enzymatic+analysis.+Vol.+2.+Weinheim:+Verlag+Chemie.&lr=&source=gbs\\_navlinks\\_s](https://books.google.co.in/books?id=GDd2zYuLpRwC&dq=Bergmeyer,+H.U.,+Grassi,+M.+(1983).+Methods+of+enzymatic+analysis.+Vol.+2.+Weinheim:+Verlag+Chemie.&lr=&source=gbs_navlinks_s)
56. Worthington V. Worthington enzyme manual: enzymes and related biochemicals. Worthington Chemical, New Jersey. 1993;399.
57. Borlongan IG. Studies on the digestive lipases of milkfish, *Chanoschanos*. *Aquaculture*. 1990;89(3-4):315-25.
58. APHA. Standard Methods for Examination of Water and Wastewater. 19th ed. Washington DC, USA: American Public Health Association; 1995.
59. Abdulrahman NM, Ameen HJH. Replacement of fishmeal with microalgae *Spirulina* on common carp weight gain, meat and sensitive composition and survival. *Pakistan Journal of Nutrition*. 2014;13(2):93.

60. Abdulrahman NM. Evaluation of *Spirulina* spp. as food supplement and its effect on growth performance of common carp fingerlings. *Int J Fish Aquat Stud* [Internet]. 2014;2(2):89–92. Available from: <https://www.fisheriesjournal.com/archives/2014/vol2issue2/PartB/2.pdf>
61. Hassanpour S, Chaichisemsari M, Riyazi SR. Effect of replacing different levels of Soybean meal with *Spirulina* on performance in Rainbow Trout. *Ann Biol Res* [Internet]. 2011;2(3):374–9. Available from: <https://d1wqtxts1xzle7.cloudfront.net/107724245/effect-of-replacing-different-levels-of-soybean-meal-with-spirulina-on-performance-in-rainbow-trout-libre.pdf?1700756584=&response-content-disposition=inline>
62. Zeinab AK, Aly MS, Faiza AK, Fatma EM. Effect of *Spirulina platensis* and *Lactobacillus rhamnosus* on growth and biochemical performance of Nile Tilapia (*Oreochromis niloticus*) fingerlings. *Int J Curr Microbiol Appl Sci* [Internet]. 2015;4(4):747–63. Available from: <http://www.ijcmas.com/vol-4-4/Zeinab>
63. Abu-Elala NM, Galal MK, Abd-Elsalam RM, Mohey-Elsaeed O, Ragaa NM. Effects of dietary supplementation of *Spirulina platensis* and garlic on the growth performance and expression levels of immune-related genes in Nile tilapia (*Oreochromis niloticus*). *J Aquac Res Dev* [Internet]. 2016;7(7):433–42. Available from: [https://www.researchgate.net/profile/Nermeen-Abu-Elala/publication/306047814\\_Effects\\_of\\_Dietary\\_Supplementation\\_of\\_Spirulina\\_platensis\\_and\\_Garlic\\_on\\_the\\_Growth\\_Performance\\_and\\_Expression\\_Levels\\_of\\_Immune-related\\_Genes\\_in\\_Nile\\_tilapia\\_Oreochromis\\_niloticus/links/5f439eec458515b729499b83/Effects-of-Dietary-Supplementation-of-Spirulina-platensis-and-Garlic-on-the-Growth-Performance-and-Expression-Levels-of-Immune-related-Genes-in-Nile-tilapia-Oreochromis-niloticus.pdf](https://www.researchgate.net/profile/Nermeen-Abu-Elala/publication/306047814_Effects_of_Dietary_Supplementation_of_Spirulina_platensis_and_Garlic_on_the_Growth_Performance_and_Expression_Levels_of_Immune-related_Genes_in_Nile_tilapia_Oreochromis_niloticus/links/5f439eec458515b729499b83/Effects-of-Dietary-Supplementation-of-Spirulina-platensis-and-Garlic-on-the-Growth-Performance-and-Expression-Levels-of-Immune-related-Genes-in-Nile-tilapia-Oreochromis-niloticus.pdf)
64. Teimouri M, Amirkolaie AK, Yeganeh S. The effects of *Spirulina platensis* meal as a feed supplement on growth performance and pigmentation of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*. 2013;396:14–9.
65. Hanel H, Broekman D, deGraaf S, Schnack D. S, Schnack D. Partial replacement of fishmeal by the lyophilized powder of the microalgae *Spirulina platensis* in Pacific white shrimp diets” *Open Mar Biol J*. 2007;1:1–5.
66. Wang L, Pan B, Sheng J, Xu J, Hu Q. Antioxidant activity of *Spirulina platensis* extracts by supercritical carbon dioxide extraction. *Food Chem*. 2007;105(1):36–41.
67. Saranraj P, Sivasakthi S. *Spirulina platensis*—food for future: a review. *Asian J Pharm Sci Technol* [Internet]. 2014;4(1):26–33. Available from: <https://d1wqtxts1xzle7.cloudfront.net/82407660/26-33-libre.pdf?1647806087=&response-content-disposition=inline>
68. Velasquez SF, Chan MA, Abisado RG, Traifalgar RFM, Tayamen MM, Maliwat GCF, et al. Dietary *Spirulina* (*Arthrospiraplatensis*) replacement enhances performance of juvenile Nile tilapia (*Oreochromis niloticus*). *J Appl Phycol*. 2016;28:1023–30.
69. Tulli F, ChiniZittelli G, Giorgi G, Poli BM, Tibaldi E, Tredici MR. Effect of the inclusion of dried *Tetraselmissuecica* on growth, feed utilization, and fillet composition of European sea bass juveniles fed organic diets. *Journal of aquatic food product technology*. 2012;21(3):188–97.

70. Nakagawa H. Effect of dietary algae on improvement of lipid metabolism in fish. *Biomedicine & pharmacotherapy*. 1997;51(8):345–8.
71. Ergün S, Soyutürk M, Güroy B, Güroy D, Merrifield D. Influence of *Ulva* meal on growth, feed utilization, and body composition of juvenile Nile tilapia (*Oreochromis niloticus*) at two levels of dietary lipid. *Aquaculture International*. 2009;17:355–61.
72. El Gammal MA, Shalaby AM, Ramadan AA. The use of spirulina platensis (microalgae) as a food additive to stimulate growth and immunity modulation for Nile tilapia (*Oreochromis niloticus*) challenged with pathogenic bacteria (*Pseudomonas fluorescens*). *Abbassa International Journal for Aquaculture* [Internet]. 2010;3(1):53–83. Available from: <https://abbassajournal.com/magazine/wp-content/uploads/2017/07/THE-USE-OF-SPIRULINA-PLATENSIS-MICROALGAE-AS-FOOD-ADDITIVES-TO-STIMULATE-GROWTH-AND-IMMUNITY-MODULATION-FOR-NILE-TILAPIA.pdf>
73. Promya J, Chitmanat C. The effects of *Spirulina platensis* and *Cladophora* algae on the growth performance, meat quality and immunity stimulating capacity of the African sharp-tooth catfish (*Clarias gariepinus*). *Int J Agric Biol* [Internet]. 2011;13:77–82. Available from: [https://www.researchgate.net/profile/Chitmanat-Chanagun/publication/279600728\\_The\\_effects\\_of\\_Spirulina\\_platensis\\_and\\_Cladophora\\_Algae\\_on\\_the\\_Growth\\_Performance\\_Meat\\_Quality\\_and\\_Immunity\\_Stimulating\\_Capacity\\_of\\_the\\_African\\_Sharptooth\\_Catfish\\_Clarias\\_gariepinus/links/641db79392cfd54f8428906c/The-effects-of-Spirulina-platensis-and-Cladophora-Algae-on-the-Growth-Performance-Meat-Quality-and-Immunity-Stimulating-Capacity-of-the-African-Sharptooth-Catfish-Clarias-gariepinus.pdf?\\_sg](https://www.researchgate.net/profile/Chitmanat-Chanagun/publication/279600728_The_effects_of_Spirulina_platensis_and_Cladophora_Algae_on_the_Growth_Performance_Meat_Quality_and_Immunity_Stimulating_Capacity_of_the_African_Sharptooth_Catfish_Clarias_gariepinus/links/641db79392cfd54f8428906c/The-effects-of-Spirulina-platensis-and-Cladophora-Algae-on-the-Growth-Performance-Meat-Quality-and-Immunity-Stimulating-Capacity-of-the-African-Sharptooth-Catfish-Clarias-gariepinus.pdf?_sg)
74. Khalil SR, Reda RM, Awad A. Efficacy of *Spirulina platensis* diet supplements on disease resistance and immune-related gene expression in *Cyprinus carpio* L. exposed to herbicide atrazine. *Fish Shellfish Immunol*. 2017;67:119–28.
75. Yu W, Wen G, Lin H, Yang Y, Huang X, Zhou C, et al. Effects of dietary *Spirulina platensis* on growth performance, hematological and serum biochemical parameters, hepatic antioxidant status, immune responses and disease resistance of Coral trout *Plectropomus leopardus* (Lacepede, 1802). *Fish Shellfish Immunol*. 2018;74:649–55.
76. Hegazi SM, Fadl SE, Gad DM, Abeerd EKM. Effect of *Spirulina* incorporated in diets of tilapia nilotica fresh water fish exposed to lead toxicity. *J Vet Sci* [Internet]. 2014;1:633–49. Available from: [https://www.researchgate.net/profile/EI-Keredy-Abeer/publication/350008720\\_Effect\\_of\\_spirulina\\_incorporated\\_in\\_diets\\_of\\_Tilapia\\_nilotica\\_fresh\\_water\\_fish\\_exposed\\_to\\_lead\\_toxicity/links/604b5def299bf1f5d84171ad/Effect-of-spirulina-incorporated-in-diets-of-Tilapia-nilotica-fresh-water-fish-exposed-to-lead-toxicity.pdf](https://www.researchgate.net/profile/EI-Keredy-Abeer/publication/350008720_Effect_of_spirulina_incorporated_in_diets_of_Tilapia_nilotica_fresh_water_fish_exposed_to_lead_toxicity/links/604b5def299bf1f5d84171ad/Effect-of-spirulina-incorporated-in-diets-of-Tilapia-nilotica-fresh-water-fish-exposed-to-lead-toxicity.pdf)
77. Sayed AEDH, Fawzy MA. Effect of dietary supplementation of *Spirulina platensis* on the growth and haematology of the catfish *Clarias gariepinus*. *J Adv Biol* [Internet]. 2014;5(2):625–35. Available from: [https://d1wqtxts1xzle7.cloudfront.net/35169553/Sayed\\_and\\_Fawzy\\_2014a-libre.pdf?1413550140=&response-content-disposition=inline](https://d1wqtxts1xzle7.cloudfront.net/35169553/Sayed_and_Fawzy_2014a-libre.pdf?1413550140=&response-content-disposition=inline)
78. Zhang C. The effects of polysaccharide and phycocyanin from *Spirulina platensis* variety on peripheral blood and hematopoietic system of bone marrow

- in mice. In: Second Asia-Pacific Conference on Alga Biotechnology. 1994. p. 25–7.
79. Abdulrahman NM, Hama Ameen HJ, Hama SR, Hassan BR, Nader PJ. Effect of microalgae *Spirulina* spp. as food additive on some biological and blood parameters of common carp *Cyprinus carpio* L. *Iraqi Journal of Veterinary Sciences* [Internet]. 2019;33(1):27–31. Available from: <https://www.iasj.net/iasj/download/15342ccc1690df08>
  80. Schäperclaus W, Kulow H, Schreckenbach K. *Fish Disease*—AA Balkema. the Netherlands: Rotterdam; 1992.
  81. Youssef IM, Saleh ES, Tawfeek SS, Abdel-Fadeel AA, Abdel-Razik ARH, Abdel-Daim AS. Effect of *Spirulina platensis* on growth, hematological, biochemical, and immunological parameters of Nile tilapia (*Oreochromis niloticus*). *Trop Anim Health Prod*. 2023;55(4):275.
  82. Abdel-Daim MM, Dawood MA, Elbadawy M, Aleya L, Alkahtani S. *Spirulina platensis* reduced oxidative damage induced by chlorpyrifos toxicity in Nile tilapia (*Oreochromis niloticus*). *Animals*. 2020;10(3):473.
  83. Akbary P, Molzaei E, Aminikhoei Z. Effect of dietary supplementation of *Ulva rigida* C. Agardh extract on several of physiological parameters of grey mullet, *Mugil cephalus* (Linnaeus). *Sustainable Aquaculture and Health Management Journal*. 2018;4(1):59–68.
  84. Yeganeh S, Adel M. Effects of dietary algae (*Sargassum ilicifolium*) as immunomodulator and growth promoter of juvenile great sturgeon (*Huso huso* Linnaeus, 1758). *J Appl Phycol*. 2019;31(3):2093–102.
  85. Baker ME. Beyond carrier proteins: albumin, steroid hormones and the origin of vertebrates. *J Endocrinol* [Internet]. 2002;175:121–7. Available from: [https://www.researchgate.net/profile/Michael-Baker-11/publication/11079965\\_Albumin\\_steroid\\_hormones\\_and\\_the\\_origin\\_of Vertebrates/links/00b4951d6f94ea8476000000/Albumin-steroid-hormones-and-the-origin-of-vertebrates.pdf](https://www.researchgate.net/profile/Michael-Baker-11/publication/11079965_Albumin_steroid_hormones_and_the_origin_of Vertebrates/links/00b4951d6f94ea8476000000/Albumin-steroid-hormones-and-the-origin-of-vertebrates.pdf)
  86. Al-Deriny SH, Dawood MA, Abou Zaid AA, Wael F, Paray BA, Van Doan H, et al. The synergistic effects of *Spirulina platensis* and *Bacillus amyloliquefaciens* on the growth performance, intestinal histomorphology, and immune response of Nile tilapia (*Oreochromis niloticus*). *Aquac Rep*. 2020;17:390–7.
  87. Xiang Y, Shuning H, Lingke Z, Cheng H. The analysis of the effect of reducing triglyceride level in serum of hyperlipidemia model rats by feeding *Spirulina maxima*. *Journal of Nanjing University (natural Sciences)* [Internet]. 2002;38(2):182–6. Available from: <https://europepmc.org/article/cba/383078>
  88. Colla LM, Muccillo-Baisch AL, Costa JA V. *Spirulina platensis* effects on the levels of total cholesterol, HDL and triacylglycerols in rabbits fed with a hypercholesterolemic diet. *Brazilian Archives of Biology and Technology*. 2008;51:405–11.
  89. Mg M. Effects of *Ascophyllum* and *Spirulina* meal as feed additives on growth performance and feed utilization of red sea bream, *Pagrus major*. *J Fac Appl Biol Sci* [Internet]. 1994;33:125–32. Available from: <https://cir.nii.ac.jp/crid/1573105975387643904>
  90. Brito R, Chimal ME, Gelabert R, Gaxiola G, Rosas C. Effect of artificial and natural diets on energy allocation in *Litopenaeus setiferus* (Linnaeus, 1767) and

- Litopenaeusvannamei (Boone, 1931) early postlarvae. *Aquaculture*. 2004;237(1-4):517-31.
91. Anand PS, Kohli MPS, Roy SD, Sundaray JK, Kumar S, Sinha A, et al. Effect of dietary supplementation of periphyton on growth performance and digestive enzyme activities in *Penaeus monodon*. *AQUACULTURE*. 2013;392:59-68.
  92. Cahu CL, Infante JZ, Peres A, Quazuguel P, Le Gall MM. Algal addition in sea bass (*Dicentrarchuslabrax*) larvae rearing: effect on digestive enzymes. *Aquaculture*. 1998;161(1-4):479-89.
  93. Lu J, Takeuchi T. Spawning and egg quality of the tilapia *Oreochromisniloticus* fed solely on raw *Spirulina* throughout three generations. *Aquaculture*. 2004;234(1-4):625-40.
  94. Teimouri M, Yeganeh S, Mianji GR, Najafi M, Mahjoub S. The effect of *Spirulina platensis* meal on antioxidant gene expression, total antioxidant capacity, and lipid peroxidation of rainbow trout (*Oncorhynchus mykiss*). *Fish PhysiolBiochem*. 2019;45:977-86.
  95. Misra S, Niyogi S. Selenite causes cytotoxicity in rainbow trout (*Oncorhynchus mykiss*) hepatocytes by inducing oxidative stress. *Toxicology in vitro*. 2009;23(7):1249-58.
  96. Abdel-Latif HM, Khalil RH. Evaluation of two phytobiotics, *Spirulina platensis* and *Origanum vulgare* extract on growth, serum antioxidant activities and resistance of Nile tilapia (*Oreochromisniloticus*) to pathogenic *Vibrio alginolyticus*. *Int J Fish Aquat Stud*. 2014;250:250-5.
  97. Xie S, Chen P, Lv Z, Lan Z, Zou J. Effects of *Spirulina platensis* on muscle nutritive components and growth of *Carassius auratus*. *Journal of Zhongkai University of Agriculture and Engineering [Internet]*. 2015;28(2):9-21. Available from: <http://znjx.cbpt.cnki.net>
  98. Tayag CM, Lin YC, Li CC, Liou CH, Chen JC. Administration of the hot-water extract of *Spirulina platensis* enhanced the immune response of white shrimp *Litopenaeusvannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol*. 2010;28(5-6):764-73.
  99. Lin YC, Tayag CM, Huang CL, Tsui WC, Chen JC. White shrimp *Litopenaeusvannamei* that had received the hot-water extract of *Spirulina platensis* showed earlier recovery in immunity and up-regulation of gene expressions after pH stress. *Fish Shellfish Immunol*. 2010;29(6):1092-8.
  100. Abdelkhalek NK, Ghazy EW, Abdel-Daim MM. Pharmacodynamic interaction of *Spirulina platensis* and deltamethrin in freshwater fish Nile tilapia, *Oreochromisniloticus*: impact on lipid peroxidation and oxidative stress. *Environmental Science and Pollution Research*. 2015;22:3023-31.
  101. Buege JA, Aust SD. [30] Microsomal lipid peroxidation Methods in enzymology. In: Academic press. Academic press; 1978. p. 302-10.
  102. Liu B, Xie J, Ge X, Xu P, Wang A, He Y, et al. Effects of anthraquinone extract from *Rheum officinale* Bail on the growth performance and physiological responses of *Macrobrachiumrosenbergii* under high temperature stress. *Fish Shellfish Immunol*. 2010;29(1):49-57.

103. Lin YH, Shiau SY. Dietary vitamin E requirement of grouper, *Epinephelus malabaricus*, at two lipid levels, and their effects on immune responses. *Aquaculture*. 2005;248(1–4):235–44.
104. Luo P. Effect of Spirulina on the growth performance of *Cyprinus carpio* var. Jian. *J Hydrol*. 2006;26(2006):41–2.
105. Moreira LM, Behling B, Rodrigues RS, Costa JA V, Souza-Soares LA. Spirulina as a protein source in the nutritional recovery of wistar Rats. *Brazilian Archives of Biology Technology*. 2013;56(3):447–56.
106. Karakatsouli N. An overview of the use of fatty acids in the fish farming research during the last decade, with particular emphasis on fish quality. *Journal of World Aquaculture Society*. 2012;43:291–320.
107. Bhavan PS, Devi VG, Shanti R, Radhakrishnan S, Poongodi R. Basic biochemical constituents and profiles of amino acids in the post larvae of *Macrobrachium rosenbergii* fed with Spirulina and yeast enriched Artemia. *J Sci Res [Internet]*. 2010;2(3):539–49. Available from: <http://dx.doi>
108. Prakoso VA, Pouil S, Cahyanti W, Sundari S, Arifin OZ, Subagja J, et al. Fluctuating temperature regime impairs growth in giant gourami (*Osphronemus goramy*) larvae. *Aquaculture*. 2021;539:606–736.
109. Syandri H, Mardiah A, Aryani N. Effect of stocking density on the performance of juvenile gurami sago (*Osphronemus goramy*) in the synthetic sheet pond. *Pak J Zool [Internet]*. 2020;52(2):717. Available from: <https://www.proquest.com/openview/82b67628eb7695544259a655627e3825/1?pq-origsite=gscholar&cbl=616534>
110. Boyd CE, Tucker CS, Somridhivej B. Alkalinity and hardness: critical but elusive concepts in aquaculture. *J World Aquac Soc*. 2016;47(1):6–41.