

Impact of *Spirulina platensis* Supplementation on Growth, Health, and Antioxidant Defence in Giant Gourami (*Osphronemus goramy*)

ABSTRACT

A 90-day trial investigated the effects of dietary *Spirulina platensis* (SP) supplementation on growth performance, haematology, serum biochemistry, digestive enzymes, and hepatic antioxidants in giant gourami (*Osphronemus goramy*). Fish (6.96 ± 0.011 g) were distributed into six treatments with 3 replicates each (15 fish per replicate) across 18, 60 L aquaria. Six isonitrogenous diets with 0% (control), 1% (SP1), 2% (SP2), 3% (SP3), 4% (SP4), and 5% (SP5) SP were provided *ad libitum* twice daily. Results showed dose-dependent improvements in final weight, weight gain, weight gain rate, protein efficiency ratio, and specific growth rate, with the highest values at 5% SP ($P < 0.05$). Feed conversion ratio decreased with higher SP, indicating enhanced efficiency. Body indices increased significantly with higher SP ($P < 0.05$). Haematological parameters, including WBC (white blood cells), HB (hemoglobin), HCT (hematocrit), monocyte %, and neutrophil %, improved with increasing SP significantly ($P < 0.05$). Serum biochemistry showed markedly reduced cholesterol and triglycerides concentration, and increased total protein, albumin, and globulin levels with higher SP ($p < 0.05$). Intestinal amylase and lipase activities increased significantly with SP ($p < 0.05$), while protease remained consistent. Hepatic antioxidants revealed decreased MDA (malondialdehyde) and increased SOD (Superoxide dismutase) with higher SP ($P < 0.05$). Carcass analysis showed increased protein, fibre, and fat content, and decreased ash and carbohydrates with higher SP. These findings suggest that 5% SP enhances growth, biochemical parameters, haematology, digestive, and antioxidant mechanisms in *O. goramy*, supporting its potential as a dietary supplement in aquaculture.

Keywords: *Spirulina*, Giant Gourami, Growth, Haematology, Immune Response, Digestive Enzyme

1. INTRODUCTION

In this modern era, the decline in the production of capture fisheries contrasts with the escalating demand for fish products for human consumption. Consequently, the Food and Agriculture Organization anticipates that aquaculture activities will contribute 60% of future fisheries production, a trend projected to persist and increase (1). The expansion of aquaculture to include a diverse range of fish species has amplified the requirement for commercial feed (2, 3, 4 and 5). Approximately 70% of animal production in aquaculture relies on commercial feeds rich in protein (2, 3, 4, 5 and 6)

The giant gourami, *Osphronemus goramy*, stands out as a favoured species for both pisciculture (7) and the ornamental industry, primarily owing to its impressive size, reaching nearly 2 feet in length, its adaptability to freshwater as well as brackish water, and its suitability for live fish trade (8). This fish exhibits omnivorous behaviour with a strong inclination towards vegetarianism, demonstrating relative robustness and adaptability to adverse environmental conditions such as aerial respiration, which renders it highly appealing for low-input aquaculture (9). While the production of giant gourami fry relies on the natural spawning of captive broodfish held in ponds, efforts to standardize commercial-scale propagation of the species are still underway (10–12). It is noteworthy that the growth rate of gourami is relatively slow, requiring 18 months to attain a weight of 500 grams (13). Native to Southeast Asia, particularly from Indonesia, the Malay Peninsula, Thailand, and the Mekong basin, the giant gourami *Osphronemus goramy* Lacépède, 1801, has been introduced to numerous other countries including India, Pakistan, Sri Lanka, Mauritius, Seychelles, Madagascar, Uganda, Philippines, Australia, New Zealand, Papua New Guinea, Japan, Hawaii, Colombia, France, Italy, among others, for aquaculture purposes (14,15) In India, the giant gourami was first introduced from Mauritius and Java (Indonesia) to Chennai during 1865–66 and subsequently distributed to various regions of the country (8). The cultivation of this species for food is particularly well-established in the state of Kerala, situated on the south-west coast. Furthermore, the National Fisheries Development Board, Government of India, is actively promoting this fish as one of the potential candidate species for both aquaculture and the ornamental fish trade in India (16).

Spirulina platensis is a type of blue-green algae characterized by its filamentous, spiral shape and remarkable nutritional value. Its cell wall possesses a soft texture due to a coating of complex

sugars and proteins, facilitating easy digestion. This alga boasts an impressive nutrient composition, with approximately 62% protein content, as well as vitamins and antioxidants such as β -carotene and xanthophyll (17). (18) highlighted the presence of Gamma Linolenic Acid (GLA) and iron in *S. platensis*, alongside chlorophyll, glycogen, minerals (19), and phycocyanin and porphirin(20). Furthermore, it is rich in essential amino acids, vitamins B12 and β -carotene, phosphorus, iron, and calcium (21). *Spirulina platensis* is also known for its bioactive compounds with antioxidant properties, making it a valuable supplement for enhancing food (22) and serving as a significant source of micronutrients due to its high protein content (23).

Spirulina, renowned for its phycocyanin content (24), particularly C-phycocyanin, demonstrates antioxidant properties and may serve as a viable alternative to heparin due to its anticoagulant effects (25). Phycocyanin has been utilized for its antioxidative, antifungal and antitumor characteristics (26,27), as well as its immunostimulatory and hepatoprotective benefits (28). Multiple studies on Nile tilapia have indicated that incorporating *Spirulina* into their diet significantly enhances growth rates, immune response, and disease resistance(29–32). Li et al. demonstrated that phycocyanin augments the activities of SOD and GPx enzymes in pancreatic beta cells(32). The inclusion of *S. platensis* in diets can modulate the immune system(33)facilitate large-scale fish farming (34), and serve as a pivotal component for animal feed (35). Moreover, supplementing fish feed with *S. platensis* has been shown to enhance growth and survival rates (36,37) and can serve as a primary ingredient for feed production (38).

The rise in giant gourami production can be attributed to advancements in production inputs such as feed, seed, disease management, and optimal culture techniques within giant gourami aquaculture. However, a significant challenge in culturing this fish lies in the inefficiency of production systems, largely due to their slow growth rate (39,40). In terms of feed technology development, numerous studies have been undertaken, particularly focusing on identifying alternative protein sources capable of promoting optimal growth performance in giant gourami (41,42). The utilization of chemicals for controlling bacterial or parasitic populations presents drawbacks such as bioaccumulation, pollution, and potential human health risks. Furthermore, the excessive reliance on antibiotics in aquaculture has led to issues of resistance, immunosuppression, and disruption of beneficial bacterial populations.

The utilization of *Spirulina*, known for its nutritional value and bioactive properties, offers a promising avenue for enhancing fish growth performance, immune response, and overall health, thereby contributing to the advancement of sustainable aquaculture practices. By investigating the effects of *S. platensis* supplementation on the growth and health parameters of *Osphronemus goramy*, a commercially important fish species, this study aims to identify potential strategies for improving aquaculture efficiency and reducing reliance on conventional feed additives and antibiotics.

2. MATERIALS AND METHODS

2.1 Water quality

Water quality analysis was conducted three times throughout the maintenance period (on days 1, 30, and last days also taken). A thermometer calibrated in Celsius was utilized to measure water temperature. Dissolved oxygen (O₂; mg L⁻¹) levels were measured using an oxygen meter (YSI Model 52, Yellow Instrument Co, Yellow Spring, OH USA). The pH values of the water were determined using a digital pH meter (Mini 0–14 pH IQ, Scientific Cemo Science, Thailand). The concentrations of nitrite (mg/L), nitrate (mg/L), TAN (mg/L), alkalinity (mg L⁻¹), and hardness (mg L⁻¹) were determined following standard procedures outlined by APHA (58).

2.2 Experimental fish

A collection of 270 superficially robust giant gourami (*Osphronemus goramy*), averaging 6.96 \pm 0.011g in weight, was procured from a seed farm. These fish were subjected to a one-week acclimatization period in laboratory conditions before being randomly distributed into six dietary groups. Each group of juvenile gouramis was accommodated in aquariums 60L, with a stocking density of 15 fish per aquarium. To maintain optimal conditions, each aquarium was equipped with a heater set to maintain a temperature of 30°C and aeration equipment. Before introducing the juveniles, the aquariums underwent sterilization using 30 ppm chlorine. Throughout the one-week acclimatization period, the juveniles were provided with a commercial diet containing 30% protein ad libitum. Body weight measurements were taken individually at the commencement and conclusion of

the maintenance period for each aquarium. To sustain water quality, 25% of the total aquarium water volume was replaced before the morning feeding. Over the 90-days experimental period, fish were weighed every 15 days to adjust the daily feed ration, following the recommendation of the National Research Council (43), which was administered twice daily at 9:00 am and 5:00 pm.

2.3 Experimental Diets

Six different diets were prepared with varying levels of *S. platensis* (SP) ranging from 0% (C), 1% (SP2), 2% (SP2), 3% (SP3), 4% (SP4) and 5% (SP5). Detailed information regarding the formulation of the feeds and their proximate compositions presented in Table 1. Each diet was carefully prepared by weighing the ingredients according to the specified proportions and thoroughly mixing them for a period of 10 minutes. Subsequently, fish oil and plant oil were incorporated into the mixture, which was then passed through a 425 µm sieve. Further mixing was carried out for components containing fish oil for an additional 10 minutes. Afterward, distilled water was added and stirred for 5 minutes before being cooked in an autoclave. Pellets of 2.0 mm to 2.5 mm diameter was produced using a pelletizer through wet-extrusion after autoclave, followed by air-drying. The resulting pellets were then packaged in plastic bags and stored at room temperature.

Table 1. Feed ingredients and the proximate chemical composition (% of dry weight) in the experimental diets.

Treatment	Control	T1	T2	T3	T4	T5
<i>Spirulina p.</i>	0	1	2	3	4	5
Soybean meal	21	21	21	21	21	21
Fish meal	10.5	10.5	10.5	10.5	10.5	10.5
Poultry by-product meal	16.5	16.5	16.5	16.5	16.5	16.5
Rice bran	21.3	20.3	19.3	19.3	18.3	18.3
Wheat flour	17.7	17.7	17.7	17.7	17.7	17.7
Tapioca powder	5	5	5	4	4	3
Plant oil	2	2	2	2	2	2
Fish oil	3	3	3	3	3	3
Vitamin mineral mix ¹	3	3	3	3	3	3
Total	100	100	100	100	100	100
Moisture	8.74	8.84	8.86	8.82	8.76	8.88
Dry matter	91.26	91.16	91.14	91.18	91.24	91.12
Protein	30.399	30.289	30.179	30.179	30.069	30.069
Ether extraction	5.24	5.33	5.27	5.38	5.28	5.3
Ash	6.34	6.25	6.25	6.38	6.38	6.4
Fiber	4.2	4.26	4.24	4.23	4.36	4.3
CHO	43.85065	43.66796	43.7554	43.29211	43.28394	43.00126
NFE** (%) ²	44.98	44.93	44.83	44.74	44.52	44.46
Gross energy (kcal/g) ³	405.26	404.8	403.95	403.08	402.61	401.58

¹Composition of vitamin mineral mixture of 1 kg: Vitamin A - 50,00,000 IU; Vitamin D3 - 10,00,000 IU; Vitamin B2 - 2.0 g; Vitamin E - 750 units; Vitamin K - 1.0 g; Calcium pantothenate 2.5 g; Nicotinamide - 10.0 g; Vitamin B12 - 6.0 g; Choline Chloride - 150.0 g; Calcium - 750.0 g; Manganese - 27.5 g; Iodine - 1.0 g; Iron - 7.5 g; Zinc - 15.0 g; Copper - 2.0 g; Cobalt - 0.45 g.

²NFE: Nitrogen free extract calculated using the following equation: NFE = 100 - (crude protein + ether extract + crude fiber + ash).

³GE: Gross energy calculated on the basis of 23.6, 39.4 and 17.2 k joule gross energy g-1 protein, ether extract and NFE, respectively.

P/E ratio: Protein energy ratio (mg crude protein kJ-1 gross energy) = CP/GE × 1000.

2.4 Growth performance

The daily feed intake (FI) of the fish was monitored consistently throughout the entire experiment. Before being weighed, the fish were carefully dried using fresh filter paper to eliminate any excess moisture. They were then weighed using a digital balance. The parameters were computed using the following formulas:

- 1) Weight gain rate (WGR, %) = $100 \times (\text{Final body weight} - \text{initial body weight}) / \text{Initial body weight}$ (45)
- 2) WG (g fish-1) = Final body weight - Initial weight of the fish (g) (45)
- 3) Specific growth rate (SGR, % day⁻¹) = $100 \times [(\ln \text{ final individual weight} - \ln \text{ initial individual weight}) / \text{number of days}]$

- 4) **Average daily growth** ADG (g fish⁻¹ day⁻¹) = (Final body weight - Initial weight of the fish (g))/Day
- 5) Feed conversion ratio (FCR) = Total diet fed/ total wet weight gain (45)
- 6) Feed efficiency ratio (FER) = 1/FCR
- 7) Survival rate (SR, %) = 100 × (Final fish number/Initial fish number).
- 8) Protein efficiency ratio (PER) = Final biomass (g) - initial biomass (g) /Protein intake (46)

Three fish from each tank were dissected to determine the Viscerosomatic index (GSI%), Hepatosomatic index (HSI%), Visceral fat-somatic indexes (VFSI%), and Bilesomatic index (BSI) utilizing the prescribed formulas:

$$\begin{aligned} \text{GSI} &= 100 [\text{viscera weight (g) / whole body weight (g)}] \\ \text{HSI} &= 100 [\text{liver weight (g) / whole body weight (g)}] \\ \text{VFSI} &= 100 [\text{visceral fat weight (g) / whole body weight (g)}] \\ \text{BSI} &= 100 [\text{Bile weight (g) / weight of liver (g)}] \end{aligned}$$

2.5 Proximate analysis

The fish body samples collected before infection, comprising three fish from each replicate (**totalling** nine from each group), underwent chemical analysis for moisture, protein, fat, and ash using the standard method recommended by the Association of Official Analytical Chemists (47). Approximately **nine** fish per treatment were included in the analysis. Dry matter (DM) content was determined by drying the samples in an oven at 105°C for 24 hours. Crude protein was quantified using the micro-Kjeldhal method, with nitrogen content multiplied by 6.25 (utilizing a Kjeltach autoanalyzer, Model VELP Scientifica, UDK 127). Crude lipids were assessed using the Soxhlet method with ether extraction (Model VELP Scientifica, SER 148), while ash content was determined by burning samples at 550°C for 16 hours. Crude fiber (CF) analysis was conducted after digestion with 5% sulfuric acid and 5% sodium hydroxide for 15 minutes; the residues were subsequently dried and ashed. The nitrogen-free extract (NFE) was calculated using the formula: $\text{NFE} = 100 - (\text{CP} + \text{EE} + \text{CF} + \text{ash})$.

2.6 Sample collection

At the end of 90-day experiment period, all fish underwent a 24-hour fasting period before being subjected to the final sampling. Fish samples were randomly collected from each treatment group and anesthetized using Tricaine methanesulfonate. Blood samples (**2ml/replica**) were then obtained from the caudal blood vessel using a 1 ml syringe (MDI Europa GmbH, Germany). After blood collection, the blood samples were immediately divided into two equal parts. One part was transferred to a tube containing an anti-coagulant (heparin) for hematological analysis, while the other part was transferred to non-heparinized tubes for biochemical and other analyses. The samples were then centrifuged at 4°C with 3000 rpm for 10 minutes and stored at -80 °C until further use, following the protocol outlined by Adel et al (44). Additionally, carcass composition analysis was conducted on three fish samples, which were stored at -20°C.

2.7 Hematological parameters analysis

Red blood cell (RBC, 10⁶ mm³) and white blood cell (WBC, 10³ mm³) counts, hemoglobin (Hb, g dL⁻¹), and hematocrit (Ht %) were measured according to the method detailed by Li et al. (54). Differential leukocytic counts (neutrophils, lymphocytes, and monocytes) were conducted using an automated blood cell counter.

2.8 Biochemical Measurements

Superoxide dismutase (SOD Umg/prot) and malondialdehyde (MDA µg/g) levels were determined following the methods outlined by Nishikimi et al(48) and Mihara & Uchiyama (49), respectively. The concentration of total protein (g dL⁻¹) in plasma samples from the various experimental groups was measured using the Biuret method as described by AG(50). Albumin concentration (g dL⁻¹) was assessed using the bromocresol green method (51), while globulin concentration (g dL⁻¹) was computed by subtracting albumin from total protein. The Albumin to Globulin ratio (A/G ratio) was obtained by dividing albumin by globulin. Total cholesterol (TC) and triglycerides (TG) levels were determined using colorimetric diagnostic kits provided by spectrum-

bioscience (Egyptian Company for Biotechnology, Cairo, Egypt), following the protocols of Allain et al(52) and McGowan et al. (53), respectively.

2.9 Enzyme activity

Enzyme activity, such as protease, amylase, and lipase, was determined using the methodologies described by (55), (56), and (57).

2.10 Statistical analysis

The values for each parameter were expressed as the arithmetic mean \pm standard error (SE). The effects of enriched diets on growth performance, hematological, biochemical, digestive enzyme activity, and water quality parameters were assessed using one-way ANOVA. Comparison of the mean values was conducted using Duncan's multiple range tests at a significance level of 0.05%. The software program SPSS (Version 14.0; SPSS) for Windows was employed for the analysis.

3. RESULTS

3.1 Water quality

The water quality of the **aquarium** where giant gourami juveniles were reared was assessed. Parameters including water temperature, dissolved oxygen (DO), total alkalinity, hardness, pH, and nitrates were measured and found to be within the typical range recommended by WHO/FAO standards, as outlined in Table 8.

Table 8. Water quality parameter observed during 90 days of experiment

	Control (0% SP)	T1 (1% SP)	T2 (2% SP)	T3 (3% SP)	T4 (4% SP)	T5 (5% SP)
Temperature (°C)	28.56 \pm 0.088 ^e	28.93 \pm 0.088 ^d	28.6 \pm 0.057 ^e	30.1 \pm 0.057 ^b	30.7 \pm 0.057 ^a	29.3 \pm 0.057 ^c
PH	7.833 \pm 0.008 ^a	7.396 \pm 0.012 ^f	7.496 \pm 0.008 ^e	7.533 \pm 0.008 ^d	7.716 \pm 0.008 ^c	7.793 \pm 0.008 ^b
DO (mg/L)	5.813 \pm 0.012 ^e	6.096 \pm 0.012 ^b	6.24 \pm 0.011 ^a	5.963 \pm 0.012 ^c	5.863 \pm 0.008 ^d	6.266 \pm 0.012 ^a
Total Alkalinity (mg/L as CaCO ₃)	55.33 \pm 0.088 ^d	56.46 \pm 0.088 ^c	59.53 \pm 0.088 ^b	54.73 \pm 0.088 ^e	60.06 \pm 0.088 ^a	59.33 \pm 0.088 ^b
Hardness (mg/L as CaCO ₃)	66.56 \pm 0.088 ^e	68.56 \pm 0.088 ^d	69.23 \pm 0.088 ^c	65.56 \pm 0.088 ^f	69.86 \pm 0.088 ^b	70.23 \pm 0.088 ^a
NO ₃ (mg/L)	0.054 \pm 0.001 ^a	0.031 \pm 0.001 ^c	0.046 \pm 0.001 ^b	0.035 \pm 0.001 ^c	0.047 \pm 0.001 ^b	0.057 \pm 0 ^a
NO ₂ (mg/L)	0.027 \pm 0.002 ^e	0.069 \pm 0.001 ^b	0.021 \pm 0.001 ^f	0.063 \pm 0.001 ^c	0.041 \pm 0.001 ^d	0.079 \pm 0 ^a
TAN (mg/L)	0.606 \pm 0.008 ^e	0.773 \pm 0.008 ^d	0.48 \pm 0.005 ^f	1.506 \pm 0.008 ^b	1.806 \pm 0.008 ^a	0.896 \pm 0.008 ^c

3.2 Growth performance

The growth performance of fish fed with various experimental diets is presented in Table 2. At the conclusion of the 90-day experiment, the results indicated a dose-dependent increase in Final Weight (FW), Weight Gain (WG), Weight Gain Rate (WG%), Feed Efficiency Ratio (FER), Protein Efficiency Ratio (PER), Average Daily Gain (ADG) (g fish⁻¹ day⁻¹), and Specific Growth Rate (SGR). The highest values for FW, WG, WG%, PER, FER, ADG, and SGR were observed in the T5 (5% SP) diet group, showing statistical significance (P<0.05). These parameters were notably higher in the T5 (5% SP) diet group compared to the control group (Control) and other treatments (P < 0.05). Conversely, a dose-related decrease was observed in Feed Conversion Ratio (FCR), with the highest decrease recorded in the T5 (5% SP) diet group (P<0.05). No significant variations were detected in Survival Rate (SR) among all treatment groups (P>0.05). According to the broken line regression model, the optimal SPC supplementation level was determined to be 5% based on the survival rate and growth parameters.

Table 2. Mean (SE) value growth parameter of *Osphronemus goramy* during the 90-day experiment after feeding trial of *Spirulina platensis*.

	Control (0% SP)	T1(1% SP)	T2(2% SP)	T3(3% SP)	T4(4% SP)	T5(5% SP)
Initial weight (IW)	6.926 \pm 0.012 ^d	6.893 \pm 0.008 ^d	6.896 \pm 0.008 ^d	7.01 \pm 0.015 ^b	7.1 \pm 0.011 ^a	6.973 \pm 0.012 ^c
Final weight (FW)	55.16 \pm 0.202 ^f	58.07 \pm 0.12 ^e	60.76 \pm 0.145 ^d	64.1 \pm 0.208 ^c	67.87 \pm 0.145 ^b	69.98 \pm 0.06 ^a

Weight gain (WG) gm	48.24±0.214 ^f	51.173±0.111 ^e	53.87±0.143 ^d	57.09±0.202 ^c	60.766±0.156 ^b	63.01±0.05 ^a
Weight gain Rate (WG%)	696.45±4.29 ^f	742.35±0.67 ^e	781.10±2.163 ^d	814.41±2.79 ^c	855.88±3.594 ^b	903.58±1.102 ^a
Total feed consumption (TFC) gm	96.5±0 ^f	97.4±0 ^e	100.7±0 ^d	104.8±0 ^c	106.9±0 ^b	107.7±0 ^a
Feed conversion ratio (FCR)	2±0.008a	1.903±0.004 ^b	1.869±0.004 ^c	1.835±0.006 ^d	1.759±0.004 ^e	1.709±0.001 ^f
FER	0.499±0.002 ^f	0.525±0.001 ^e	0.534±0.001 ^d	0.544±0.001 ^c	0.568±0.001 ^b	0.585±0 ^a
Protein efficiency ratio (PER)	1.663±0.009 ^f	1.751±0.006 ^e	1.783±0.008 ^d	1.813±0.007 ^c	1.895±0.008 ^b	1.950±0.002 ^a
ADG (g fish-1 day-1)	0.536±0.002 ^f	0.568±0.001 ^e	0.598±0.001 ^d	0.634±0.002 ^c	0.675±0.001 ^b	0.7±0 ^a
Specific growth rate (SGR) (%)	2.305±0.005 ^f	2.367±0 ^e	2.417±0.002 ^d	2.459±0.003 ^c	2.508±0.004 ^b	2.562±0.001 ^a
Survival rate (SR%) (%)	100±0 ^a	100±0 ^a	100±0 ^a	100±0 ^a	100±0 ^a	100±0 ^a




Fig 1. Body weight and weight gain of *Osphronemus goramy* during the 90-day experiment after feeding trial of *Spirulina platensis*.




Fig1: FCR, FER, PER and DGR of *Osphronemus goramy* during the 90-day experiment after feeding trial of *Spirulina platensis*.




Fig2: Specific growth rate (SGR) of *Osphronemus goramy* during the 90-day experiment after feeding trial of *Spirulina platensis*.

3.2. Body indices

The body indices of fish fed with varying *spirulina* concentration diets are listed in Table 3. The hepatosomatic index (HSI) exhibited a significant increase from **Control** (0% SP) to T5 (5% SP) over the experimental period ($P < 0.05$). Similarly, the viscerosomatic index (GSI%) showed a significant upward trend from **Control** (0% SP) to T5 (5% SP), indicating an increase in visceral mass relative to body weight ($P < 0.05$). The visceral fat-somatic indexes (VFSI%) displayed a significant rise from **Control** to T5 (5% SP), signifying an augmentation in visceral fat deposition over time ($P < 0.05$). The bile somatic index (BSI%) demonstrated a statistically significant increase from **Control** to T5 (5% SP), indicating an elevation in bile accumulation relative to body weight ($P < 0.05$).

Table 3. Mean (SE) value condition factor and body indices of *Osphronemus goramy* during the 90-day experiment after feeding trial of *Spirulina platensis*.

		Control (0% SP)	T1 (1% SP)	T2 (2% SP)	T3 (3% SP)	T4 (4% SP)	T5 (5% SP)
Hepatosomatic index (HSI) (%)		0.926±0.012 ^c	0.956±0.008 ^d	0.976±0.008 ^d	0.993±0.008 ^c	1.1±0.011 ^b	1.29±0.005 ^a
Viscerosomatic index (GSI%)		2.706±0.029 ^f	2.78±0.017 ^e	2.853±0.014 ^d	2.983±0.014 ^c	3.216±0.02 ^b	3.493±0.02 ^a
Visceral fat-somatic indexes (VFSI%)		2.09±0.005 ^f	2.133±0.008 ^e	2.17±0.005 ^d	2.21±0.005 ^c	2.29±0.005 ^b	2.41±0.005 ^a
Bilesomatic (BSI%)		9.916±0.014 ^f	10±0.025 ^e	10.206±0.008 ^d	10.296±0.014 ^c	10.343±0.008 ^b	10.443±0.008 ^a

3.3. Carcas composition

Protein content exhibited a significant increase ($P < 0.05$) as *spirulina* content increased, from 17.07% ± 0.03 at **Control** to 18.6% ± 0.01 at T5 (Table 7). Fat content showed a similar trend, with significant differences observed between each treatment ($P < 0.05$). Fiber content also increased significantly with higher SP content, ranging from 0.96% ± 0 at **Control** to 1.2% ± 0.01 at T5. ASH (Ash content) decreased significantly ($P < 0.05$) with increasing SP content, with values ranging from 5.13% ± 0.01 at **Control** to 4.23% ± 0.01 at T5. Carbohydrate (CHO) content displayed a significant decrease ($P < 0.05$) as SP content increased, with values ranging from 1.98% ± 0.01 at **Control** to 1.21% ± 0 at T5. Dry matter content showed significant variability across SP levels, and Moisture content exhibited an inverse relationship with *spirulina* content. Overall, the results indicate that increasing SP content in the body composition led to significant changes in nutritional composition, particularly in protein, fat, fibre, ash, and carbohydrate contents.

Table 7. Effects of dietary *S. platensis* on Body composition of *Osphronemus goramy*.

	Control (0% SP)	T1 (1% SP)	T2 (2% SP)	T3 (3% SP)	T4 (4% SP)	T5 (5% SP)
Crude Protein%	17.07±0.03 ^f	17.36±0.01 ^e	17.51±0.01 ^d	17.79±0.01 ^c	18.19±0 ^b	18.6±0.01 ^a
Crude Fat	4.2±0.01 ^d	4.29±0.01 ^c	4.28±0.01 ^c	4.51±0.02 ^b	4.53±0.02 ^b	4.61±0.01 ^a

Crude Fiber	0.96±0 ^d	0.98±0.01 ^d	1.04±0 ^c	1.08±0.01 ^b	1.11±0.01 ^b	1.2±0.01 ^a
ASH	5.13±0.01 ^f	4.91±0.01 ^e	4.81±0.01 ^d	4.51±0.01 ^c	4.3±0.01 ^b	4.23±0.01 ^a
CHO	1.98±0.01 ^a	1.96±0 ^a	1.95±0.01 ^a	1.97±0 ^a	1.52±0 ^b	1.21±0 ^c
Dry matter	29.35±0.06 ^d	29.52±0.02 ^c	29.61±0.04 ^c	29.87±0.03 ^a	29.66±0.02 ^b	29.87±0.05 ^a
Moisture content	70.64±0.06 ^a	70.48±0.02 ^b	70.39±0.04 ^{bc}	70.12±0.03 ^d	70.33±0.02 ^c	70.13±0.05 ^d

3.4. Hematological parameters

As indicated in Table 4, no significant differences were observed in RBC levels among all treatments ($P > 0.05$). However, WBC, Hb, HCT, Monocyte, and Neutrophil levels exhibited a significant increase from **Control** to T5 (5% SP) as *spirulina* concentration increased ($P < 0.05$). Lymphocyte levels demonstrated significant fluctuations across different concentrations of *spirulina*, with the highest level observed at T5 (5% SP) and the lowest in **Control** ($P < 0.05$).

Table4. Effects of dietary *S. platensis* on hematological parameters in *Osphronemus goramy*.

	Control (0% SP)	T1 (1% SP)	T2 (2% SP)	T3 (3% SP)	T4 (4% SP)	T5 (5% SP)
RBC (mm ³ . 10 ⁶)	3.216±1.608 ^a	4.933±0.008 ^a	4.996±0.008 ^a	5.123±0.012 ^a	5.263±0.008 ^a	5.413±0.008 ^a
WBC (mm ³ . 10 ⁴)	8.886±0.02 ^f	9.316±0.02 ^e	9.686±0.012 ^d	10.516±0.02 ^c	11.203±0.017 ^b	12.303±0.024 ^a
(Hb, g dL-1)	5.526±0.012 ^e	5.67±0.011 ^d	5.793±0.008 ^c	5.766±0.014 ^c	5.906±0.008 ^b	6±0.011 ^a
HCT (%)	33.513±0.008 ^f	33.813±0.008 ^e	34.1±0.005 ^d	34.423±0.008 ^c	34.846±0.008 ^b	35.216±0.008 ^a
Monocyte (%)	2.51±0.011 ^f	2.776±0.017 ^e	2.916±0.014 ^d	3.33±0.015 ^c	3.79±0.015 ^b	4.21±0.011 ^a
Lymphocyte (%)	76.233±0.12 ^a	73.066±0.12 ^d	75.166±0.088 ^b	72.733±0.088 ^d	74.333±0.12 ^c	70.733±0.12 ^e
Neutrophil (%)	10.116±0.012 ^f	11.423±0.014 ^e	12.663±0.008 ^d	12.983±0.017 ^c	13.65±0.015 ^b	14.306±0.008 ^a

3.5. Serum biochemical parameters

The serum biochemical parameters of fish fed with *Spirulina* are shown in Table 5. Fish fed the control diet exhibited the highest levels of cholesterol and triglyceride, with these levels decreasing as the dietary *S. platensis* levels increased ($P < 0.05$). Total protein (g dL-1), albumin (g/dL), and globulin (g/dL) levels increased with the increase in *spirulina* concentration. The significantly highest levels of protein (g dL-1), albumin (g/dL), and globulin (g/dL) were observed in the T5 (5% SP) diet group ($P < 0.05$). The A:G ratio (g/dL) was significantly highest in the **Control** (0% SP) diet group and lowest in the T1 (1% SP) diet group ($P < 0.05$).

Table5. The effects of dietary *S. platensis* (SP) on the blood biochemical parameters of *Osphronemus goramy*.

	Control (0% SP)	T1 (1% SP)	T2 (2% SP)	T3 (3% SP)	T4 (4% SP)	T5 (5% SP)
Total protein (g dL-1)	3.53±0.015 ^f	3.94±0.017 ^e	4.186±0.012 ^d	4.36±0.011 ^c	4.663±0.008 ^b	4.833±0.014 ^a
Albumin (g/dL)	1.843±0.012 ^f	1.91±0.011 ^e	1.97±0.005 ^d	2.093±0.008 ^c	2.153±0.008 ^b	2.27±0.011 ^a
Globulin(g/dL)	1.823±0.014 ^f	2.033±0.02 ^e	2.093±0.008 ^d	2.176±0.008 ^c	2.256±0.014 ^b	2.33±0.015 ^a
A:G ratio (g/dL)	1.011±0.003 ^a	0.939±0.015 ^c	0.941±0.006 ^c	0.961±0.001 ^{bc}	0.954±0.005 ^{bc}	0.974±0.007 ^c
Triglyceride (Mg/dl)	315.366±0.384 ^a	300.3±0.115 ^b	293.633±0.12 ^c	289.7±0.115 ^d	283.733±0.12 ^e	280.066±0.12 ^f
Cholesterol (mg/dl)	191.9±0.115 ^a	187.9±0.057 ^b	181.1±0.115 ^c	178±0.057 ^d	172.933±0.088 ^e	168.566±0.145 ^f

3.6. Intestinal digestive enzymes and Hepatic antioxidant activities

Table 6 illustrates the digestive enzymes activities and antioxidant activities of fish fed with *Spirulina*. Both Amylase and Lipase activities experienced a noteworthy increase as the concentration of *spirulina* increased from 0% to 5% ($P < 0.05$). These enzymatic activities demonstrated a gradual elevation with the increasing *spirulina* concentration in the diet. However, there was no significant variation observed in Protease activity across different concentrations of *spirulina* ($P > 0.05$). MDA levels showed a significant decrease from **Control** (0% SP) to T5 (5% SP) with the increasing concentration of *spirulina* ($P < 0.05$). Conversely, SOD levels exhibited a significant increase from **Control** (0% SP) to T5 (5% SP) with the increasing *spirulina* concentration ($P < 0.05$).

Table 6. The effects of dietary *S. platensis* (SP) on Intestinal digestive enzymes activities and Hepatic antioxidant activities of *Osphronemus goramy*.

	Control (0% SP)	T1(1% SP)	T2(2% SP)	T3(3% SP)	T4(4% SP)	T5(5% SP)
Amilase (U/mg substrate)	2.11±0.011 ^f	2.326±0.014 ^c	2.25±0.011 ^d	2.456±0.008 ^c	2.656±0.02 ^b	2.873±0.012 ^a
Lipase (U/mg substrate)	6.803±0.017 ^f	7.346±0.014 ^e	7.126±0.017 ^d	7.833±0.02 ^c	7.553±0.017 ^b	7.973±0.008 ^a
Protease (U/mg substrate)	0.019±0 ^c	0.022±0 ^b	0.019±0 ^c	0.02±0 ^c	0.023±0 ^b	0.025±0 ^a
MDA (µg/g)	2.02±0.017 ^a	1.96±0.011 ^b	1.906±0.008 ^c	1.836±0.014 ^d	1.496±0.017 ^e	1.356±0.012 ^f
SOD (Umg/prot)	6.406±0.008 ^f	7.25±0.011 ^e	7.73±0.011 ^d	8.03±0.011 ^c	8.36±0.011 ^b	8.576±0.008 ^a

4. DISCUSSION

4.1. Water quality

In The present study found that water quality parameters, including temperature (°C), dissolved oxygen (mg/L), total alkalinity (mg/L as CaCO₃), hardness (mg/L as CaCO₃), Ph and nitrates (mg/L), were all within acceptable limits according to WHO/FAO standards (Table 9).

Table 9. standard water quality parameter range according to WHO/FAO limits of fish culture

Water quality parameters	WHO/FAO limits	References
Water temperatures (°C)	25–33	(108)
Dissolved oxygen (mg/L)	3–5	(109)
Total alkalinity (mg/L as CaCO ₃)	120	(110)
Hardness (mg/L as CaCO ₃)	168	(110)
pH	6.5–9.0	(110)
Nitrates (mg/L)	0.2–219	(110)

4.2. Growth performance

During feeding trial, observed a significant increase in the growth performance of *O. goramy* when the *Spirulina* concentration in the feed increased from 1% to 5% over a span of 90 days. The inclusion of *Spirulina* in aquafeed led to an increase in hunger and notably elevated feed intake, resulting in improvements in parameters such as Final weight (FW), weight gain (WG), weight gain% (WG%), feed efficiency ratio (FER), average daily growth (ADG), and specific growth rate (SGR). These findings are consistent with previous research (29,36,59,60). Studies by Hassanpour et al. (61) and Zeinab et al. (62) found that the supplementation of *Spirulina* in the diets of rainbow trout and Nile Tilapia, respectively, resulted in higher body weights compared to control groups. Similarly, research by Jana et al. (37), Simanjuntak et al. (13), and Abu-Elala et al. (63) demonstrated significant growth enhancement in fish fed with *Spirulina*, without any adverse effects (64) and with potential as a feed material (38). Additionally, the inclusion of *Spirulina* in aquafeed improved survival, growth rate, and feed utilization in *Macrobrachium rosenbergii* prawn culture at concentration level from 5% to 20%, as observed in similar studies on *Litopenaeus vannamei* (65). Overall, *Spirulina* shows promise as a natural supplement for enhancing growth and health in aquaculture.

S. platensis enhances growth in fish due to its highly digestible protein content, abundant essential fatty acids, vitamins (such as vitamin B12), and minerals (66,67). Its cellular structure lacks cellulose, making it easily digestible by fish digestive enzymes, aided by the presence of mucopolymer murein (64). Additionally, *Spirulina* promotes the growth of beneficial intestinal flora in fish and aids in the breakdown of indigestible feed components (36,64). Parameters like protein efficiency ratio (PER), survival rate (SR%), and FCR significantly improve with increased *Spirulina* concentration.

4.3. Body indices

At the end of experiment result was found that the body indices (HIS%, VFSI%, GSI% and BSI%) was increased with increase in concentration of *spirulina* in diet from (0% to 5%). Significant highest Hepatosomatic index (HSI) (%), Viscerosomatic index (GSI%), Visceral fat-somatic indexes (VFSI%) and Bilesomatic (BSI%) observed in T5 (5% SP) diet ($P < 0.05$). A higher hepatosomatic index (HSI) reflects increased glycogen levels and favorable environmental conditions. This relationship was observed in separate studies conducted by Velasquez et al. (68) on Nile tilapia and Tulli et al. (69) on European sea bass juveniles (*Dicentrarchus labrax*), where diets containing 20%

Tetraselmis suecica microalgae resulted in lower HSI compared to control diets without microalgae. This connection between the level of microalgae inclusion and improved lipid mobilization efficiency in fish liver has also been noted by (70) and (71). The viscerosomatic index (VSI) serves as an indicator of fish digestive capacity. Consistent findings were reported by (68) in Nile tilapia, where the highest VSI was observed in fish fed diets with 0% to 30% microalgae inclusion, showing a downward trend as the level of *Arthrospira* inclusion increased. Nevertheless, it's essential to recognize that VSI represents just one aspect of fish digestive capacity, as other factors likely play a role.

4.4. Body composition

In this study, findings revealed that protein deposition in the body increased with higher *spirulina* content, notably peaking at $18.6 \pm 0.01a$ in the T5 group (5% SP) ($P > 0.05$). Likewise, lipid ($4.61 \pm 0.01a$) and fibre ($1.2 \pm 0.01a$) content rose with escalating *spirulina* concentration, while carbohydrate (CHO) ($1.21 \pm 0c$) and ash (ASH) ($4.23 \pm 0.01a$) levels declined in *O.goramy* bodies with increasing *spirulina* content. Overall, the body composition exhibited optimal results in the high 5% (*spirulina*) concentration diet, indicating efficient utilization of *spirulina*-infused feed by *O.goramy* bodies. These results are consistent with past research; Moreira et al. (105) proposed that *spirulina* LEB-18 is a valuable dietary addition for addressing nutritional deficiencies due to its rich mineral and nutritional content. Changes in body fat and protein content are linked to variations in synthesis and deposition levels within muscle tissues (29,106). (107) noted significant enhancements in the biochemical composition of *Macrobrachium rosenbergii* PL bodies when fed *Spirulina*-enriched *Artemia* nauplii. Feeding larvae of *C.carpio* with 20% *Spirulina* led to considerable improvements in body composition, particularly in protein composition and body fat (59).

4.5. Hematological parameters

In present study, we observed a dose-dependent increase in blood parameters, including, WBC, HCT, Hb, Monocyte (%) and Neutrophil (%), in fish fed with 5% *S. platensis* compared to the control group. But there is no significant difference was found in the RBC count among fish fed different concentrations of *Spirulina* ($P > 0.05$) and Lymphocyte (%), count is significantly decrease with increase in the concentration of *spirulina* ($P < 0.05$). the RBC count Similar findings were reported by (72) and (34). Promya and Chitmant(73) demonstrated higher WBC and RBC counts in African sharp tooth catfish fed diets containing 5% *A. platensis* + basal. (44) found increased RBC and Hb levels in Great Sturgeon fed with *S. platensis*. *Spirulina*'s high iron content significantly affects erythropoiesis, as seen in studies on anaemic rats (74). (75) observed the highest levels of RBC, WBC, and Hb in Nile tilapia with 10% *Spirulina* supplementation. Similar increases in WBC were reported in large yellow croaker and carp with *Spirulina* supplementation(54,74). This finding was also supported by (76) and (77). This is attributed to polypeptide-phycoerythrin, enhancing WBC production (78). (62) noted increased hemoglobin levels in Nile tilapia fed *Spirulina*, while (36) found 10% *S. platensis* supplementation increased hematocrit values. However, carp RBCs were unaffected by 3.0–5.0 g/kg *Spirulina*(79).

4.6. Blood biochemical parameters

In fish, the levels of total protein, albumin, and globulin in serum provide valuable diagnostic insights into overall nutritional status, vascular system integrity, and liver function (80). The current study observed that supplementation with 5% *spirulina* resulted in higher levels of total protein, albumin, and globulin compared to the control group. Conversely, previous research by (13) found that diets containing 2.0–6.0% *Spirulina* significantly increased serum levels of total protein, globulin, and the A/G ratio. Similar findings were reported by (81), where feeding with 10.0% *Spirulina* significantly elevated serum albumin levels, as also observed by (29). Additionally, (62) noted that increased levels of *S. platensis* correlated with higher serum total protein and globulin, although there was no significant difference in albumin levels ($P > 0.05$). The rise in serum albumin levels could be attributed to *Spirulina*'s various active compounds, including carotenoids, polysaccharides, vitamins, minerals, and linoleic acid, which act as immunostimulants, enhancing the immune system (82). *Spirulina*'s potent antioxidant properties contribute to its ability to improve immune capacity and health status in *Osphronemus gourami*, similar to findings in *O. niloticus*(22,63). Furthermore, (83) demonstrated that different levels of *S. platensis* improved total serum protein, albumin, globulin, and biochemical responses in *Mugil cephalus*. Including *S. platensis* and *Sargassum ilicifolium* extract in diets for *Huso huso* and *O. niloticus* also increased serum total protein and globulin levels (62,84). The serum albumin level plays a crucial role in regulating metabolism, maintaining osmotic pressure in the blood, and facilitating the transport of both exogenous and endogenous metabolites (85). Globulin serves as a prominent carrier/transporter in living organisms Simanjuntak et al.(13).

In the present result cholesterol and triglyceride level decrease with increase in *Spirulina* concentration. Similarly, (86) discovered that incorporating 1.0 g/kg *Spirulina* into tilapia diets did not impact the serum levels of total cholesterol and triglycerides. However, diets containing 10.0% *Spirulina* significantly raised the serum levels of cholesterol and triglycerides(29). *S. platensis* contains α -linolenic acid, linoleic acid, and β -carotene, which have the potential to lower cholesterol and triglyceride levels and prevent hypercholesterolemia and atherosclerosis (87,88).

4.7. Digestive enzyme

Algae are abundant in both macro and micro elements, which have the potential to stimulate digestive enzyme activity (89). Low concentrations of microalgae in the gut may trigger the secretion of endogenous digestive enzymes and promote fish growth(90,91). (92) noted that *Isochrysis galbana* algae enhanced pancreatic and intestinal enzyme production in sea bass (*Dicentrarchus labrax*) larvae. While fish typically struggle to digest plant matter due to the cellulose cell wall, *S. platensis* is known to have a thin cell wall composed primarily of 80% pectin and 20% cellulose, facilitating improved digestion and assimilation (93). Consequently, *S. platensis* is easily digested and may enhance the assimilation of other nutrients, including antioxidants.

In the current experiment, the activity of amylase (U/mg substrate), lipase (U/mg substrate), and protease (U/mg substrate) significantly increased with increase the concentration of *spirulina* in diet compared to the control group ($P>0.05$). Similarly, Biabani Asrami et al. observed that dietary supplementation of phycocyanin extracted from *Spirulina* enhanced the activities of amylase, protease, and lipase in Guppy fish (*Poecilia reticulata*(27)). Additionally, Teimouri et al. (94) found that *S. platensis* enhanced digestive enzyme activity, thereby improving nutrient assimilation. Therefore, increasing the inclusion of *spirulina* in the diet improved the digestive enzyme activity in *Osphronemus gourami*.

4.8. Antioxidant activity

Superoxide dismutase (SOD) play crucial roles in scavenging free radicals and protecting essential cellular macromolecules and organelles from oxidative damage (95). Previous studies have indicated that dietary supplementation of *spirulina* can reduce antioxidant activities in Nile tilapia (*Oreochromis niloticus*)(96) and goldfish (*Carassius auratus*) (97). Additionally, Li et al. (2009) demonstrated that phycocyanin enhances the activities of SOD enzymes in pancreatic beta cells. Tayag et al. (98) and Lin et al. (99) reported a direct increase in SOD enzyme activity with *spirulina* extract in white shrimp. Abdelkhalek et al. (100) illustrated that *spirulina* supplementation enhances SOD enzyme activity in *Oreochromis niloticus*. Furthermore, Yu et al.(75) observed that rainbow trout fed with 10% *spirulina* exhibited the highest SOD activity.

MDA serves as the end product of lipid peroxidation, leading to cellular toxicity and accelerating damage to cells and tissues(101–103). In the current study, MDA enzyme activity decreased with an increase in *spirulina* concentration, consistent with previous findings in Nile tilapia (*O. niloticus*) (96), and Carp (*Cyprinus carpio var*) (104). Similarly, Abdelkhalek et al.(100) reported that dietary supplementation with *S. platensis* reduced MDA levels in the liver, kidney, and gills of Nile tilapia.

5. CONCLUSIONS

In conclusion, recent experiment showed that *Osphronemus gourami* fed a diet supplemented with *S. platensis* at 5% for 90 days improved fish growth performance (FW, WG, WG%, PER, FCR, FER, ADG, SGR, and SR%), increased feed and protein utilization, improve the body deposition (Protein, fat, ASH, CHO, fiber), elevated body indices (HIS%, VFSI%, GSI%, and BSI%), enhanced hematological parameters (RBC, WBC, HCT, Hb, Monocyte %, and Neutrophil %), improved biochemical responses (Total serum protein, Albumin, and Globulin), decreased levels of cholesterol and triglyceride, and increased digestive (Amylase, Lipase, and Protease) and antioxidant (SOD & MDA) enzyme activity in fish and improve the immunity.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. For writing the review of literature and few information of introduction the Chat GTP is used.
2. The Grammarly is used to check the grammar and scientific writing.

REFERENCES

1. FAO. (Food and Agriculture Organization). 2008 The state of world fisheries and aquaculture: Opportunities and challenges. Food and Agriculture Organization of the United Nations. 2018;209.
2. Hua K, Cobcroft JM, Cole A, Condon K, Jerry DR, Mangott A, et al. The future of aquatic protein: implications for protein sources in aquaculture diets. *One Earth*. 2019;1(3):316–29.
3. Gómez B, Munekata PE, Zhu Z, Barba FJ, Toldrá F, Putnik P, et al. Challenges and opportunities regarding the use of alternative protein sources: Aquaculture and insects. *Adv Food Nutr Res*. 2019;89:259–95.
4. Zhang Y, Lu R, Qin C, Nie G. Precision nutritional regulation and aquaculture. *Aquac Rep*. 2020;18:100496.
5. Nagappan S, Das P, AbdulQuadir M, Thaher M, Khan S, Mahata C, et al. Potential of microalgae as a sustainable feed ingredient for aquaculture. *J Biotechnol*. 2021;341:1–20.
6. Dawood MA, Koshio S. Application of fermentation strategy in aquafeed for sustainable aquaculture. *Rev Aquac*. 2020;12(2):987–1002.
7. Azrita HS, Syandri H. Morphological character among five strains of giant gourami, *Oshpronemusgouramy* Lacepede, 1801 (Actinopterygii: Perciformes: Osphronemidae) using a truss morphometric system. *Int J Fish Aquat Stud* [Internet]. 2015;2(6):344–50. Available from: <https://www.fisheriesjournal.com/vol2issue6/Pdf/2-6-58.1.pdf>
8. Sundara RB. Notes on the freshwater fishes of Madras. *Records Indian Museum* 7; 1916.
9. Caruso D, Arifin ZO, Subagja J, Slembrouck J, New M. *Osphronemusgoramy*. Cultured Aquatic Species Information Programmed. Rome: FAO Fisheries and Aquaculture Department; 2019.
10. Amornsakun T, Kullai S, Hassan A. Feeding behavior of giant gourami, *Osphronemusgoramy* (Lacepede) larvae. *Songklanakarin Journal of Science and Technology* [Internet]. 2014;36(3):261–4. Available from: <https://www.thaiscience.info/Journals/Article/SONG/10968306.pdf>

11. Arifin OZ, Imron I, Asependi A, Hendri A, Muslim N, Yani A. Hibridisasi intraspesifik antar dua populasi ikan gurami Galunggung (*Osphronemus goramy*, Lacepede, 1801). *Jurnal Riset Akuakultur*. 2018;12(4):315–23.
12. Prakoso VA, Pouil S, Prabowo MNI, Sundari S, Arifin OZ, Subagja J, et al. Effects of temperature on the zootechnical performances and physiology of giant gourami (*Osphronemus goramy*) larvae. *Aquaculture*. 2019;510:160–8.
13. Simanjuntak SBI, Indarmawan I, Wibowo ES. Impact of fed containing different levels of diets supplementation *Spirulina platensis* on growth, haematological, body composition and biochemical parameters, of Gurami (*Osphronemus goramy*). *Turk J Fish Aquat Sci*. 2018;18(5):681–90.
14. FAO. Cultured aquatic species information programme *Osphronemus goramy* (Lacépède, 1801). In: FAO Fisheries and Aquaculture Department [Internet]. 2020. Available from: <http://www.FAO.org/fishery/culturedspecies/Osphr>
15. RI W. International introduction of Inland Aauatic species. Rome, Fao Fisheries Technical Papers. 1988;(294):318.
16. NFDB. Matsya Bharat. National Fisheries Development Board, Government of India. 2016;8.
17. Kumari DJ, Babitha B, Jaffar S, Prasad MG, Ibrahim MD, Khan MS. Potential health benefits of *Spirulina platensis*. *Int J Adv Pharm Sci* [Internet]. 2011;2:417–22. Available from: https://www.researchgate.net/profile/Mohammad-Ibrahim-38/publication/215961590_Potential_Health_Benefits_of_Spirulina_platensis/links/06ab3257153eddb43a4d41ff/Potential-Health-Benefits-of-Spirulina-platensis.pdf
18. Henrikson R. *Spirulina*: Health discoveries from the source of life. 2000.
19. Allen D. Nutrient comparison of *Spirulina* with carrot. Denver, Colorado. webmaster@microhydron.com. 2000.
20. Belay A. The potential application of *Spirulina* (*Arthrospira*) as a nutritional and therapeutic supplement in health management. *Journal of the american nutraceutical association (jana)* [Internet]. 2002;5(-),. Sid:26–48. Available from: <https://sid.ir/paper/546615/en>
21. Gutiérrez-Salmeán G, Fabila-Castillo L, Chamorro-Cevallos G. Aspectos nutricionales y toxicológicos de *Spirulina* (*arthrospira*). *Nutr Hosp*. 2015;32(1):34–40.
22. Bermejo P, Piñero E, Villar ÁM. Iron-chelating ability and antioxidant properties of phycocyanin isolated from a protean extract of *Spirulina platensis*. *Food Chem* [Internet]. 2008;110(2):436–45. Available from: <http://dx.doi>
23. Phang SM, Miah MS, Yeoh BG, Hashim MA. *Spirulina* cultivation in digested sago starch factory wastewater. *J Appl Phycol*. 2000;12:395–400.
24. Jaime-Ceballos BJ, Hernández-Llamas A, Garcia-Galano T, Villarreal H. Substitution of *Chaetoceros muelleri* by *Spirulina platensis* meal in diets for *Litopenaeus schmitti* larvae. *Aquaculture*. 2006;260(1–4):215–20.
25. Kamble SP, Gaikar RB, Padalia RB, Shinde KD. Extraction and purification of C-phycocyanin from dry *Spirulina* powder and evaluating its antioxidant, anticoagulation and prevention of DNA damage activity. *J Appl Pharm Sci*. 2013;3(8):149–53.

26. Bhat VB, Madyastha KM. Scavenging of peroxynitrite by phycocyanin and phycocyanobilin from *Spirulina platensis*: protection against oxidative damage to DNA. *BiochemBiophys Res Commun*. 2001;285(2):262–6.
27. BiabaniAsrami M, Sudagar M, Shahraki N, Vahdat S. Effect of extracted phycocyanin from *Spirulina platensis* on growth parameters, colorations, digestive enzymes and body chemical compositions of Guppy fish (*Poecilia reticulata*). *Survey in Fisheries Sciences* [Internet]. 2019;6(1):1–8. Available from: https://web.archive.org/web/20200715221920id_/http://sifisheriessciences.com/files/site1/user_files_d60aad/fallahi-A-10-444-56-b3344d0.pdf
28. Fernández-Rojas B, Medina-Campos ON, Hernández-Pando R, Negrette-Guzmán M, Huerta-Yepez S, Pedraza-Chaverri J. C-phycocyanin prevents cisplatin-induced nephrotoxicity through inhibition of oxidative stress. *Food Funct*. 2014;5(3):480–90.
29. Abdel-Tawab M, Ahmed H. Live *Spirulina* (*Arthrospira platensis*) as a growth and immunity promoter for Nile tilapia, *Oreochromis niloticus* (L.), challenged with pathogenic *Aeromonas hydrophila*. *Aquac Res* [Internet]. 2009;40(9):1037–46. Available from: <http://dx.doi>
30. Awad LZ, El-Mahallawy HS, Abdelnaeim NS, Mahmoud MM, Dessouki AA, ElBanna NI. Role of dietary *Spirulina platensis* and betaine supplementation on growth, hematological, serum biochemical parameters, antioxidant status, immune responses, and disease resistance in Nile tilapia. *Fish Shellfish Immunol*. 2022;126:122–30.
31. Mahmoud MM, El-Lamie MM, Kilany OE, Dessouki AA. *Spirulina* (*Arthrospira platensis*) supplementation improves growth performance, feed utilization, immune response, and relieves oxidative stress in Nile tilapia (*Oreochromis niloticus*) challenged with *Pseudomonas fluorescens*. *Fish Shellfish Immunol*. 2018;72:291–300.
32. Li XL, Xu G, Chen T, Wong YS, Zhao HL, Fan RR, et al. Phycocyanin protects INS-1E pancreatic beta cells against human islet amyloid polypeptide-induced apoptosis through attenuating oxidative stress and modulating JNK and p38 mitogen-activated protein kinase pathways. *Int J Biochem Cell Biol*. 2009;41(7):1526–35.
33. Rabadiya B, Patel P. *Spirulina: Potential* clinical therapeutic application. *J of Pharmacy Research*. 2010;3(8):1726–32.
34. El-Sheekh M, El-Shourbagy I, Shalaby S, Hosny S. Effect of feeding *Arthrospira platensis* (*Spirulina*) on growth and carcass composition of hybrid red tilapia (*Oreochromis niloticus* x *Oreochromis mossambicus*). *Turk J Fish Aquat Sci*. 2014;14(2):471–8.
35. Kim SS, Rahimnejad S, Kim KW, Lee KJ. Partial replacement of fish meal with *Spirulina pacifica* in diets for parrot fish (*Oplegnathus fasciatus*). *Turk J Fish Aquat Sci*. 2013;13(2).
36. Ibrahim MD, Mohamed MF, Ibrahim MA. The role of *Spirulina platensis* (*Arthrospira platensis*) in growth and immunity of Nile tilapia (*Oreochromis niloticus*) and its resistance to bacterial infection. 5(6). 2013;109–17.
37. Jana A, Saroch JD, Borana K. Effect of *Spirulina* as a feed supplement on survival and growth of *Pangasius sutchi*. *Int J Fish Aquat Stud* [Internet]. 2014;1(5):77–9. Available from: <https://www.fisheriesjournal.com/vol1issue5/pdf/98.1.pdf>

38. Simanjuntak SBI, Soedibya PHT, Wibowo ES. Growth performance of gurami (*Osphronemus gouramy* Lac.) the given phytoplankton *Spirulina platensis* and *Chlorella vulgaris*. 2014.
39. Aryani N, Mardiah A, Syandri H. Influence of feeding rate on the growth, feed efficiency and carcass composition of the Giant gourami (*Osphronemus goramy*). Pak J Zool [Internet]. 2017;49(5). Available from: https://www.researchgate.net/profile/Ainul-Mardiah-3/publication/319604781_Influence_of_Feeding_Rate_on_the_Growth_Feed_Efficiency_and_Carcass_Composition_of_the_Giant_Gourami_Osphronemus_goramy/links/5ab5caf745851515f59a8124/Influence-of-Feeding-Rate-on-the-Growth-Feed-Efficiency-and-Carcass-Composition-of-the-Giant-Gourami-Osphronemus-goramy.pdf
40. Budi DS, Alimuddin A, Suprayudi MA. Growth response and feed utilization of giant gourami (*Osphronemus goramy*) juvenile feeding different protein levels of the diets supplemented with recombinant growth hormone. Hayati. 2015;22(1):12–9.
41. Nofyan E. Pengaruh Pemberian Pakan Dari Sumber Nabati Dan Hewani Terhadap Berbagai Aspek Fisiologi Ikan Gurami (*Osphronemus Gouramy* L.) [The Effect of Plant and Animal Food on Various Physiological Aspects of Giant Gourami (*Osphronemus Gouramy* L.)]. J Iktiologi Indones [Internet]. 2005;5(1):19–23. Available from: <https://doi.org/10.32491/jii.v5i1.297>
42. Yandes Z, Affandi R. Pengaruh pemberian selulosadalam pakan terhadap kondisi biologis benih ikan gurami (*Osphronemus gouramy* Lac) [Effect of cellulose in dietary on the biological condition of giant gourami fry (*Osphronemus goramy* Lac)]. J Iktiologi Indones [Internet]. 2003;3(1):27–33. Available from: <https://doi.org/10.32491/jii.v3i1.269>
43. Nutrient N. requirements of fish [Internet]. Washington DC: National Academy Press; 1993. Available from: [https://books.google.co.in/books?hl=en&lr=&id=M2aBtOUhRagC&oi=fnd&pg=PA1&dq=NRC+Nutrient+requirements+of+fish.++\(1993\).+Washington+DC:+National+Academy+Press.&ots=ZXhiJPPIso&sig=ZyWycMmB4zMySw5YLTh1YqYXfgE&redir_esc=y#v=onepage&q=NRC](https://books.google.co.in/books?hl=en&lr=&id=M2aBtOUhRagC&oi=fnd&pg=PA1&dq=NRC+Nutrient+requirements+of+fish.++(1993).+Washington+DC:+National+Academy+Press.&ots=ZXhiJPPIso&sig=ZyWycMmB4zMySw5YLTh1YqYXfgE&redir_esc=y#v=onepage&q=NRC)
44. Adel M, Yeganeh S, Dadar M, Sakai M, Dawood MA. Effects of dietary *Spirulina platensis* on growth performance, humoral and mucosal immune responses and disease resistance in juvenile great sturgeon (*Huso huso* Linnaeus, 1754). Fish Shellfish Immunol. 2016;56:436–44.
45. Castell JD, Tiews K. Report of the EIFAC, IUNS and ICES Working Group on Standardization of Methodology in Fish Nutrition Research, Hamburg, Federal Republic of Germany, 21-23 March 1979 (No [Internet]. 1980. Available from: <https://agris.fao.org/search/en/providers/122621/records/64774bfaf2e6fe92b36378d3>
46. Stuart JS, Hung SS. Growth of juvenile white sturgeon (*Acipenser transmontanus*) fed different proteins. Aquaculture. 1989;76(3–4):303–16.
47. AOAC. Official methods of analysis. In: W Horwitz (Ed). 13th ed. Washington, DC: Academic Press; 2000.
48. Nishikimi M, Rao NA, Yagi K. The occurrence of superoxide anion in the reaction of reduced phenazine methosulfate and molecular oxygen. Biochem Biophys Res Commun. 1972;46(2):849–54.

49. Mihara M, Uchiyama M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid test. *Anal Biochem.* 1978;86(1):271–8.
50. Ag G. Determination of serum proteins by means of the biuret reactions. *J Biol Chem*, [Internet]. 1949;177:51. Available from: https://cir.nii.ac.jp/crid/1573387450479762048#citations_container
51. Doumas BT, Watson WA, Biggs HG. Albumin standards and the measurement of serum albumin with bromocresol green. *Clinicachimica acta.* 1997;258(1):21–30.
52. Allain CC, Poon LS, Chan CS, Richmond WFPC, Fu PC. Enzymatic determination of total serum cholesterol. *Clin Chem.* 1974;20(4):470–5.
53. McGowan MW, Artiss JD, Strandbergh DR, Zak B. A peroxidase-coupled method for the colorimetric determination of serum triglycerides. *Clin Chem.* 1983;29(3):538–42.
54. Li M, Wu W, Zhou P, Xie F, Zhou Q, Mai K. Comparison effect of dietary astaxanthin and *Haematococcus pluvialis* on growth performance, antioxidant status and immune response of large yellow croaker *Pseudosciaenacrocea*. *Aquaculture.* 2014;434:227–32.
55. Bergmeyer HU, Grassi M. Methods of enzymatic analysis [Internet]. 1983. Available from: [https://books.google.co.in/books?id=GDd2zYuLpRwC&dq=Bergmeyer,+H.U.,+Grassi,+M.+\(1983\).+Methods+of+enzymatic+analysis.+Vol.+2.+Weinheim:+V+erlag+Chemie.&lr=&source=gbs_navlinks_s](https://books.google.co.in/books?id=GDd2zYuLpRwC&dq=Bergmeyer,+H.U.,+Grassi,+M.+(1983).+Methods+of+enzymatic+analysis.+Vol.+2.+Weinheim:+V+erlag+Chemie.&lr=&source=gbs_navlinks_s)
56. Worthington V. Worthington enzyme manual: enzymes and related biochemicals. Worthington Chemical, New Jersey. 1993;399.
57. Borlongan IG. Studies on the digestive lipases of milkfish, *Chanoschanos*. *Aquaculture.* 1990;89(3–4):315–25.
58. APHA. Standard Methods for Examination of Water and Wastewater. 19th ed. Washington DC, USA: American Public Health Association; 1995.
59. Abdulrahman NM, Ameen HJH. Replacement of fishmeal with microalgae *Spirulina* on common carp weight gain, meat and sensitive composition and survival. *Pakistan Journal of Nutrition.* 2014;13(2):93.
60. Abdulrahman NM. Evaluation of *Spirulina spp.* as food supplement and its effect on growth performance of common carp fingerlings. *Int J Fish Aquat Stud* [Internet]. 2014;2(2):89–92. Available from: <https://www.fisheriesjournal.com/archives/2014/vol2issue2/PartB/2.pdf>
61. Hassanpour S, Chaichisemsari M, Riyazi SR. Effect of replacing different levels of Soybean meal with *Spirulina* on performance in Rainbow Trout. *Ann Biol Res* [Internet]. 2011;2(3):374–9. Available from: <https://d1wqtxts1xzle7.cloudfront.net/107724245/effect-of-replacing-different-levels-of-soybean-meal-with-spirulina-on-performance-in-rainbow-trout-libre.pdf?1700756584=&response-content-disposition=inline>
62. Zeinab AK, Aly MS, Faiza AK, Fatma EM. Effect of *Spirulina platensis* and *Lactobacillus rhamnosus* on growth and biochemical performance of Nile Tilapia (*Oreochromis niloticus*) fingerlings. *Int J Curr Microbiol Appl Sci* [Internet]. 2015;4(4):747–63. Available from: <http://www.ijcmas.com/vol-4-4/Zeinab>
63. Abu-Elala NM, Galal MK, Abd-Elsalam RM, Mohey-Elsaeed O, Ragaa NM. Effects of dietary supplementation of *Spirulina platensis* and garlic on the

growth performance and expression levels of immune-related genes in Nile tilapia (*Oreochromis niloticus*). *J Aquac Res Dev* [Internet]. 2016;7(7):433–42. Available from: https://www.researchgate.net/profile/Nermeen-Abu-Elala/publication/306047814_Effects_of_Dietary_Supplementation_of_Spirulina_platensis_and_Garlic_on_the_Growth_Performance_and_Expression_Level_s_of_Immune-related_Genes_in_Nile_tilapia_Oreochromis_niloticus/links/5f439eec458515b729499b83/Effects-of-Dietary-Supplementation-of-Spirulina-platensis-and-Garlic-on-the-Growth-Performance-and-Expression-Levels-of-Immune-related-Genes-in-Nile-tilapia-Oreochromis-niloticus.pdf

64. Teimouri M, Amirkolaie AK, Yeganeh S. The effects of *Spirulina platensis* meal as a feed supplement on growth performance and pigmentation of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*. 2013;396:14–9.
65. Hanel H, Broekman D, deGraaf S, Schnack D. S, Schnack D. Partial replacement of fishmeal by the lyophilized powder of the microalgae *Spirulina platensis* in Pacific white shrimp diets” *Open Mar Biol J*. 2007;1:1–5.
66. Wang L, Pan B, Sheng J, Xu J, Hu Q. Antioxidant activity of *Spirulina platensis* extracts by supercritical carbon dioxide extraction. *Food Chem*. 2007;105(1):36–41.
67. Saranraj P, Sivasakthi S. *Spirulina platensis*—food for future: a review. *Asian J Pharm Sci Technol* [Internet]. 2014;4(1):26–33. Available from: <https://d1wqtxts1xzle7.cloudfront.net/82407660/26-33-libre.pdf?1647806087=&response-content-disposition=inline>
68. Velasquez SF, Chan MA, Abisado RG, Traifalgar RFM, Tayamen MM, Maliwat GCF, et al. Dietary *Spirulina (Arthrospira platensis)* replacement enhances performance of juvenile Nile tilapia (*Oreochromis niloticus*). *J Appl Phycol*. 2016;28:1023–30.
69. Tulli F, Chini Zittelli G, Giorgi G, Poli BM, Tibaldi E, Tredici MR. Effect of the inclusion of dried *Tetraselmis suecica* on growth, feed utilization, and fillet composition of European sea bass juveniles fed organic diets. *Journal of aquatic food product technology*. 2012;21(3):188–97.
70. Nakagawa H. Effect of dietary algae on improvement of lipid metabolism in fish. *Biomedicine & pharmacotherapy*. 1997;51(8):345–8.
71. Ergün S, Soyutürk M, Güroy B, Güroy D, Merrifield D. Influence of *Ulva* meal on growth, feed utilization, and body composition of juvenile Nile tilapia (*Oreochromis niloticus*) at two levels of dietary lipid. *Aquaculture International*. 2009;17:355–61.
72. El Gammal MA, Shalaby AM, Ramadan AA. The use of *spirulina platensis* (microalgae) as a food additive to stimulate growth and immunity modulation for Nile tilapia (*Oreochromis niloticus*) challenged with pathogenic bacteria (*Pseudomonas fluorescens*). *Abbassa International Journal for Aquaculture* [Internet]. 2010;3(1):53–83. Available from: <https://abbassajournal.com/magazine/wp-content/uploads/2017/07/THE-USE-OF-SPIRULINA-PLATENSIS-MICROALGAE-AS-FOOD-ADDITIVES-TO-STIMULATE-GROWTH-AND-IMMUNITYMODULATION-FOR-NILE-TILAPIA.pdf>
73. Promya J, Chitmanat C. The effects of *Spirulina platensis* and *Cladophora* algae on the growth performance, meat quality and immunity stimulating capacity of the African sharp-toothed catfish (*Clarias gariepinus*). *Int J Agric Biol* [Internet]. 2011;13:77–82. Available from:

https://www.researchgate.net/profile/Chitmanat-Chanagun/publication/279600728_The_effects_of_Spirulina_platensis_and_Cladophora_Algae_on_the_Growth_Performance_Meat_Quality_and_Immunity_Stimulating_Capacity_of_the_African_Sharptooth_Catfish_Clarias_gariepinus/links/641db79392cfd54f8428906c/The-effects-of-Spirulina-platensis-and-Cladophora-Algae-on-the-Growth-Performance-Meat-Quality-and-Immunity-Stimulating-Capacity-of-the-African-Sharptooth-Catfish-Clarias-gariepinus.pdf?_sg

74. Khalil SR, Reda RM, Awad A. Efficacy of *Spirulina platensis* diet supplements on disease resistance and immune-related gene expression in *Cyprinus carpio* L. exposed to herbicide atrazine. *Fish Shellfish Immunol.* 2017;67:119–28.
75. Yu W, Wen G, Lin H, Yang Y, Huang X, Zhou C, et al. Effects of dietary *Spirulina platensis* on growth performance, hematological and serum biochemical parameters, hepatic antioxidant status, immune responses and disease resistance of Coral trout *Plectropomus leopardus* (Lacepede, 1802). *Fish Shellfish Immunol.* 2018;74:649–55.
76. Hegazi SM, Fadl SE, Gad DM, Abeerd EKM. Effect of *Spirulina* incorporated in diets of tilapia nilotica fresh water fish exposed to lead toxicity. *J Vet Sci [Internet].* 2014;1:633–49. Available from: https://www.researchgate.net/profile/EI-Keredy-Abeer/publication/350008720_Effect_of_spirulina_incorporated_in_diets_of_Tilapia_nilotica_fresh_water_fish_exposed_to_lead_toxicity/links/604b5def299bf1f5d84171ad/Effect-of-spirulina-incorporated-in-diets-of-Tilapia-nilotica-fresh-water-fish-exposed-to-lead-toxicity.pdf
77. Sayed AEDH, Fawzy MA. Effect of dietary supplementation of *Spirulina platensis* on the growth and haematology of the catfish *Clarias gariepinus*. *J Adv Biol [Internet].* 2014;5(2):625–35. Available from: https://d1wqtxts1xzle7.cloudfront.net/35169553/Sayed_and_Fawzy_2014a-libre.pdf?1413550140=&response-content-disposition=inline
78. Zhang C. The effects of polysaccharide and phycocyanin from *Spirulina platensis* variety on peripheral blood and hematopoietic system of bone marrow in mice. In: *Second Asia-Pacific Conference on Alga Biotechnology.* 1994. p. 25–7.
79. Abdulrahman NM, Hama Ameen HJ, Hama SR, Hassan BR, Nader PJ. Effect of microalgae *Spirulina spp.* as food additive on some biological and blood parameters of common carp *Cyprinus carpio* L. *Iraqi Journal of Veterinary Sciences [Internet].* 2019;33(1):27–31. Available from: <https://www.iasj.net/iasj/download/15342ccc1690df08>
80. Schäperclaus W, Kulow H, Schreckenbach K. *Fish Disease—AA Balkema.* the Netherlands: Rotterdam; 1992.
81. Youssef IM, Saleh ES, Tawfeek SS, Abdel-Fadeel AA, Abdel-Razik ARH, Abdel-Daim AS. Effect of *Spirulina platensis* on growth, hematological, biochemical, and immunological parameters of Nile tilapia (*Oreochromis niloticus*). *Trop Anim Health Prod.* 2023;55(4):275.
82. Abdel-Daim MM, Dawood MA, Elbadawy M, Aleya L, Alkahtani S. *Spirulina platensis* reduced oxidative damage induced by chlorpyrifos toxicity in Nile tilapia (*Oreochromis niloticus*). *Animals.* 2020;10(3):473.
83. Akbary P, Molazaei E, Aminikhoei Z. Effect of dietary supplementation of *Ulva rigida* C. Agardh extract on several of physiological parameters of grey mullet,

- Mugil cephalus (Linnaeus). Sustainable Aquaculture and Health Management Journal. 2018;4(1):59–68.
84. Yeganeh S, Adel M. Effects of dietary algae (*Sargassum ilicifolium*) as immunomodulator and growth promoter of juvenile great sturgeon (*Huso huso* Linnaeus, 1758). J Appl Phycol. 2019;31(3):2093–102.
 85. Baker ME. Beyond carrier proteins: albumin, steroid hormones and the origin of vertebrates. J Endocrinol [Internet]. 2002;175:121–7. Available from: https://www.researchgate.net/profile/Michael-Baker-11/publication/11079965_Albumin_steroid_hormones_and_the_origin_of_vert_ebrates/links/00b4951d6f94ea8476000000/Albumin-steroid-hormones-and-the-origin-of-vertebrates.pdf
 86. Al-Deriny SH, Dawood MA, Abou Zaid AA, Wael F, Paray BA, Van Doan H, et al. The synergistic effects of *Spirulina platensis* and *Bacillus amyloliquefaciens* on the growth performance, intestinal histomorphology, and immune response of Nile tilapia (*Oreochromis niloticus*). Aquac Rep. 2020;17:390–7.
 87. Xiang Y, Shuning H, Lingke Z, Cheng H. The analysis of the effect of reducing triglyceride level in serum of hyperlipidemia model rats by feeding *Spirulina maxima*. Journal of Nanjing University (natural Sciences) [Internet]. 2002;38(2):182–6. Available from: <https://europemc.org/article/cba/383078>
 88. Colla LM, Muccillo-Baisch AL, Costa JA V. *Spirulina platensis* effects on the levels of total cholesterol, HDL and triacylglycerols in rabbits fed with a hypercholesterolemic diet. Brazilian Archives of Biology and Technology. 2008;51:405–11.
 89. Mg M. Effects of *Ascophyllum* and *Spirulina* meal as feed additives on growth performance and feed utilization of red sea bream, *Pagrus major*. J Fac Appl Biol Sci [Internet]. 1994;33:125–32. Available from: <https://cir.nii.ac.jp/crid/1573105975387643904>
 90. Brito R, Chimal ME, Gelabert R, Gaxiola G, Rosas C. Effect of artificial and natural diets on energy allocation in *Litopenaeus setiferus* (Linnaeus, 1767) and *Litopenaeus vannamei* (Boone, 1931) early postlarvae. Aquaculture. 2004;237(1–4):517–31.
 91. Anand PS, Kohli MPS, Roy SD, Sundaray JK, Kumar S, Sinha A, et al. Effect of dietary supplementation of periphyton on growth performance and digestive enzyme activities in *Penaeus monodon*. AQUACULTURE. 2013;392:59–68.
 92. Cahu CL, Infante JZ, Peres A, Quazuguel P, Le Gall MM. Algal addition in sea bass (*Dicentrarchus labrax*) larvae rearing: effect on digestive enzymes. Aquaculture. 1998;161(1–4):479–89.
 93. Lu J, Takeuchi T. Spawning and egg quality of the tilapia *Oreochromis niloticus* fed solely on raw *Spirulina* throughout three generations. Aquaculture. 2004;234(1–4):625–40.
 94. Teimouri M, Yeganeh S, Mianji GR, Najafi M, Mahjoub S. The effect of *Spirulina platensis* meal on antioxidant gene expression, total antioxidant capacity, and lipid peroxidation of rainbow trout (*Oncorhynchus mykiss*). Fish Physiol Biochem. 2019;45:977–86.
 95. Misra S, Niyogi S. Selenite causes cytotoxicity in rainbow trout (*Oncorhynchus mykiss*) hepatocytes by inducing oxidative stress. Toxicology in vitro. 2009;23(7):1249–58.
 96. Abdel-Latif HM, Khalil RH. Evaluation of two phytobiotics, *Spirulina platensis* and *Origanum vulgare* extract on growth, serum antioxidant activities and

- resistance of Nile tilapia (*Oreochromis niloticus*) to pathogenic *Vibrio alginolyticus*. *Int J Fish Aquat Stud*. 2014;250:250–5.
97. Xie S, Chen P, Lv Z, Lan Z, Zou J. Effects of *Spirulina platensis* on muscle nutritive components and growth of *Carassius auratus*. *Journal of Zhongkai University of Agriculture and Engineering* [Internet]. 2015;28(2):9–21. Available from: <http://znjx.cbpt.cnki.net>
 98. Tayag CM, Lin YC, Li CC, Liou CH, Chen JC. Administration of the hot-water extract of *Spirulina platensis* enhanced the immune response of white shrimp *Litopenaeus vannamei* and its resistance against *Vibrio alginolyticus*. *Fish Shellfish Immunol*. 2010;28(5–6):764–73.
 99. Lin YC, Tayag CM, Huang CL, Tsui WC, Chen JC. White shrimp *Litopenaeus vannamei* that had received the hot-water extract of *Spirulina platensis* showed earlier recovery in immunity and up-regulation of gene expressions after pH stress. *Fish Shellfish Immunol*. 2010;29(6):1092–8.
 100. Abdelkhalek NK, Ghazy EW, Abdel-Daim MM. Pharmacodynamic interaction of *Spirulina platensis* and deltamethrin in freshwater fish Nile tilapia, *Oreochromis niloticus*: impact on lipid peroxidation and oxidative stress. *Environmental Science and Pollution Research*. 2015;22:3023–31.
 101. Buege JA, Aust SD. [30] Microsomal lipid peroxidation Methods in enzymology. In: Academic press. Academic press; 1978. p. 302–10.
 102. Liu B, Xie J, Ge X, Xu P, Wang A, He Y, et al. Effects of anthraquinone extract from *Rheum officinale* Bail on the growth performance and physiological responses of *Macrobrachium rosenbergii* under high temperature stress. *Fish Shellfish Immunol*. 2010;29(1):49–57.
 103. Lin YH, Shiau SY. Dietary vitamin E requirement of grouper, *Epinephelus malabaricus*, at two lipid levels, and their effects on immune responses. *Aquaculture*. 2005;248(1–4):235–44.
 104. Luo P. Effect of *Spirulina* on the growth performance of *Cyprinus carpio* var *jian*. *J Hydrol*. 2006;26(2006):41–2.
 105. Moreira LM, Behling B, Rodrigues RS, Costa JA V, Souza-Soares LA. *Spirulina* as a protein source in the nutritional recovery of wistar Rats. *Brazilian Archives of Biology Technology*. 2013;56(3):447–56.
 106. Karakatsouli N. An overview of the use of fatty acids in the fish farming research during the last decade, with particular emphasis on fish quality. *Journal of World Aquaculture Society*. 2012;43:291–320.
 107. Bhavan PS, Devi VG, Shanti R, Radhakrishnan S, Poongodi R. Basic biochemical constituents and profiles of amino acids in the post larvae of *Macrobrachium rosenbergii* fed with *Spirulina* and yeast enriched *Artemia*. *J Sci Res* [Internet]. 2010;2(3):539–49. Available from: <http://dx.doi>
 108. Prakoso VA, Pouil S, Cahyanti W, Sundari S, Arifin OZ, Subagja J, et al. Fluctuating temperature regime impairs growth in giant gourami (*Osphronemus goramy*) larvae. *Aquaculture*. 2021;539:606–736.
 109. Syandri H, Mardiah A, Aryani N. Effect of stocking density on the performance of juvenile gourami sago (*Osphronemus goramy*) in the synthetic sheet pond. *Pak J Zool* [Internet]. 2020;52(2):717. Available from: <https://www.proquest.com/openview/82b67628eb7695544259a655627e3825/1?pq-origsite=gscholar&cbl=616534>

110. Boyd CE, Tucker CS, Somridhivej B. Alkalinity and hardness: critical but elusive concepts in aquaculture. *J World Aquac Soc.* 2016;47(1):6–41.

UNDER PEER REVIEW