

Effect of chemical mutagens on vegetative growth of *Calendula officinalis* L. (cv. Calypso orange)

ABSTRACT

The present investigation entitled “Effect of chemical mutagens on vegetative growth of *Calendula officinalis* L. cv. Calypso orange” was carried out at the laboratory of floriculture and experimental farm of Mata Gujri college, Fatehgarh Sahib, Punjab. The experiment was laid out in randomized block design (RBD) with nine treatments and three replications. The treatments were T₁ i.e., Sodium azide @1500ppm, T₂ i.e., Sodium azide @2500ppm, T₃ i.e., Sodium azide @3500ppm, T₄ i.e., Sodium azide @4500ppm, T₅ i.e., Diethyl sulphate @1500ppm, T₆ i.e., Diethyl sulphate @2500ppm, T₇ i.e., Diethyl sulphate @3500ppm, T₈ i.e., Diethyl sulphate @4500ppm and T₉ i.e., control. Observations were recorded for vegetative among all treatments maximum plant height (24.84cm), number of leaves per stem (8.95), number of stems per plant (7.48), stem length (22.85cm), leaf length (9.48cm), plant spread (24.89cm²) and diameter of main stem (9.39mm) were found maximum in T₁ i.e., Sodium azide @1500ppm. Best vegetative parameters were recorded in Sodium azide @1500ppm. Therefore, Sodium azide @1500ppm is best dosage for *Calendula officinalis* cv. Calypso orange.

Keywords: Calendula, Sodium azide, Diethyl sulphate.

INTRODUCTION

In recent decades, the floriculture business has grown significantly. A breeder goals are always to generate new and unusual cultivars of ornamentals, and this process is never-ending (Datta, 2009). Pot marigold, or *Calendula officinalis* L., belongs to the Asteraceae family and has a chromosomal number of 2n=32. The plant's vegetative sections are green, and the stems have small hairs covering them. According to Filipovic *et al.*, (2016), the flower's colour varies from yellow to orange. The flower contains disc and ray florets, which may open up in the spring (Jan *et al.*, 2017).

The herb *Calendula officinalis* is native to southern Europe and the Eastern Mediterranean region. It can be an annual or short-lived perennial. It is a popular garden escape in cold, temperate areas and has been cultivated for a very long time. It has been utilized for a variety of gastronomic and therapeutic uses for many ages. Common names for this species include pot marigold, English marigold, and Scotch marigold. It is important to distinguish this species from marigolds, which are a group of numerous ‘*Tagetes*’ species. Field marigold is a branching annual plant native to southern Europe. Its daisies are one inch in size and are either yellow or orange in colour.

The herbaceous plant *Calendula officinalis* reaches a height of 30 to 60 cm. The plant is heterophyllous in a single individual, with lanceolate upper and lower leaves, and an

Comment [R1]: Can improve grammatically.

Comment [R2]: Also add future direction and what impact this result can make on market or farms.

Comment [R3R2]: Like Suggested addition: These findings can inform floriculturists and plant breeders about optimal mutagen dosages for enhancing the growth of *Calendula officinalis*.

Comment [R4]: I won't use word unusual here as it can have both negative and positive meanings.

inflorescence resembling a basket with two distinct types of flowers. Its distinctive features include ligular blooms on the outside and tubular flowers on the inside.

Comment [R5]: Citation missing in this part. There are certain informations here which needs citation.

MATERIAL AND METHODS

Comment [R6]: Statement of the problem and rationale for doing this research are all missing. Please complete that part.

The present investigation was conducted at Research Farm, Mata Gujri College, Fatehgarh Sahib, Punjab during 2023-2024. Field of experimental site lies at 30.6435° North latitude and 76.3970° East longitudes. The altitude of the location is 246 meter above the mean sea level. Seeds of the calendula flower cv. Calypso orange was purchased from the Biocarve seeds, Dhablan (Patiala).

Comment [R7]: Missing statistical analysis part here. Include what tool and methods you used for collecting and analyzing data

Experimental design

The experiment was laid out in Randomized Block Design (RBD) with three replications. The seedlings of uniform size (3-5cm) used for treatment of chemicals on vegetative growth and then planted in the beds with a spacing of 25 cm plant to plant and 30 cm row to row during November 2023.

Experimental details

There were 9 treatments of chemicals dosages which were applied to calypso orange cultivar of calendula. The treatments were T₁ i.e., Sodium azide @1500ppm, T₂ i.e., Sodium azide @2500ppm, T₃ i.e., Sodium azide @3500ppm, T₄ i.e., Sodium azide @4500ppm, T₅ i.e., Diethyl sulphate @1500ppm, T₆ i.e., Diethyl sulphate @2500ppm, T₇ i.e., Diethyl sulphate @3500ppm, T₈ i.e., Diethyl sulphate @4500ppm and T₉ i.e., control.

Comment [R8]: If possible, lay out these treatments in a table so its easier to read.

RESULTS AND DISCUSSIONS

Plant height (cm)

The maximum plant height was observed in T₁ i.e., @1500ppm Sodium Azide (24.84cm) which was statistically superior than other treatments. At low concentration of Sodium azide increase in the rate of cell elongation or division as well as auxin activation may be responsible for this mutagen's effect. This finding was similar with Zaka *et al.*, (2004) and Joshi *et al.*, (2011). T₄ i.e., Sodium Azide @ 4500 ppm recorded minimum plant height (21.03cm) of calendula cv. calypso orange which was statistically at par with T₆ (21.86 cm @2500 ppm Diethyl Sulphate) due to the physiological harm brought on by Sodium azide and its hydrolysis products may be the cause of these outcomes at higher concentration of sodium azide. The present study's findings are also in line with observations made *Asparagus densiflorus* by Asrar and El- Nashar (2016).

Number of leaves per stem

There was significant effect on the number of leaves per stem of calendula cv. calypso orange. The T₁ treatment i.e., @1500ppm Sodium Azide (8.95) produced maximum number of leaves per stem on calendula cv. calypso orange which was statistically at par with T₂ i.e.,

Sodium Azide @2500ppm (8.33), T₃ i.e., Sodium Azide@3500ppm (8.82), T₄ i.e., Sodium Azide @4500ppm (8.86), T₅ i.e., Diethyl Sulphate @ 1500ppm (8.55), T₆ i.e., Diethyl Sulphate @2500ppm (8.48), T₇ i.e., Diethyl Sulphate @3500ppm (8.87) and T₉ i.e., control (8.38) due to the low concentration of sodium azide the number of leaves per stem was increased due to increase the metabolic activities of plants, and it also enhance nutrient absorption and photosynthesis process. The same results were also reported by Mohamed *et al.*, 2019. Minimum number of leaves per stem was discovered in T₈ i.e., Diethyl Sulphate @ 4500ppm (7.58) which was statistically inferior. The higher concentration of chemical mutagen is toxic for the calendula plant that can cause severe damage to plant cells and disrupt essential biological processes. The same findings were also stated by Gvozdenovic *et al.*, (2009) on the sunflower.

Number of stems per plant

The maximum number of stems per plant was recorded in T₁ i.e., @1500ppm Sodium Azide (7.48) which was statistically at par with T₅ i.e., Diethyl Sulphate @2500ppm (7.12), T₇ i.e., Diethyl Sulphate @ 3500ppm (7.25), and T₉ i.e., control (7.08). These results were observed might be due to the physiological effects of sodium azide and the products of their hydrolysis may be responsible for these results, as well as the reason for the increase in stem count at low concentration of chemical mutagen. The result was closely related with El-Nashar (2006). The less number of stems per plant was found in T₈ i.e., Diethyl Sulphate @4500ppm (6.50) which was statistically at par with T₂ (6.88) i.e., Sodium Azide @2500ppm, T₄ (6.72) i.e., Sodium Azide @4500ppm, and T₆ (6.72) i.e., Diethyl Sulphate @2500ppm. Diethyl sulphate is a powerful mutagen at higher concentration it can cause changes in the DNA of plants, disrupt the cell division, hormonal balance with chemical mutagens on plant genetics and development. In connection with this, Sharma *et al.*, 2023 also report the same conclusion.

Stem length (cm)

This data shows significant effect of chemical mutagens on the stem length of calendula cv. calypso orange. The maximum stem length was observed in T₁ i.e., Sodium Azide @1500 ppm Sodium Azide (22.85 cm) which was statistically at par with T₄ i.e., Sodium Azide @4500ppm (21.85cm) and T₇ i.e., Diethyl Sulphate @3500ppm (21.94cm). These results might be due to effect of the mutagen, resulting in a higher rate of cell elongation or division, could be linked to either auxin or cytokinin at low concentration of chemical mutagen. The same study was reported by Khan *et al.*, (2024). The shortest stem length per plant was observed T₈ i.e., Diethyl Sulphate@ 4500ppm (20.01cm). Diethyl sulphate due to the higher concentration of Diethyl sulphate interference with cell elongation pathways and hormonal balance was highlighted in the study as a major factor contributing to the decrease in stem length and this chemical can also disrupt plant respiration by inhibiting cytochrome oxidase, an enzyme crucial for the electron transport chain in mitochondria. As discovered and reported by Gupta *et al.*, 2023.

Leaf length (cm)

In T₁ i.e., @1500ppm Sodium Azide (9.48cm) the maximum leaf length of calendula cv. calypso orange was observed. At low concentration of sodium azide, expression of related genes and the molecular level that affects the gene or groups of genes and these genes could be those controlling the synthesis of growth regulators such as auxins and cytokinins and same result mentioned by Cain *et al.*, 2006 and Joshi *et al.*, 2011. The smallest leaf length of calendula cv. calypso orange per plant was observed in T₈ i.e., Diethyl Sulphate@ 4500ppm (6.92cm), which was statistically inferior due to its cell elongation processes are inhibited by high concentrations of diethyl sulphate, which limits growth and prevents plants from developing their leaves to their normal length and these findings as reported by Sharma *et al.*, (2023).

Plant spread (cm²)

The maximum plant spread data was observed in T₁ i.e., @1500 ppm Sodium Azide (24.89 cm²) which was statistically at par with T₇ i.e., Diethyl Sulphate@ 3500ppm (23.07cm²). The stimulatory effect of low dose sodium azide may contribute to better nutrient absorption and hormonal signalling as reported by Kayalvizhi *et al.*, 2023. The lowest data of plant spread was noted in T₈ i.e., Diethyl Sulphate@ 4500ppm (18.90cm²) which was statistically at par with T₂ i.e., Sodium Azide @2500ppm (21.02cm²) and T₉ i.e., control (21.32cm²) the study demonstrated the toxicity of diethyl sulphate at high concentrations, which can cause physiological disturbances in plant cells and metabolic processes. This study was reported by Sharma *et al.*, (2023).

Diameter of main stem (mm)

There was significant effect of chemical mutagens on calendula cv. calypso orange. the main stem's maximum diameter was recorded in T₁ 1500ppm Sodium Azide (9.39mm), which was statistically at par with T₅ i.e., Diethyl Sulphate@ 1500ppm. (8.57mm). T₈ Diethyl Sulphate@4500ppm (7.18mm). Which might be due to the stimulatory effect of the mutagen may be linked to an increase in the rate of cell division or elongation as well as an activation of auxin (Zaka *et al.*, 2004 and Joshi *et al.*, 2011). The minimum diameter of main stem was noted in T₈ Diethyl Sulphate@4500ppm (7.18mm) which was statistically at par with T₂ i.e., Sodium Azide @ 2500ppm (7.52mm) T₄ i.e., Sodium Azide @ 4500ppm (7.97mm) T₉ i.e., control (7.70mm), diethyl sulphate is the highly toxic compound at higher concentration that can lead to cell death, inhibition of cell division, and disruption of vascular tissues in the main stem. These findings are related with Kannan *et al.*, 2002.

Table 1: Effect of chemical mutagens on vegetative growth of *Calendula officinalis* L. (cv. Calypso orange):

TREATMENTS	Plant height(cm)	Number of leaves per stem	Number of stems per plant	Stem length (cm)	Leaf length (cm)	Plant spread (cm ²)	Diameter of main stem (mm)
T ₁ Sodium azide@1500ppm	24.84	8.95	7.48	22.85	9.48	24.89	9.39
T ₂ Sodium azide@2500ppm	22.23	8.33	6.88	21.38	8.23	21.02	7.52
T ₃ Sodium azide@3500ppm	22.85	8.82	7.03	21.61	8.62	22.43	8.35
T ₄ Sodium azide@4500ppm	21.03	8.86	6.72	21.85	8.12	21.79	7.97
T ₅ Diethyl sulphate@1500ppm	23.48	8.55	7.12	22.30	8.52	21.76	8.57
T ₆ Diethyl sulphate@2500ppm	21.86	8.48	6.72	21.36	8.39	21.68	8.11
T ₇ Diethyl sulphate@3500ppm	23.58	8.87	7.25	21.94	8.38	23.07	8.34
T ₈ Diethyl sulphate@4500ppm	22.13	7.58	6.50	20.01	6.92	18.90	7.18
T ₉ Control	23.40	8.38	7.08	21.29	8.13	21.32	7.70

SEm±	0.32	0.24	0.15	0.40	0.25	0.81	0.29
CD _{0.05}	0.97	0.73	0.44	1.21	0.76	2.43	0.87

CONCLUSION

From the present study it can be concluded that the lowest concentration of Sodium Azide (1500ppm) reported the best performance of vegetative parameters such as plant height, number of leaves per stem, number of stems per plant, stem length, leaf length, plant spread, and diameter of main stem.

From the results, it can be concluded that Sodium Azide @1500ppm is best dosage for *Calendula officinalis* cv. Calypso orange.

REFERENCES

- Badr M, EL-Shennaway O, Mostafa M and EL-Tony F. (2000) Effect of gamma irradiation, ethyl methane sulphate and their combinations on growth, flowering and induced variability in *Tagetes erecta* L. *Journal Agriculture Science*. **25**,3587- 3604.
- Bhat T A, Sharma M, and Anis M. (2007). Comparative analysis of meiotic aberrations induced by diethyl sulphate and sodium azide in broad bean (*Vicia faba* L.). *Asian Journal of Plant Sciences*, **6** (7), 1051- 1057.
- El-Khateeb, M A, El-Attar, A B, Fayed, R G. (2022). Comparative study on the effect of chemical mutagens of sodium azide and di-ethyl sulphate on improving morphological traits and yield components of *Borago officinalis* L, plant. *International Journal of Health Sciences*, **6**(S4), 10881-10898.
- El-Mokadem Hoda E, Mostafa Gehan G (2014). Induction of mutations in *Browallia speciosa* using sodium azide and identification of the genetic variation by peroxidase isozyme. *African J. of Biotechnology*, **13**(1): 106-111.
- El-Nashar, Y I and Ammar, M H (2016). Mutagenic influences of colchicine on phenological and molecular diversity of *Calendula officinalis* L. *Genetics and Molecular Research*, **15**(2):1-15.

Comment [R9]: You have made conclusions based on only ANOVA. I would suggest to perform post-hoc tests (eg. Turkey) to identify which specific treatments differs from each other. Also, perform normality and homogeneity of variance test to check normality of your data. AND one more suggestion will be to present some data in graphs as well.

El-Nashar, Y I, and Asrar, A A (2016). Phenotypic and biochemical profile changes in calendula (*Calendula officinalis* L.) plants treated with two chemical mutagenesis. Genetics and Molecular Research, **15**(2):1-14.

El-Nashar, Y I. (2006). Effect of chemical mutagens (sodium azide and diethyl sulphate) on growth, flowering and induced variability in *Amaranthus caudatus* L. and *A. hypochondriacus* L. Ph. D. Thesis Faculty of Agriculture, Alex. Univ. A.R.E.

Gupta, M and Singh S. (2010). *Borago officinalis* L., An important medicinal plant of Mediterranean region: a review. International Journal Pharmacy Science Rev. Res., **5**: 27- 34.

Gvozdenovic, S, Bado, S, Afza, R, Jovic, S and Mba, C. 2009. Interval differences in response of sunflower (*Helianthus annuus* L.) to different mutagenic treatments. In: Induced Plant Mutations in the Genomics Era, Q.Y. Shu (ed.), Food and Agriculture Organization of the United Nations, Rome, 358-360.

Jan N, Andrabi K I and Riffat J. 2017. *Calendula officinalis* - An important medicinal plant with potential, biological properties. Proc. Indian National Sci. Acad., **83**(4), 769-787.

Joshi, N, Ravindran A and Mahajan V (2011). Investigations on chemical mutagen sensitivity in onion (*Allium cepa* L.). Int. J. Bot., **7**: 243-248. 27.

Kannan, M, Sathiyamurthy, V A and Sankar, V (2002). Mutagenetic studies on *Jasminum sambac*. Floriculture research trend in India. Proceedings of the National Symposium on Indian Floriculture in the New Millennium, Lal Bagh, Bangalore, 25- 27, February, 209-211.

Kapadiya D B, Chawla SL, Patel A L Ahlawat TR (2014). Exploitation of variability through mutagenesis in *Chrysanthemum morifolium* Ramat.) var. Maghi, Journal of Life Sci., **9**(4):1799-1804.

Kayalvizhi K, Kannan M, Ganga M. Effect of Physical and Chemical Mutagens on Morphological Characters in M1V2 Generation of Tuberose (*Polianthes tuberosa* L.). International Journal of Current Microbiology and Applied Sciences. 2017, **6**(4):2492-2499.

Khan S, Al-Qurainy F, Anwar F. Sodium azide: a chemical mutagen for enhancement of agronomic traits of crop plants, Environmental and We an International Journal of Science and Technology, **4**, 2009, 1-21. 17.

Mohamed, M A H, M K Aly, G G, Mostafa and H.R AbdAlaziz. 2019. Evaluation of *Chrysanthemum morifolium* cv. Maghi plants after sodium azide treatment. Scientific Journal of Agricultural Sciences, **1**(1): 14-20.