

Analysis of the phytochemical composition of extracts of the aerial parts of *Ageratum conyzoides* L. (Asteraceae)

ABSTRACT

Introduction: *Ageratum conyzoides* L. (Family Asteraceae) is an annual herb with a long history of traditional medicinal use in the tropical and sub-tropical region of the world. The present study was designed to screen out the major groups of phytochemicals found in the aerial part of the plant.

Methodology: The aqueous and ethanolic extracts of the leaves, stem and flowers of *A. conyzoides* as well as the entire aerial part of the plant as a whole underwent qualitative using specific standard methods. The identified classes of metabolites in the different extracts were then quantified using spectrophotometric methods.

Results: Phytochemical analysis of the *A. conyzoides* extracts revealed the presence of tannins, alkaloids, flavonoids, phenol, steroids, phlobatannins, saponins, and coumarins. The quantitative phytochemical analysis of the extracts of *A. conyzoides* revealed that the aqueous maceration of the leaves of *A. conyzoides* had the highest content in polyphenols, flavonoids, tannins with the exception of the total alkaloid content which was highest in the hydroethanolic maceration of the aerial part of the plant.

Conclusion: The quantitative phytochemical analysis of the extracts of *A. conyzoides* revealed that the aqueous maceration of the leaves of *A. conyzoides* had the highest content in most of the metabolites. This study also shows that the extracts of *A. conyzoides* are extensively rich in secondary metabolites which could explain the application of the plant in the treatment of several ailments in traditional medicine.

Keywords: *Ageratum conyzoides*, Aerial, Phytochemical analysis, Plant metabolites, Maceration.

1. INTRODUCTION

The plant kingdom is a treasure house of potential drugs and in the recent years there has been an increasing awareness about the importance of medicinal plants [1]. According to the World Health Organization more than 80% of the world's people depend on traditional medicine for their

primary healthcare needs[2]. The reason being that mankind, in his attempts to find solutions to his healthcare problems, has resorted to using plant-derived therapies to remedy the shortage of reliable cures. Medicinal plants used in different diseases and ailments are the richest bio reservoirs of various phytochemicals[3]. Therefore, a systematic approach should be made to unlock the hidden potential of these ethnomedicinal plants[4].

Ageratum conyzoides is one of the well-known plants which has been reported to possess diverse medicinal properties. *A. conyzoides* (billy goat weed; Asteraceae) is native to the tropical Americas, but it is widely distributed in tropical and sub-tropical regions throughout the world[5]. It has a peculiar odor likened in Australia to that of a male goat and hence its name 'goatweed'[6]. It is an annual branching herb which grows to approximately 1 m in height. The stems and leaves are covered with fine white hairs, the leaves are ovate and up to 7.5 cm long. The flowers are purple to white, less than 6 mm across and arranged in close terminal inflorescences[7]. The plant grows commonly in the proximity of habitation, thrives in any garden soil and is very common in waste places and on ruined sites[8].

A. conyzoides has been known since ancient times for its curative properties and has been utilized for the treatment of various ailments, such as burns and wounds, headaches, pneumonia, analgesic, inflammation, asthma, spasmotic and hemostatic effects, stomach ailments, gynecological diseases, leprosy and other skin diseases[9]. In Cameroon and Congo, it is used to treat fever, rheumatism, headache and colic[10]. The aqueous extract of leaves or whole plant is utilized to treat colic, cold and fevers, rheumatism in Brazil [11]. Due to the fact that different parts of *A. conyzoides* have traditional medicinal applications, the present study was therefore planned to define and quantify the phytochemicals present in the aerial parts (leaves, flowers, stem) of the plant in order to provide a scientific basis to justify its traditional therapeutic application.

MATERIALS AND METHODS

Sample collection and authentication

The plant specimen was deposited and identified in the National Herbarium of Cameroon (NHC) by comparing with reference specimen number 61801/NHC. Fresh mature plants were harvested in Yaoundé in January 2023. The harvested plant was separated into individual aerial parts (leaves, stems and flowers) and aerial part as a whole of *A. conyzoides*. The plant material was air-dried separately away from sunlight at room temperature under shade and ground to fine powder using a milling machine. The fine powder was stored in clean labeled airtight containers till required for analysis.

Preparation of plant extracts

In this study, aqueous and hydroethanolic extracts of each aerial part (leaves, stem and flowers) as well as the whole aerial part of *A. Conyzoides* were prepared separately using the maceration technique.

Maceration

In maceration (for fluid extract), whole or coarsely powdered plant-drug is kept in contact with the solvent in a stoppered container for at least two days with frequent agitation until soluble matter is dissolved. At the end of extraction, the micelle is separated from marc by filtration and from the menstruum by evaporation in an oven.[16]

Procedure

In this process, 100 g of coarsely powdered plant's aerial part, leaves, stem and flowers were placed in a stoppered container with 1000 mL of solvent (distilled water, hydro-ethanolic solution 50:50) and allowed to stand at room temperature for a period of 72 hours with frequent agitation until the soluble matter has dissolved. The mixture was then strained, the marc (the damp solid material) pressed, and the combined liquids were clarified by filtration using Whatman paper. At the end of extraction, the micelle is separated from marc by filtration and from the menstruum by evaporation in an oven at 50°C[13]. The dried extract was stored in an air-tight container and kept in refrigerator at 4°C for their future use in phytochemical analysis.

Determination of extraction yield (%)

Each extract of *A. conyzoides* was weighed in order to determine the percentage yield obtained from the initial powder quantity. The yield of evaporated dried extracts based on dry weight basis was calculated from the equation:

$$\text{Percentage yield (\%)} = \frac{\text{mass of the extract obtained}}{\text{mass of the initial powder}} \times 100$$

Phytochemical screening

Qualitative analysis

The various extracts of the leaves, stem and flowers of *A. conyzoides* as well as the aerial part as a whole were screened qualitatively for the phytochemical constituents utilizing standard methods of analysis by Sofowara (1993), Savithrama et al. (2011) and Balamurugan (2019) [14–16]. The filtrates of the *A. conyzoides* extracts were analyzed for the following phytochemicals: carbohydrates, proteins, glycosides, tannins, alkaloids, flavonoids, saponins, resins, quinones, cardiac glycosides, coumarins, steroids, phenols, anthraquinones and phlobatannins. The tests were based on the visual observation of color change or formation of a precipitate after the addition of specific reagents.

Qualitative analysis of primary metabolites

Test for carbohydrates

Fehling's test: About 1 mL of the filtrate was taken to which 1 mL of Fehling's reagent is added and boiled in a water bath. The appearance of red precipitate indicates the presence of sugars.

Molisch's test: To about 2 mL of the sample, 2 drops of alcoholic solution of α -naphthol are added and to the mixture after being shaken well. Few drops of conc. H_2SO_4 was added along the sides of the test tube. A violet ring indicates the presence of sugars. [15]

Test for proteins

Biuret test: To 2 mL of filtrate was added 1 drop of 2% copper sulphate solution; and 1 mL of 95% ethanol was added. This was followed by excess addition of KOH. The appearance of pink color indicates the presence of protein.

Millon's test: To 2 mL of filtrate was added a few drops of Millon's reagent. The appearance of a white precipitate indicates the presence of proteins. [15]

Test for lipids

To the extract, few drops of 0.5N alcoholic KOH and few drops of phenolphthalein were added. This mixture was heated for about 2 hours. The formation of soap or partial neutralization of alkali indicates the presence of fixed oils or fats. [15]

Miscellaneous compounds

Test for resins

1 mL of extract was taken and to this few mL of acetic anhydride was added to 1 mL of conc. H_2SO_4 . The appearance of orange to yellow color indicates the presence of resins. [15]

Test for gums and mucilage

To 1 mL of extract, distilled water, 2 mL of absolute ethanol was added accompanied by constant stirring white or cloudy precipitate indicates the presence of gums or mucilage. [15]

Vitamin C

In 5ml of distilled water, 1ml of the sample was diluted and a drop of 5% sodium nitroprusside and 2ml of NaOH is added. Few drops of HCl are added dropwise, the yellow color turns blue. This indicates the presence of vitamin- C. [15]

Carotenoids

To two test tubes containing 2 mL of the plant extract and 2 mL of water each, 2 mL of HCL and 2 mL of H_2SO_4 were added. The appearance of a blue-green color in both test tubes confirms the presence of carotenoids. [16]

Qualitative analysis of secondary metabolites

Test for anthraquinones

To 5ml of extract, few ml of conc. H_2SO_4 was added and 1ml of diluted ammonia was added to it. The appearance of rose pink confirms the presence of anthraquinones. [15]

Test for quinones

To 1 mL of extract, alcoholic KOH was added. The presence of red to blue color indicates the presence of quinones.[16]

Test for alkaloids

Mayer's test: to a few mL of filtrate, 2 drops Mayer's reagent was added. A creamy or white precipitate shows a positive result for alkaloids.

Wagner's test: (iodine – potassium iodine reagent): To about an mL of extract few drops of Wagner's reagent was added. Reddish – brown precipitate indicates presence of alkaloids.

Hager's test: To a few mL of the filtrate was added 2 mL of Hager's reagent. A creamy white/ yellow precipitate shows a positive result for alkaloids.

Tannic acid test: To the acidified extract, 10% tannic acid solution was added. A buff color precipitate indicates the presence of alkaloids. [15]

Test for glycosides

2 mL of extract is mixed with a few drops of glacial acetic acid containing traces of ferric chloride and 0.5 mL of conc. H_2SO_4 was added. The production of blue color is positive for glycosides. [15]

Test for cardiac glycosides (Keller-Killani test)

5mL of solvent extract was mixed with 2mL of glacial acetic acid and a drop of ferric chloride solution followed by the addition of 1mL of conc. H_2SO_4 . A brown ring in the interface indicates the presence of deoxy sugars of cardenolides. A violet ring may appear beneath the brown ring while acetic acid layer a green ring may also form just gradually towards the layer. [15]

Test for polyphenols

To the extract was added few drops of 5% ferric chloride. The formation of dark green color indicates presence of polyphenols. To 1 mL of extract, few drops of 5% solution of lead acetate were added. The appearance of yellow precipitate indicates the positive results for polyphenols. [15]

Test for tannins

To 5mL of extract, few drops of neutral 5% ferric chloride solution was added. The production of dark green color indicates the presence of tannins. [14]

Differentiation of catechic and gallic tannins

It is obtained by STIASNY reaction, which is carried out in the following manner. To 30mL of the extract, we add 15mL of STIASNY reagent (10mL of 40 % formalin and 5 mL of concentrated HCl) and heated for 15 minutes in a water bath at 90 °C.

Catechic tannins: The obtaining of a precipitate shows their presence;

Gallic Tannins: After filtration, we saturate the filtrate with powdered sodium acetate, then 1 mL of a solution of 1 % ferric perchloride (FeCl_3). The presence of gallic tannins not precipitated by the STIASNY reagent is indicated by the development of a shade dark blue. [14]

Test for flavonoids

In a test tube, put 2 mL of the 1 % extract. Add 1 mL of sodium hydroxide (NaOH) of 2N. The formation of a yellow-orange coloration indicates the presence of flavonoids. The addition of a few drops of sulfuric acid and the change of coloring confirm the presence of flavonoids. [14]

Test for phlobatannins

Aqueous extract was boiled with diluted HCl leading to the deposition of reddish precipitate indicates the presence of phlobatannins. [15]

Test for beta cyanide

2 mL of the extract was introduced in a test tube and 2 mL of 2N NaOH was added and heated the tube in a boiling water bath for 5 minutes. The appearance of a yellow coloration indicates the presence of betacyanin. [16]

Test for saponins

0.5 mg of extract was vigorously shaken with few mL of distilled water. The formation of frothing is positive for saponins. The froth from the above reaction was taken and few drops of olive oil were added and shaken vigorously and observed for the formation of emulsion. [16]

Test for steroids

2 mL of extract with 2 mL of chloroform and 2mL of concentrated H_2SO_4 was added, the appearance of red color and yellowish green fluorescence indicates the presence of steroids. [15]

Test for anthocyanins

2 mL of aqueous extract was taken to which was added 2N HCl and it was followed by the addition of ammonia, the conversion of pink-red turns blue-violet indicates the presence of anthocyanins. [15]

Test for coumarins

To 2 mL of the extract, 3 mL of 10% aqueous solution of NaOH was added. The production of yellow color indicates the presence of coumarins. [15]

Test for oxalates

In a test tube, put 2 mL of the extract. A few drops of ethanoic acid were added. Obtaining a greenish-black color indicates the presence of oxalates. [14]

Test for chalcones


To 5 mL of the plant extract was added 2 mL of ammonia solution. The appearance of a red color confirms the presence of chalcones. [16]

QUANTITATIVE ANALYSIS

Depending on the qualitative results, the quantitative assay was carried out for carbohydrates, proteins, polyphenols, alkaloids and flavonoids.

Determination of total soluble carbohydrates

Picric acid is reduced into picramic acid by glucose. The intensity of the orange color of picramic acid will be proportional to the concentration of glucose. The orange color of picramic acid produced, whose maximum absorption is around 570 nm, is proportional to the quantity of phenolic compounds present in the plant extract [17]. The reaction is as follows:



In a test tube, 1000 μ L of 13% picric acid and 1000 μ L of 4% sodium hydroxide were added to 1000 μ L of the extract at a concentration of 1000 μ g/mL. The tube was placed in a boiling water

bath for 10 minutes. The calibration curve was used during the reading of the optical density at 570 nm against the extract and blank concentration. The amount of carbohydrate content was expressed as equivalent of glucose per mg of dry extract of the plant (EG/g) [17].

Determination of total protein content

Lowry's method is another colorimetric protein assay method, complementary to that of Biuret. In fact, the protein first reacts with an alkaline cupric reagent (Gornall's reagent of the biuret method) then a second reagent, called phosphotungstomolybdic (Folin-Ciocalteu's reagent), is added. This reagent allows the reduction of aromatic amino acids (tyrosine and tryptophan) leading to the formation of a dark blue colored complex whose absorbance is measured between 650 and 750 nm [18].

BSA was used as standard reagent for preparing the standard curve against which the unknown concentration of proteins was estimated. In a test tube, 1000 μ L of alkaline reagent was added to 1000 μ L of the extract at 1000 μ g/mL and incubated for 10 minutes followed by the addition of 500 μ L of the Folin-Ciocalteu reagent (diluted to 1/10th) was followed immediately by incubation at room temperature for 20 minutes in darkness to allow for the development of the blue color. The absorbance was read at 760 nm on a spectrophotometer. The quantity of protein content after that the absorbance was measured at 760 nm and the amount of protein expressed as μ g/mL of BSA (EBSA/g). [18]

Evaluation of total phenolic contents

To determine the total phenolic content, we used the method of Folin-Ciocalteu. Folin-Ciocalteu reagent is a yellow-colored acid consisting of a mixture of two acids: phosphotungstic acid ($H_3PW_{12}O_{40}$) and phosphomolybdic acid ($H_3PMO_{12}O_{40}$). It is reduced during the oxidation of polyphenols to form a stable blue complex of tungsten and molybdenum oxides. The color produced, whose maximum absorption is around 760 nm, is proportional to the quantity of phenolic compounds present in the plant extract.[19]

1000 μ L of the Folin-Ciocalteu reagent (diluted to 1/10th) was added to 1000 μ L of the extract at 1000 μ g/mL and the whole was incubated for two minutes at room temperature. Then, 1000 μ L of sodium bicarbonate at 75 g/L was added, followed immediately by incubation at room temperature for 90 minutes. The absorbance was read at 760 nm on a spectrophotometer. The quantity of phenolic compounds, expressed in milligrams of gallic acid equivalent per gram of

dry weight of the plant (GAE/g), was determined by the calibration curve varying from 0 to 1000 mg/mL, made with different concentrations of gallic acid.[20]

Evaluation of total flavonoid contents

The quantification of flavonoids was carried out using aluminum trichloride and sodium hydroxide. The aluminum trichloride forms a yellow complex with flavonoids and the soda forms a pink colored complex which absorbs in the visible electromagnetic spectrum at 510 nm [21].

In each tube, we introduced 1000 μ L of the extract at 1000 μ g/mL, followed by the addition of 150 μ L of sodium nitrite (NaNO_2) at 5%. After 5 minutes, 150 μ L of aluminum trichloride (AlCl_3) at 10% (m/v) was added to the mixture. After 6 minutes of incubation at room temperature, 500 μ L of 4% sodium hydroxide (NaOH) was added. The mixture was stirred immediately to homogenize its content. The absorbance of the pinkish colored solution was read at 510 nm against a blank. A calibration curve was produced in parallel under the same operating conditions using quercetin as a positive control. The total flavonoid content of the plant extracts studied was expressed in milligram equivalent of quercetin per gram of dry plant matter (mg EQ/g). [21]

Evaluation of total flavonols content

The sample containing flavonols results in the formation of a green color when reacted with aluminum chloride and sodium acetate, and the samples are read at 440 nm in a UV-Vis spectrophotometer [22].

In a 10 ml test tube, 1000 μ l of extracts at a dose of 1000 μ g/ml, and 1000 μ l of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ 2%, 600 μ l of sodium acetate 50g/l, was completed with distilled water to the volume of 3000 μ l. The tubes were incubated at room temperature for 2.5 hours. The solution was mixed well and the absorbance was measured against the blank not containing the extract at 440 nm. The standard curve for flavonols was carried out using a standard solution of quercetin (0 to 200 μ g/l) according to the same procedure as that which was previously described. The total flavonols were expressed in milligrams of quercetin equivalents per g of dried fraction (mg EQ/g). [22]

Evaluation of total alkaloids content

The alkaloid, in contact with concentrated sulfuric acid and potassium dichromate, develops a violet line which turns blue then green, so the maximum absorption proportional to the intensity of the color developed is 650 nm [23].

In a glass test tube, to 1000 μ l of the sample at 1000 μ g/ml, was added 1000 μ L of 5% potassium dichromate and incubation was allowed for 5min at 30°C. 8000 μ l of concentrated sulfuric acid was then added to the tubes content and mixed. The tubes at room temperature were allowed for 20min to observe any color change and then read at the optical density of 650nm against the blank or the sample. The alkaloid concentration was obtained from the calibration curve and expressed in milligrams of quinine hydrochloride equivalents per gram of dried fraction (QHE/g). [23]

Evaluation of total tannins content

The technique for assaying condensed tannins by the Folin-Ciocalteu method is based on the reduction of phosphomolybdic and tungstic acid in an alkaline medium, in the presence of tannins, to give a blue color whose intensity is measured between 640 and 760 nm [24].

In a 10 ml test tube, 1000 μ l of extracts at a dose of 1000 μ g/ml, 200 μ l of Folin's reagent diluted to one tenth and 1000 μ l of 35% Na₂CO₃ were added. The solution was mixed well and the absorbance was measured against the blank containing no extract at 700 nm. The standard curve for total tannins was produced using a standard solution of tannic acid (0 to 500 μ g/ml) according to the same procedure as that described previously. Total tannins were expressed in milligrams of tannic acid equivalents per g of dried fraction (TAE/g). [24]

Ethical considerations

The study was conducted after having approval from the institutional review board of the Faculty of Medicine and Biomedical Sciences. An authorization was obtained from the head of animal house affiliated with the laboratory of pharmaco-toxicology and pharmacokinetics.

RESULTS

EXTRACTION YIELD OF EXTRACTS

The extraction yield of each extract of *A. conyzoides* is presented in table 1. The extraction yield of the aerial part of *A. conyzoides* was the lowest among the aqueous and hydroethanolic extracts

of the plant. While the flower hydroethanolic extract had the highest yield among the hydroethanolic extracts of the plant, the leaf aqueous extract had the highest yield among the plant's aqueous extracts.

Table 1: Extraction yield of the extracts of *A. conyzoides*.

Solvent	Aqueous maceration				Hydroethanolic maceration			
	AP of plant	Leaf	Stem	Flower	AP of plant	Leaf	Stem	Flower
Extraction yield (%)	9.2	13.1	10.8	11.3	8.3	10.3	13.4	16.3
Texture	viscous	viscous	powder	powder	viscous	viscous	viscous	viscous

Legend: AP = Aerial part

PHYTOCHEMICAL ANALYSIS

Evaluation of the primary metabolites and vitamins

Analysis of *A. conyzoides* extracts revealed the presence of carbohydrates, total proteins, vitamin C and the absence of total lipids and carotenoids in both its aqueous and hydroethanolic extracts as shown in table 2.

Table 2: Qualitative analysis of primary metabolites in *A. conyzoides* extracts.

Metabolites	Test	Aqueous maceration				Hydroethanolic maceration			
		AP of plant	Leaf	Stem	Flower	AP of plant	Leaf	Stem	Flower
Vitamin C	DNPH	+	+	+	+	+	+	+	+
Carotenoids		-	-	-	-	-	-	-	-
Carbohydrates	Molisch test	+	+	+	+	+	+	+	+
	Fehling's test	+	+	+	+	+	+	+	+
Proteins	Biuret test	+	+	+	+	+	+	+	+
	Millon's test	+	+	+	+	+	+	+	+
Lipids	Ethanol/H ₂ O	-	-	-	-	-	-	-	-
	NaOH/Ethanol	-	-	-	-	-	-	-	-

Key : - = negative, + = positive, AP= Aerial part

Qualitative analysis of secondary metabolites

Preliminary phytochemical screening of the aqueous and ethanolic extracts of the aerial part of the plant and its individual parts (leaf, flower and stem) revealed the presence of various bioactive components which include polyphenols, alkaloids, flavonoids, coumarins and tannins though there were some inconsistencies with regard to a few metabolites as shown in Table 3. Saponins and mucilages were found to be present in all the plant extracts except in the aqueous maceration of the aerial part of the plant and stem hydroethanolic extract. Steroids and cardiac glycosides were present only in the aqueous and hydroethanolic extracts of *A. conyzoides* leaves.

Table 3: Qualitative analysis of secondary metabolites of *A. conyzoides* extracts.

Metabolite	Reagents	Aqueous maceration				Hydroethanolic maceration			
		AP of plant	Leaf	Stem	Flower	AP of plant	Leaf	Stem	Flower
Polyphenols	FeCl ₃	+	+	+	+	+	+	+	+
	Lead acetate	+	+	+	+	+	+	+	+
Alkaloids	Wagner	+	+	+	+	+	+	+	+
	Hager	+	+	+	+	+	+	+	+
	Valse-Mayer	+	+	+	+	+	+	+	+
	Tannic acid	+	+	+	+	+	+	+	+
Flavonoids	AlCl ₃	+	+	+	+	+	+	+	+
	NaOH/H ₂ SO ₄	+	+	+	+	+	+	+	+
Tannins	Catechic tannins	+	+	+	+	+	+	+	+
	Gallic tannins	+	+	+	+	+	+	+	+
Saponins	Distilled water	-	+	+	+	+	+	-	+
Mucilage	Absolute ethanol	-	+	+	+	+	+	-	+
Steroids	Acetic anhydride	-	+	-	-	-	+	-	-
Coumarines	HNO ₃	+	+	+	+	+	+	+	+
Oxalate	Ethanoic acid	-	-	-	-	-	-	-	-
Quinones	Sulfuric acid	-	-	-	-	-	-	-	-
Betacyane	NaOH	+	+	+	+	+	+	+	+
Phlobotannins	HCl	+	+	+	+	+	+	+	+
Anthocyanes	NH ₄ OH	-	-	-	-	-	-	-	-
Cardiac glycosides	Glacial acetic	-	-	-	-	-	-	-	-
Resins	Anhydrous acetic	-	-	-	-	-	-	-	-

3.3. Quantitative analysis of plant metabolites

The various extracts of the plant contained carbohydrates, proteins, tannins, flavonoids, flavonols and alkaloids which were then quantified using a UV spectrophotometer. Their concentrations are shown in table 4.

Table 4: Concentration of metabolites in *A. conyzoides* extracts.

Metabolites	AP of Plant		Leaf		Stem		Flower	
	AQ	HET	AQ	HET	AQ	HET	AQ	ET
Carbohydrates (EG/g)	85,57 ± 2,52 ^{fi}	92,45 ± 2,90	157,77 ± 4,25	59,20 ± 2,36 ^{ci,j}	34,68 ± 2,08 ^{fi}	80,06 ± 1,25 ^c	82,63 ± 1,95 ^l	74,46 ± 1,96 ^{cl}
Total proteins (EBSA/g)	561,79 ± 107,30 ^{fi}	767,18 ± 26,89	1059,23 ± 4,68	530,26 ± 25,68 ^{ci,j}	343,85 ± 6,30 ^{fi}	681,79 ± 28,91 ^c	541,79 ± 8,44 ^f	521,03 ± 23,09 ^{cl}
Total polyphenols (GAE/g)	430,27 ± 14,83 ^{fi}	364,93 ± 5,66 ^l	633,87 ± 5,60	356,00 ± 3,12 ^l	288,13 ± 17,22 ^{fi}	366,67 ± 7,49 ^l	422,27 ± 3,78 ^f	408,80 ± 7,11
Total flavonoids (EQ/g)	407,71 ± 16,29 ^f	425,35 ± 3,07	524,37 ± 15,63	361,73 ± 10,37 ^c	181,73 ± 19,28 ^{c,fi}	311,43 ± 7,96 ^{c,fi}	388,29 ± 14,64 ^f	362,02 ± 4,87 ^c
Total flavonols (EQ/g)	637,14 ± 14,38 ^f	590,36 ± 3,44	779,76 ± 19,12	435,95 ± 7,41	261,67 ± 7,90 ^{c,fi}	255,24 ± 22,38	755,71 ± 24,77	451,79 ± 3,11
Total tannins (TAE/g)	170,56 ± 5,70 ^f	128,93 ± 1,65	352,81 ± 2,82	117,66 ± 2,34	153,35 ± 5,30 ^{a,f,k}	131,53 ± 7,00 ^{a,g,l}	171,72 ± 5,74 ^f	139,78 ± 3,93
Total alkaloids (QHE/g)	1090,56 ± ± 34,05	1891,67 ± 35,28	1152,78 ± 65,18	1535,00 ± 131,70 ^b	506,11 ± 39,77 ^{c,fi}	1483,89 ± 30,25 ^{c,e}	899,44 ± 11,71 ^{b,c}	1162,78 ± 35,64 ^{c,e,k}

Key: AP = Aerial Part, AQ = aqueous, HET = hydro ethanol

The results are expressed as mean ± SEM with n = 3; Data analysis was performed using the ANOVA test, followed by Turkey Kramer post hoc multiple comparison test. The differences were considered significant from $a < 0,05$; $b < 0,01$; $c < 0,001$ with the aerial part of the plant, $d < 0,05$; $e < 0,01$; $f < 0,001$ for the leaf; $g < 0,05$; $h < 0,01$; $i < 0,001$ for the stem; $j < 0,05$; $k < 0,01$; $l < 0,001$ for the flower

Total carbohydrate content

The distribution of total carbohydrates in the different plant extracts shows that in the aqueous extracts, the leaves contain the highest concentration in total carbohydrates with a significant difference compared to the other parts of the plant as well as the aerial part of the plant as a whole. with a $p\text{-value} < 0.001$. In the hydro-ethanolic extracts, we note the preponderance of the aerial part of the plant compared to the other parts which are the leaves with a significant difference with a $p\text{-value} < 0.001$. Among the aqueous extracts of *A. conyzoides*, the stem has the lowest content in carbohydrates meanwhile among the hydroethanolic extracts of the plant, the flowers have the lowest content in soluble carbohydrates.

Total protein content

The distribution of total proteins in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaves present the highest concentration in comparison with the entire aerial part of the plant, the flowers and the stems with a $p\text{-value} < 0.001$. We note however in the hydro-ethanolic extracts, a higher concentration of proteins in the aerial part of plant in comparison with the leaves, stems and flowers. The total protein content in the flower ethanolic extract is low as compared to the other ethanolic extracts of the plant while the stem aqueous extract is the lowest in comparison to the other aqueous extracts of the plant.

Total polyphenols content

The distribution of total polyphenols in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaves presented the highest concentration in comparison with the aerial part of the plant, flower and stem with a $p\text{-value} < 0.001$. However, we note that in the hydro-ethanolic extracts, their concentration was highest in the flowers as compared to the leaf, stem and the aerial part of the plant with a $p\text{-value} < 0.001$. The stem aqueous extract has a significantly low content in polyphenols amongst the aqueous *A. conyzoides* extracts while the leaf of *A. conyzoides* ethanolic extract has the lowest concentration in polyphenols as compared to other ethanolic extracts of the plant.

Total flavonoid content

The distribution of total flavonoids in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaf presents the highest concentration in comparison with the whole aerial part of the plant, flower and stem with a $p\text{-value} < 0.001$. Meanwhile, we notice

that in the hydro-ethanolic extracts, total proteins are highest in the aerial part of the plant in comparison with the leaves, stems and flowers with a p-value of < 0.001 . Generally, both the aqueous and ethanolic extracts of the stem of *A. conyzoides* presents the lowest content in flavonoids as compared to the leaves, flowers and aerial part of the plant's aqueous and ethanolic extracts respectively.

Total flavonols content

The distribution of total flavonols in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaves present the highest concentration in comparison with the whole aerial part of the plant, the flowers and stem with a p-value < 0.001 whereas, in the hydro-ethanolic extracts, we observe that the aerial part of the whole plant has the highest concentration in flavonols in comparison with the leaves, stems and flowers with a p-value of < 0.001 . We also notice that the stem aqueous and ethanolic extracts have the lowest in flavonols in comparison to the aqueous and ethanolic extracts of the leaves, flowers and aerial part as a whole of *A. conyzoides* respectively.

Total tannins content

The distribution of total tannins in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaves present the highest concentration in comparison with the entire aerial part of the plant, flowers and stems with a *p-value* < 0.001 whereas, we note in the hydro-ethanolic extracts, a slightly higher concentration in the flowers in comparison to the leaves, stems and the aerial part of the plant. We equally note that total tannins content among the aqueous extracts of *A. conyzoides* is lowest in the stem meanwhile among the ethanolic extracts of the plant, it is lowest in the leaves.

Total alkaloid content

The distribution of total alkaloids in the different extracts of the plant shows that in the aqueous extracts obtained by maceration, the leaves present the highest concentration in comparison to the entire aerial part of the plant, flowers and stems with a p-value < 0.001 . However, in the hydro-ethanolic extracts, the aerial part of the plant in comparison to the leaves, stems and flowers had the highest concentration in alkaloids. Also, we note that, the total alkaloid content

among the aqueous extracts of *A. conyzoides* is lowest in the stem meanwhile among the ethanolic extracts of the plant, it is lowest in the flowers.

DISCUSSION

Phytochemicals are naturally occurring substances found in plants which provides health benefits. These are known as secondary metabolites and may often be created by modified synthetic pathways from primary metabolite or share substrates of primary metabolite origin [25]. *Ageratum conyzoides* L., Asteraceae, is an annual herbaceous plant with a long history of traditional medicinal uses in several countries of the world. Earlier studies conducted by researchers such as Fatima *et al.* (2016), Odeleye *et al.* (2014) and Onerijiofi and Isola (2019) on *A. conyzoides* [1,26,27] had indicated the presence of phytochemicals in the *A. conyzoides*. The minor differences observed between the phytochemicals detected in this study and those of earlier authors could have arisen from differences in extraction methods, nature, concentration and polarity of solvent used, environmental factors and time of harvest [28].

This study revealed the presence of phytochemicals in all aerial parts of *A. conyzoides* such as polyphenols, alkaloids, flavonoids, coumarins and tannins. Most of these phytochemicals are known to possess pharmacological applications. For instance, flavonoids are a large class of natural compounds that are widely found in the plant kingdom found to possess a diverse range of bioactivities, such as antioxidation, antihyperlipidemic, antifatigue, antiaging, and atherosclerosis-prevention activities [29]. Meanwhile, alkaloids possess antimicrobial activity which could explain why the plant is used for the traditional treatment of infectious diseases [30]. In addition, saponins, tannins and terpenoids have been reported to have dramatic physiological antibacterial, antifungal and antioxidant properties [31]. In all, this study shows that the extracts of *A. conyzoides* are extensively rich in secondary metabolites thereby justifying its use in the treatment of several ailments by traditional practitioners.

The quantitative phytochemical analysis of the extracts of *A. conyzoides* revealed that the aqueous maceration of the leaves of *A. conyzoides* had the highest content in most of the metabolites. It contained $157,77 \pm 4,25$ mg EG/mg of total carbohydrate content, $1059,23 \pm 4,68$ μ g BSA of total protein content, $633,87 \pm 5,60$ mg GAE/g of total phenolic content, $524,37 \pm 15,63$ mg EQ/g of total flavonoid content, $779,76 \pm 19,12$ mg EQ/g of total flavonols content and

352,81 ± 2,82 mg TAE/gm of total tannins content. However, an exception was observed with the total alkaloid content which was highest in the hydroethanolic maceration of the aerial part of the plant at a concentration of 1891,67 ± 35,28 mg QHE/g. In this study, the total phenolic content in the aqueous and ethanolic maceration of the leaves was similar to what was obtained in a study conducted by Pawan Kumar *et al.* (2012) [32] on the acetone and hexane extracts of the leaves of *A. conyzoides* and higher than what was found by Puro *et al.* (2018) [33] in the methanolic leaf extract of the plant. However, the amount of total flavonoid content in the aqueous and ethanolic extracts of the leaves in this study was much lower than what was obtained by Pawan Kumar *et al.* 2012 [32]. The differences are could be related to the difference in the extraction solvents. Also, due to the fact that the leaf extracts contained the highest contents in most secondary metabolites, this could explain why it is the most commonly applied part of *A. conyzoides*.

CONCLUSION

In conclusion, the present study shows that *A. conyzoides* aqueous and hydro-ethanolic maceration extracts of the leaves, stem and flowers as well as the aerial part as a whole contain polyphenols, saponins, alkaloids, flavonoids, catechic tannins, gallic tannins, coumarins, quinones and phlobatannins. The present work also shows that most primary and secondary metabolites are predominantly found in aqueous maceration of the leaves of *A. conyzoides* compared to the aqueous and ethanolic extracts of all other parts of the plant as well as its aerial part as a whole. This study could serve as a benchmark for further pharmacological studies on the plant that may be harnessed for drug development in the future.

REFERENCES

1. Mary APF, Giri DRS. PHYTOCHEMICAL SCREENING AND GC-MS ANALYSIS IN ETHANOLIC LEAF EXTRACTS OF AGERATUM CONYZOIDES (L.). WJPR. 5(7).
2. Uddin G, Rauf A. Phytochemical Screening, Antimicrobial and Antioxidant Activities of Aerial Parts of *Quercus robur* L. MEJMPR. 2012;1(1):1–4.
3. Shaikh JR, Patil M. Qualitative tests for preliminary phytochemical screening: An overview. Int J Chem Stud [Internet]. 2020 Mar 1 [cited 2021 Nov 25];8(2):603–8. Available from:

<http://www.chemijournal.com/archives/?year=2020&vol=8&issue=2&ArticleId=8834&si=false>

4. Al-Brashdi AS, Al-Ariymi H, Al Hashmi M, Khan SA. Evaluation of antioxidant potential, total phenolic content and phytochemical screening of aerial parts of a folkloric medicine, *Haplophyllum tuberculatum*. *J Coast Life Med* [Internet]. 2016 Apr [cited 2024 Mar 3];4(4):315–9. Available from: <http://www.jclmm.com/qk/20164/13.pdf>
5. Leke WN, Brown JK, Lighthart ME, Sattar N, Njuaem DK, Kvarnheden A. *Ageratum conyzoides*: A host to a unique begomovirus disease complex in Cameroon. *Virus Res* [Internet]. 2012 Jan 1 [cited 2024 Mar 2];163(1):229–37. Available from: <https://www.sciencedirect.com/science/article/pii/S0168170211003923>
6. Patel F, Modi N. Antioxidant activity and FTIR analysis of *Ageratum conyzoides* L. *Int J Bot Stud*. 2021 Oct 11;6(5):1152–61.
7. Okunade AL. *Ageratum conyzoides* L. (Asteraceae). *Fitoterapia* [Internet]. 2002 Feb [cited 2022 Nov 6];73(1):1–16. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0367326X01003641>
8. Chauhan A, Rijhwani DS. A Comprehensive Review on Phytochemistry of *Ageratum conyzoides* linn. (Goat weed). *IJETMAS*. 2015;3.
9. Amadi B, Majesty D, Nnabugwu A. Chemical profiles of leaf, stem, root and flower of *Ageratum conyzoides*. *AJPSKY*. 2012 Jan 1;2:428–32.
10. Chandra Jagetia G, Shirwaikar A, Rao SK, Prasad MB. Evaluation of the radioprotective effect of *Ageratum conyzoides* Linn. extract in mice exposed to different doses of gamma radiation. *JPP* [Internet]. 2003 [cited 2024 Mar 2];55:1151–8. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1211/0022357021576>
11. Galati EM, Miceli N, Taviano MF, Sanogo R, Raneri E. Anti-inflammatory and Antioxidant Activity of *Ageratum conyzoides*. *Pharm Biol* [Internet]. 2001 Jan 1 [cited 2024 Mar 2];39(5):336–9. Available from: <https://doi.org/10.1076/phbi.39.5.336.5891>
12. Pougoue KJ. Investigation of the antiulcer, and toxicity of the stem bark extract of *Ficus thonningii* Blume Linn. (Moraceae) on Wistar rats. university of Yaounde 1; 2017.
13. Tembe EF, Pougoue KJ, Borgia N, Ngoupayo J, Gatsing D, Tomkins P, et al. Phytochemical screening and in vivo evaluation of antiulcer properties of secondary metabolites in aqueous extracts of *Ficus thonningii* Blume tested on Wistar rats. *Int J Biol Chem Sci*. 13(1). 2019;475–92.
14. Sofowora A. *Medicinal Plants and Traditional Medicine in Africa*. Ibadan, Nigeria: Spectrum Books Ltd; 1993. 191–289 p.
15. Balamurugan V. A guide to phytochemical analysis. *Int J Adv Res Innov Ideas Educ*. 2019;5(1):10.

16. Savithrama N., Rao M.L., Suhrulatha D. Screening of Medicinal Plants for Secondary Metabolites. Middle East J Sci Res. 2011;8(3):579–84.
17. Thomas W, Dutcher RA. The colorimetric determination of carbohydrates in plants by the picric acid reduction method I. The estimation of reducing sugars and sucrose. J Am Chem Soc [Internet]. 1924 [cited 2022 Jun 15];46(7):1662–9. Available from: <https://pubs.acs.org/doi/abs/10.1021/ja01672a015>
18. Lowry OH, Rosebrough NJ, Farr AL, Randall R J. Protein measurement with the Folin phenol reagent. J Biol Chem. 193(1). 1951;265–75.
19. Galvão MAM, Arruda AO, Bezerra ICF, Ferreira MRA, Soares LAL. Evaluation of the Folin-Ciocalteu Method and Quantification of Total Tannins in Stem Barks and Pods from *Libidibia ferrea* (Mart. ex Tul) L. P. Queiroz. Braz Arch Biol Technol [Internet]. 2018 [cited 2022 Apr 22];61(0). Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1516-89132018000100431&lng=en&tlng=en
20. Amari NO, Bouzouina M, Berkani A, Lotmani B. Phytochemical screening and antioxidant capacity of the aerial parts of *Thymelaea hirsuta* L. Asian Pac J Trop Dis [Internet]. 2014 Apr 1 [cited 2024 Mar 3];4(2):104–9. Available from: <https://www.sciencedirect.com/science/article/pii/S2222180814603248>
21. Kosalec I. Quantitative analysis of the flavonoids in raw propolis from northern Croatia. Acta Pharm. 54(1). 2004;65–72.
22. Miliuskas G, Venskutonis, P.R, Van Beek, T. A. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. Food Chem. 2004;85:235–7.
23. Graham, H, Thomas L. Color reaction of alkaloids with dichromate-sulfuric acid and its use for quantitative assay. J Pharm Sci. 1961;50:901–4.
24. Ali-Rachedi F, Meraghni S, Touaibia N, Mesbah S. Analyse quantitative des composés phénoliques d'une endémique algérienne *Scabiosa Atropurpurea* sub. *Maritima* L. Bull Société R Sci Liège. 2018;
25. Silva GOD, Abeyundara AT, Aponso MMW. Extraction methods, qualitative and quantitative techniques for screening of phytochemicals from plants. AJEONP. 2017;5(2):29–32.
26. Odeleye OP, Oluyeye JO, Aregbesola OA, Odeleye PO. Evaluation of preliminary phytochemical and antibacterial activity of *Ageratum conyzoides* (L) on some clinical bacterial isolates. IJES. 2014;3(6):1–5.
27. ENERIJIOFI, KE, ISOLA, OB. Preliminary Phytochemical screening and invitro antibacterial activities of aqueous and ethanol extracts of *Ageratum conyzoides* L. Leaf, Stem, Flower and Root on some Bacterial isolates associated with Diarrhoea. Nig J Pure Appl Sci. 2019;32(2):3480–9.

28. Kunle OF, Egharevba O.H., Ahmadu PO. Standardization of herbal medicines - A review. *Int J Biodivers Conserv* [Internet]. 2012 Mar [cited 2023 Sep 2];4(3). Available from: <http://www.academicjournals.org/ijbc/abstracts/abstracts/abstracts2012/March/Kunle%20et%20al.htm>
29. Lin Z, Lin Y, Shen J, Jiang M, Hou Y. Flavonoids in *Ageratum conyzoides* L. Exert Potent Antitumor Effects on Human Cervical Adenocarcinoma HeLa Cells *In Vitro* and *In Vivo*. *Biomed Res Int* [Internet]. 2020 May 4 [cited 2024 Mar 2];2020:e2696350. Available from: <https://www.hindawi.com/journals/bmri/2020/2696350/>
30. Uddin G, Rauf A. Phytochemical screening and biological activity of the aerial parts of *Elaeagnus umbellata*. *Sci Res Essays*. 2012;7(43).
31. Munazir M, Qureshi R, Munir M. PRELIMINARY PHYTOCHEMICAL SCREENING OF ROOTS AND AERIAL PARTS OF LEPTADENIA PYROTECHNICA. *Pak J Bot*. 2015;47(2):659–64.
32. Pawan Kumar V, Raina R, Prawez S, Dar M. Quantitative Analysis of Total Phenolic, Flavonoids and Tannin Contents in Acetone andn-hexane Extracts of *Ageratum conyzoides*. *IntJChemTech Res*. 2012;4(3):pp 996-999.
33. Puro K, Mazumder MU, Khazeo P, Jyrwa R, Jamir N, Sailo L. Qualitative and quantitative analysis of phytochemicals of crude extracts of *Ageratum conyzoides* L. leaves. *MSC*. 2018;178:164–8.