

What are the particularities of hepatitis B virus awareness, among care workers and medical Students in an endemic country?

Abstract:

The hepatitis B virus is well known for its contagiousness, especially among healthcare workers due to their direct contact with blood products, affecting both healthcare personnel and patients. This highlights the importance of our study, which sheds light on the underestimation of the seriousness of this virus's spread and emphasizes the significance of vaccination by sampling students starting from the third year, interns, and residents.

Aims: To assess knowledge about hepatitis B virus (HBV), its infectious risk, vaccination against HBV, and the vaccination coverage rate within our population.

Study design: We conducted a descriptive cross-sectional study

Place and duration of the study: The study was conducted from October 2022 to July 2023, in Hassan II regional hospital in Agadir, Morocco.

Methodology: We were using an anonymous questionnaire that was completed by 267 participants as our foundation.

Results: Less than half (48.3%) were aware that complete vaccination requires at least 3 doses, with only 27.3% reporting being fully vaccinated. 40% were aware of the existence of anti-HBV Ig. Regarding practical attitudes towards Blood borne exposure incident, 25% of our study population had been exposed, with causes of non-reporting including underestimation of risk in 59% of cases and administrative difficulties in 49% of cases.

Conclusion: Our study highlighted the lack of information regarding basic HBV knowledge and the severity of its transmission, leading to a very low vaccination rate among healthcare personnel. This underscores the importance of thorough awareness campaigns targeted at those affected.

Keywords: *Hepatitis B vaccine, medical students, health care workers, accidental exposure to blood.*

Introduction:

Chronic hepatitis B virus (HBV) infection affects about 296 million people worldwide and is the leading etiology of cirrhosis and liver cancer globally; other medical complications also include acute flares and extrahepatic manifestations (1)

In the Moroccan context, our country is classified as having a moderate level of endemicity. According to the latest national seroprevalence survey on viral hepatitis conducted in 2019 by the National Ministry of Health, the prevalence of hepatitis B in general population was 0.7%[2,3,12-14].

Healthcare personnel, including medical students, represent a population at high risk of contamination by the virus, particularly through contact with contaminated materials and blood exposure incidents (BEI). A national multicenter study revealed a higher prevalence of 2.2% among physicians[2], this suggests several reflections: is it due to insufficient awareness despite the availability of the vaccine? Or is it a negligence of the infectious risk?

Materiel and Methods:

-We conducted a descriptive cross-sectional study from October 2022 to July 2023, including medical students from the third to the seventh year of medical studies (externs), interns, and residents from all medical and surgical specialties, practicing at Hassan II Regional Hospital in Agadir.

-We calculated the sample size using a prevalence of vaccinated professionals in Cameroon at 18%, an alpha risk of 5%, and a precision of 0.05. The minimum sample size required was 227[4-6].

-Data collection was done using an anonymous online questionnaire, and statistical analysis was performed using Excel and Jamovi software. Participants were asked for consent before filling out the questionnaire.

-Our questionnaire consisted of 25 questions, organized into three main chapters:

- ✓ Sociodemographic and professional data,
- ✓ Assessment of knowledge and practical attitudes related to the risk of HBV infection, especially during a Blood Exposure Incident (BEI),
- ✓ Evaluation of knowledge about HBV vaccination (adverse effects and vaccination contraindications), verification of vaccination status, suggestions and proposals to improve the awareness and its vaccination among hepatitis B virus

-To assess the level of general knowledge about the virus, its infectious risk, and vaccination, we calculated the average of correct responses. Knowledge level was noted "Good" if the average of correct answers was 75% or higher; "Moderate" if correct responses were between 50% and 75%; and "Low" if the knowledge level did not exceed 50%.

Results :

-Our population consisted of 267 individuals who completed the questionnaire, with a median age of 23.7 years ranging from 19 to 36 years. Most of our participants were female (65%), and 35% were male, resulting a sex ratio (M/F) of 0.5. Among the respondents, 152 were medical externs, accounting 57% of the sample, while 23.2% (n=62) were residents. There was a lower representation of interns, with 52 participants accounting for 19.5% (Figure 1)




Figure 1: Distribution of the target population according to level of education

Knowledge about HBV:

-The first question focused on the virus transmission routes: we observed that the contamination through blood, sexual fluids, and vertical routes were mostly mentioned by 98%, 93%, and 89% respectively. However, the salivary route was only known by 23.6% of participants, primarily among interns, and 29% responded that the virus has an oro-fecal transmission.

-Overall, the level of knowledge about HBV transmission routes was good.

Knowledge about HBV Vaccination:

-In this section, we asked 4 questions regarding:

- ✓ The World Health Organization's recommendation of this vaccination for healthcare personnel,
- ✓ The necessity of 3 doses for complete vaccination,
- ✓ The vaccine's efficiency,
- ✓ The existence of immunoglobulins against HBV.

-We concluded that less than half (48.3%) knew that complete vaccination requires at least 3 doses, and only 40% were aware of the existence of anti-HBV immunoglobulins, indicating a low level of knowledge regarding these two points. 66.3% (n=177) responded that vaccination is mandatory for medical professionals, and 77.5% (n=207) knew that HBV vaccination is effective.

-Regarding knowledge about vaccine adverse effects: the two main answers were allergic reaction and multiple sclerosis, respectively at 35% and 34% (Figure 2).

-For vaccine contraindications, we included among the choices: pregnancy, Human Immunodeficiency Virus (HIV) co infection, multiple sclerosis, history of severe allergic reaction, and elderly status. The majority response was a history of severe allergic reaction at 38%, followed by HIV coinfection at 27% (Figure 3).

-The level of knowledge for both vaccine adverse effects and its contraindications was poor, among our population study.




Figure 2: Knowledge about Vaccine Adverse Effects




Figure 3: Respondents' Knowledge about Vaccine Contraindications

Vaccination Status

-Out of the total 267 individuals surveyed, only 27.3% (n=73) reported being fully vaccinated, while 38% were incompletely vaccinated and 25% were not vaccinated at all. Among those vaccinated, vaccination was done through the national immunization program in 56.3% of cases, during a mass campaign in 32.5% of cases, and personally for 10.7%.

-The main reason for non-vaccination among the unvaccinated was negligence in 82% of respondents (142 individuals), followed by fear of adverse effects in 9.4% (16 individuals). The remaining 13 respondents were distributed among vaccine cost (6.4%) and HBV infection concerns (1.2%) (Figure 4).




Figure 4: Reasons for Abstaining from Vaccination among the Unvaccinated

Practical Attitudes during Blood Exposure Incidents (BEIs):

-Regarding practical attitudes during a BEI, which represents the main risk of contamination among health care professional: we noticed that 25% of our study population had already been exposed; residents were the most exposed accounting for 48.4% of cases? However, only 25% of individuals who experienced a BEI reported it to the occupational health physician.

-The reasons for non-reporting were attributed to underestimating the risk in 59% of cases and administrative difficulties in 49% of cases.

Discussion:

-HBV is a DNA virus, posing a serious public health problem on a global scale. Its risk lies in the progression towards severe and deadly complications. 5 to 10% of infected adults become chronic carriers, with about 20% developing cirrhosis or HCC due to continuous virus replication.

-The virus reservoir is human, and its transmission occurs interpersonally. In infected individuals, the virus is present in blood and its derivatives, semen, vaginal secretions, breast milk, saliva, and to a lesser extent, tears, sweat, and urine.

-Healthcare professionals are the most exposed to HBV infection due to direct contact with infected patients' fluids and medical equipment. Understanding the infectious risk of HBV and the importance of vaccination is crucial for reducing infection rates.

-In our study, the level of knowledge regarding HBV transmission routes was good, the majority recognized the blood (98%), sexual (93%), and vertical (89%) routes. However, a minority mentioned the possibility of oro-fecal transmission. These figures are almost similar to those reported in the study conducted by Fortes Déguénonvo et al. in Senegal [7].

-Hepatitis B immune globulin is an injection of immunoglobulin G (IgG) to protect against infection and disease progression. The administration of immunoglobulins may be used in certain situations, such as post-exposure prophylaxis after contact with contaminated blood or in newborns of HBV-positive mothers to prevent vertical transmission. These immunoglobulins provide temporary protection against infection and are often used in combination with the hepatitis B vaccine to boost short-term immunity. However, it is important to note that the effectiveness of immunoglobulins decreases over time, and complete vaccination remains necessary to ensure long-term protection against HBV.

-The World Health Organization (WHO) states that there is no causal link between the HBV vaccine and demyelinating neurological conditions such as multiple sclerosis, autoimmune diseases, diabetes, and chronic fatigue syndrome. The only confirmed adverse effect of the vaccine is the occurrence of mild allergic reactions within 24 hours of injection. Regarding contraindications, the WHO confirms that the only contraindication to the vaccine is a history of severe allergic reaction. Neither pregnancy, HIV co-infection, age, nor other comorbidities are contraindications to the vaccine, which is considered to have high safety and efficacy [8].

-Regarding BEI, one-quarter of our target population had already been victim of at least one blood exposure incident. A study conducted at the hospital in Casablanca in 1998 targeting healthcare workers found that 81% of participants had been exposed to blood at least once [2]. Thus, the knowledge about post-exposure management is crucial.

-Regarding vaccination, only 27.3% were fully vaccinated in our study, which is a very low rate. In Nepal, in 2020, 60.8% of students were vaccinated, with 60.9% of them receiving at least 3 complete doses, but in Saudi Arabia, 38% had received all 3 doses of the vaccine; and in Cameroon, only 18% of participants were fully vaccinated, according to their findings [4,9,10].

-Concerning abstention among the unvaccinated, the predominant reason was negligence in 82% of cases, a common argument with the study by Altamimi et al. conducted at public universities in Riyadh [11]. This underscores the importance of collective awareness about the complete vaccination for healthcare professional, and the necessity of ensuring widespread availability of the vaccine in Moroccan healthcare, and regulating it through Moroccan law.

-Perspectives: We intend to extend our study to a larger sample and throughout the country, for a more thorough and effective sensitization.

Conclusion:

- Our study demonstrated a lack of information regarding basic HBV knowledge, particularly a significant unawareness of the existence of hepatitis B immunoglobulins.
- The fear of developing multiple sclerosis remains a concern even among medical personnel, highlighting the need to increase awareness about vaccine adverse effects and contraindications during theoretical training and vaccination campaigns.
- The risk of HBV transmission is underestimated during a Blood Exposure Incident (BEI), and administrative difficulties hinder medical personnel from reporting such incidents.
- There is a low rate of complete vaccination against hepatitis B among the studied population, underscoring the need for increased awareness and initiatives to improve vaccination coverage.

Consent:

All the participants in our study were asked for written consent before filling out the questionnaire

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References:

1. Hsu, Y.-C., Huang, D.Q., Nguyen, M.H., 2023. Global burden of hepatitis B virus: current status, missed opportunities and a call for action. *Nat Rev Gastroenterol Hepatol* 20, 524–537.
2. Laraqui O, Laraqui S, Laraqui S, Tripodi D, Chahed Ouazzani L, Caubet A, et al. Évaluation des connaissances, attitudes et pratiques sur les hépatites virales B et C en milieu de soins au Maroc: *Santé Publique*. 11 août 2009;Vol. 21(3):271-86.
3. Enquête Nationale de séroprévalence des hépatites virales 2019.
4. Noubiap JJN, Nansseu JRN, Kengne KK, Tchokfe Ndoula S, Agyingi LA. Occupational exposure to blood, hepatitis B vaccine knowledge and uptake among medical students in Cameroon. *BMC Med Educ*. 8 nov 2013;13(1):148.
5. Cochran WG (William G. Sampling techniques [Internet]. New York, NY : Wiley; 1977 [cité 2 mai 2024]. 462 p. Disponible sur: http://archive.org/details/samplingtechniqu0000coch_t4x6
6. Bartlett JE, Kotrlík JW, Higgins CC. *Organizational Research: Determining Appropriate Sample Size in Survey Research*.
7. Fortes Déguénonvo L, Massaly A, Ngom Gueye N, Diallo Mbaye K, Cisse Diallo V, Lakhe N, et al. Assessment of Knowledge, Attitudes and Practices of Medical Students Regarding Hepatitis B Infection at a Private University of Medicine in Senegal. *J Infect Dis Epidemiol* [Internet]. 19 déc 2019 [cité 29 avr 2024];5(6). Disponible sur: <https://www.clinmedjournals.org/articles/jide/journal-of-infectious-diseases-and-epidemiology-jide-5-103.php?jid=jide>
8. Gerlich WH. Medical Virology of Hepatitis B: how it began and where we are now. *Virology*. déc 2013;10(1):239.

9. Shrestha DB, Khadka M, Khadka M, Subedi P, Pokharel S, Thapa BB. Hepatitis B vaccination status and knowledge, attitude, and practice regarding Hepatitis B among preclinical medical students of a medical college in Nepal. *Chemin I, éditeur. PLOS ONE*. 23 nov 2020;15(11):e0242658.
10. Alhowaish MA, Ali Alhowaish J, Hamoud Alanazi Y, Mana Alshammari M, Saeid Alshammari M, Ghadeer Alshamari N, et al. Knowledge, attitudes and practices toward prevention of hepatitis B virus infection among medical students at Northern Border University, Arar, Kingdom of Saudi Arabia. *Electron Physician*. 25 sept 2017;9(9):5388-94.
11. Altamimi A, Alqahtani T, Ahmed J, Aldosari L, Alzahrani M, Alotaibi G, et al. Knowledge, awareness, and vaccination compliance of hepatitis B among medical students in Riyadh's governmental universities. *J Fam Med Prim Care*. 2021;10(1):485.
12. Obi RK, Oke BO, Anjorin AA, Bidmos IK, Salu OB, Omilabu SA. Hepatocarcinogenic Properties of Hepatitis B Virus. *Int. J. Trop. Dis. Health*. [Internet]. 2016 Apr. 26 [cited 2024 May 25];16(2):1-12. Available from: <https://journalijtdh.com/index.php/IJTDH/article/view/353>
13. Chaudhry K, Kaur G, Kumar P. The Association of Hepatitis B and Polymyositis: Rare but Real. *Asian J. Res. Infect. Dis*. [Internet]. 2023 Apr. 13 [cited 2024 May 25];13(1):1-4. Available from: <https://journalajrid.com/index.php/AJRID/article/view/254>
14. Yuen MF, Chen DS, Dusheiko GM, Janssen HL, Lau DT, Locarnini SA, Peters MG, Lai CL. Hepatitis B virus infection. *Nature reviews Disease primers*. 2018 Jun 7;4(1):1-20.