

COMPLETION OF WEAKLY SIGN SYMMETRIC P_0 -MATRIX PROBLEM FOR 5×5 MATRICES SPECIFYING diJABB_121114S OF ORDER 5 WITH UP TO 5 ARCS

ABSTRACT

An $n \times n$ matrix is a weakly sign symmetric matrix if the off-diagonal elements have the property that if the entry in row i and column j is non-zero, then the entry in row j and column i must have same sign or zero. A diJABB_121114 D has a Wss P_0 -matrix completion if every partial weakly sign symmetric P_0 -matrix that describes D can be extended to a complete weakly sign symmetric P_0 -matrix. This paper investigates the problem of completing weakly sign symmetric P_0 -matrices. It demonstrates that partial matrices representing all directed JABB_121114s of order 5 with edge strengths from 0 to 5 can indeed be completed to a weakly sign symmetric P_0 -matrix. Moreover, we established diJABB_121114 characteristics that the partial Wss P_0 -matrices specifying diJABB_121114s of order 5 with up to 5 arcs which have a clique and are cyclic or acyclic have zero completion into a Wss P_0 -matrix. Insights gained from this class of matrix could be applied to fill gaps in data surveys, and business analytics by analysing complex relationships, allocating resources, network modelling, optimizing processes and managing risks.

Keywords: cyclic diJABB_121114s; acyclic diJABB_121114s; clique diJABB_121114s; Matrix completion; DiJABB_121114; weakly sign symmetric P_0 -matrix.

1. INTRODUCTION

Completion of Wss P_0 -matrix for 4×4 matrices has been explored by [1,2] using diJABB_121114s of order 4 with 4 arcs. However, the case of diJABB_121114s of order 5×5 has not been investigated. In this study we determine Wss P_0 -matrix completion of diJABB_121114s of order 5 with up to 5 arcs. A P_0 -matrix A is classified as a weakly sign symmetric P_0 -matrix if $a_{ij} a_{ji} \geq 0$ for all i and j [3]. A sub matrix that has no unspecified entry is said to be fully specified according to [6,8]. Let α be a subset of $N = 1, 2, \dots, n$. the principal submatrix obtained by deleting all columns and rows not in α from A is denoted as $A(\alpha)$. Similarly, $P(\alpha)$ represents the principal sub pattern obtained from the pattern P by removing all positions where the first and second coordinates are not in α . A principal minor of A is the determinant of a principal sub-matrix [5]. A square matrix A is $n \times n$, with equal rows and columns. A partial matrix is an array of numbers with some specified entries while others unspecified. A partial P_0 -matrix has non-negative principal submatrices. A fully specified submatrix of A has all entries defined [10,11].

2. PRELIMINARIES

Basic concepts in linear algebra, group theory and JABB_121114 theory that are commonly used in Wss P_0 -matrix completion problems are defined in the section below.

Definition: 2.1 JABB_121114s and diJABB_121114s is used in matrix completion of various classes of matrices. A JABB_121114, denoted as $G = (V_G, E_G)$ comprises a finite non-empty set of positive integers V_G , where vertices are the members, along with a set of unordered pairs $\{u, v\}$ of vertices called edges. A null JABB_121114 is a JABB_121114 devoid of edges [7,8].

Definition: 2.2 A diJABB_121114 $D = (V_D, E_D)$ is like a JABB_121114 G but includes directed edges (u, v) where u is the start vertex and v is the end vertex. A diJABB_121114 with cycles is termed a cycle diJABB_121114; without cycles, it's acyclic. The order of a diJABB_121114 is its vertex count, and its size is the number of arcs. A clique in a diJABB_121114 contains all possible arcs between vertices [5].

Definition: 2.3 Matrix completion involves identifying feasible positions in matrices using patterns. Patterns like symmetric pairs (i, j) and (j, i) are crucial for $n \times n$ submatrices, focusing on diagonal elements. Symmetric properties in weakly sign symmetric P_o -matrices facilitate problem-solving in completing matrices, utilizing specified entries corresponding to outlined patterns [4].

Definition: 2.4 A pattern D is considered a permutation similar to a pattern B if there exists a permutation ϕ of $\{1, 2, \dots, n\}$ such that B is formed by mapping the pairs (i, j) in D to $\phi(i), \phi(j)$ [6].

Lemma: 2.5 Weakly sign symmetric P_o -matrices exhibit closure under similarity transformations by permutations.

Weakly sign symmetric P_o -matrices, due to their closure under permutation similarity, allow diJABB_121114 relabeling. If a P_o -matrix A has non-negative elements a_{ij} and there exists a permutation matrix P such that PAP^T is sign symmetric, then A is considered a weakly sign symmetric P_o -matrix [1,2].

Definition: 2.6 Let A be a Wss P_o - matrix. Then

(i) If P is a permutation matrix, then PAP^T is a Wss P_o - matrix

(ii) Any principal sub-matrix of A is Wss P_o - matrix. The set of Wss P_o - matrices is closed under permutation similarity and left and right diagonal multiplication.

Theorem 2.7 A permutation matrix P is obtained by interchanging row on the identity matrix. The permutation matrix P is given by PAP^T . This is represented on the diJABB_121114 by relabeling the vertices of the diJABB_121114 [13].

Proposition: 2.8 Every $n \times n$ partial Wss P_o - matrix with all unspecified off diagonal entries has Wss P_o -matrix completion.

Theorem: 2.9 Let $A = [a_{ij}]$ be a partial Wss P_o -matrix and $\alpha = \{i: a_{ij}\}$ is specified. If the principal partial submatrix $A(\alpha)$ Of A has a Wss P_o -matrix completion, then A has Wss P_o -completion [12].

3. ANALYSIS OF 5x5 MATRICES SPECIFYING DIJABB_121114S WITH 5 VERTICES

In the process of constructing a partial Wss P_o - matrix, we use a diJABB_121114 where vertices correspond to diagonal entries d_{ii} . This class of matrices allows 0 as an entry, and by definition, diagonal entries are non-negative (≥ 0). DiJABB_121114s are utilized to create partial Wss P_o -matrices, where specific entries were known denoted by a_{ij} corresponding to existing arcs in the diJABB_121114, and other entries are unspecified denoted by X_{ij} representing missing arcs. diJABB_121114s with 5 vertices and 0 to 5 arcs are considered. In this paper we take our d_{ii} and a_{ij} to be 1 respectively.

DiJABB_121114 D of order 5 without an arc; consider the diaJABB_121114 $D = \{(1,1) (2,2), (3,3), (4,4), (5,5)\}$ with 5 vertices and no arc given by:




Figure 1
Digraph
D of
order 5

The partial matrix that specifies the diJABB_121114 D is

$$A = \begin{pmatrix} d_{11} & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & d_{22} & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & d_{33} & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & d_{44} & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix}$$

By definition of partial Wss P_o - matrix $d_1 \geq 0, d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0$.

For $d_{ii} = 1$, $a_{ij} = 1$, then the partial matrix becomes $A = \begin{pmatrix} 1 & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & 1 & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & 1 & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & 1 & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix}$. Next we

compute the principal minors.

Det (1,2) = $d_{11}d_{22} - x_{12}x_{21}$. Setting the unspecified entry to zero, and $d_{ii} = 1$, i.e., Det (1,2) = $1 - 0 = 1 > 0$. Similarly,

Det (1,3) = $d_{11}d_{33} - x_{13}x_{31}$, Det (1,4) = $d_{11}d_{44} - x_{14}x_{41}$

Det (1,5) = $d_{11}d_{55} - x_{15}x_{51}$, Det (2,3) = $d_{22}d_{33} - x_{23}x_{32}$, Det (2,4) = $d_{22}d_{44} - x_{24}x_{42}$

Det (2,5) = $d_{22}d_{55} - x_{25}x_{52}$, Det (3,4) = $d_{33}d_{44} - x_{34}x_{43}$, Det (3,5) = $d_{33}d_{55} - x_{35}x_{53}$

Det (4,5) = $d_{44}d_{55} - x_{45}x_{54}$.

TABLE 1: Determinants of 2×2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	Det A (1,2) = $d_{11}d_{22} \geq 0$.
A (1,3)	Det A (1,3) = $d_{11}d_{33} \geq 0$.
A (1,4)	Det A (1,4) = $d_{11}d_{44} \geq 0$.
A (1,5)	Det A (1,5) = $d_{11}d_{55} \geq 0$.
A (2,3)	Det A (2,3) = $d_{22}d_{33} \geq 0$.
A (2,4)	Det A (2,4) = $d_{22}d_{44} \geq 0$.
A (2,5)	Det A (2,5) = $d_{22}d_{55} \geq 0$.
A (3,4)	Det A (3,4) = $d_{33}d_{44} \geq 0$.
A (3,5)	Det A (3,5) = $d_{33}d_{55} \geq 0$.
A (4,5)	Det A (4,5) = $d_{44}d_{55} \geq 0$.

By definition of completion, determinants of 2×2 submatrices are obtained. i.e., $\det A (1,3) = d_{11}d_{33} - x_{13}x_{31} \geq 0$, since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero then; $\det A (1,3) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 2: Determinants of 3×3 sub-matrices

Principal submatrix	Principal minor
A (1,2,3)	Det A (1,2,3) = $d_{11}d_{22}d_{33} \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{11}d_{22}d_{44} \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{11}d_{22}d_{55} \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{33}d_{44}d_{55} \geq 0$.

Determinants of 3×3 submatrices are then obtained. i.e., $\det A(1,2,3) = d_{11}(d_{22}d_{33} - x_{23}x_{32}) - x_{12}(x_{21}d_{33} - x_{23}x_{31}) + x_{13}(x_{21}x_{32} - d_{22}x_{31}) \geq 0$, since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero then; $\det A(1,2,3) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 3: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A(1,2,3,4)	$A(1,2,3,4) = d_{11}d_{22}d_{33}d_{44} \geq 0$.
A(1,2,3,5)	$A(1,2,3,5) = d_{11}d_{22}d_{33}d_{55} \geq 0$.
A(1,2,4,5)	$A(1,2,4,5) = d_{11}d_{22}d_{44}d_{55} \geq 0$.
A(1,3,4,5)	$A(1,3,4,5) = d_{11}d_{33}d_{44}d_{55} \geq 0$.
A(2,3,4,5)	$A(2,3,4,5) = d_{22}d_{33}d_{44}d_{55} \geq 0$.


$\det(A) = d_{11}d_{22}d_{33}d_{44}d_{55} \geq 0$. Therefore $\det(A) = 1 > 0$.

Determinants of 4×4 submatrices are obtained. i.e., $\det A(1,2,3,4) = \det(1,2,3,4) = d_{11}[d_{22}(d_{33}d_{44} - x_{34}x_{43}) - x_{23}(d_{44}x_{32} - x_{34}x_{42}) + x_{24}(x_{32}x_{43} - d_{33}x_{42})]$

$$\begin{aligned}
 & - x_{12}[x_{21}(d_{33}d_{44} - x_{34}x_{43}) - x_{23}(d_{44}x_{31} - x_{34}x_{41}) + x_{24}(x_{31}x_{43} - d_{33}x_{41})] \\
 & + x_{13}[x_{21}(x_{32}d_{44} - x_{34}x_{42}) - d_{22}(d_{44}x_{31} - x_{34}x_{41}) + x_{24}(x_{31}x_{42} - x_{32}x_{41})] \\
 & - x_{14}[x_{21}(x_{32}x_{43} - d_{33}x_{42}) - d_{22}(x_{43}x_{31} - d_{33}x_{41}) + x_{23}(x_{31}x_{42} - x_{32}x_{41})] \geq 0.
 \end{aligned}$$

since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero then; $\det A(1,2,3,4) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 4×4 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

DiJABB_121114 D of order 5 and 1 arc: Consider the diJABB_121114 $D = \{(1,1), (2,2), (2,5), (3,3), (4,4), (5,5)\}$ with 5 vertices and one arc given by:



The Partial matrix that specifies the diJABB_121114 D is

$$A = \begin{pmatrix} d_{11} & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & d_{22} & x_{23} & x_{24} & a_{25} \\ x_{31} & x_{32} & d_{33} & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & d_{44} & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix}$$

Figure 2 Digraph D of order 5 and 1 arc

By definition of partial Wss P_0 - matrix $d_1 \geq 0, d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0$.

For $d_{ii} = 1, a_{ij} = 1$, then the partial matrix becomes $A = \begin{pmatrix} 1 & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & 1 & x_{23} & x_{24} & 1 \\ x_{31} & x_{32} & 1 & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & 1 & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix}$. Next, we

compute the principal minors.

$\det(1,2) = d_{11}d_{22} - x_{12}x_{21}$. Setting the unspecified entry to zero, and $d_{ii} = 1, a_{ij} = 1$, i.e., $\det(1,2) = 1 - 0 = 1 > 0$. $\det(2,5) = d_{22}d_{55} - a_{25}x_{52}$, i.e., $a_{25}x_{52} = 1 \cdot 0 = 0$. Similarly,

$$\text{Det (1,3)} = d_{11} d_{33} - x_{13} x_{31}, \text{Det (1,4)} = d_{11} d_{44} - x_{14} x_{41}$$

$$\text{Det (1,5)} = d_{11} d_{55} - x_{15} x_{51}, \text{Det (2,3)} = d_{22} d_{33} - x_{23} x_{32}, \text{Det (2,4)} = d_{22} d_{44} - x_{24} x_{42}$$

$$\text{Det (2,5)} = d_{22} d_{55} - x_{25} x_{52}, \text{Det (3,4)} = d_{33} d_{44} - x_{34} x_{43}, \text{Det (3,5)} = d_{33} d_{55} - x_{35} x_{53}$$

$$\text{Det (4,5)} = d_{44} d_{55} - x_{45} x_{54}.$$

TABLE 4: Determinants of 2×2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	Det A (1,2) = $d_{11}d_{22} \geq 0$.
A (1,3)	Det A (1,3) = $d_{11}d_{33} \geq 0$.
A (1,4)	Det A (1,4) = $d_{11}d_{44} \geq 0$.
A (1,5)	Det A (1,5) = $d_{11}d_{55} \geq 0$.
A (2,3)	Det A (2,3) = $d_{22}d_{33} \geq 0$.
A (2,4)	Det A (2,4) = $d_{22}d_{44} \geq 0$.
A (2,5)	Det A (2,5) = $d_{22}d_{55} \geq 0$.
A (3,4)	Det A (3,4) = $d_{33}d_{44} \geq 0$.
A (3,5)	Det A (3,5) = $d_{33}d_{55} \geq 0$.
A (4,5)	Det A (4,5) = $d_{44}d_{55} \geq 0$.

By definition of completion, determinants of 2×2 submatrices are obtained. i.e. $\det (2,5) = d_{22} d_{55} - x_{25} x_{52} \geq 0$, since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,3) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 5: Determinants of 3×3 sub-matrices

Principal submatrix	Principal minor
A (1,2,3)	Det A (1,2,3) = $d_{11}d_{22}d_{33} \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{11}d_{22}d_{44} \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{11}d_{22}d_{55} \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{33}d_{44}d_{55} \geq 0$.

Determinants of 3×3 submatrices are obtained. i.e., $\det A (1,2,3) = d_{11} (d_{22}d_{33} - x_{23} x_{32}) - x_{12} (x_{21} d_{33} - x_{23} x_{31}) + x_{13} (x_{21} x_{32} - d_{22}x_{31}) \geq 0$, since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2,3) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 6: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A (1,2,3,4)	A (1,2,3,4) = $d_{11}d_{22}d_{33}d_{44} \geq 0$.

A (1,2,3,5)	A (1,2,3,5) = d ₁₁ d ₂₂ d ₃₃ d ₅₅ ≥ 0.
A (1,2,4,5)	A (1,2,4,5) = d ₁₁ d ₂₂ d ₄₄ d ₅₅ ≥ 0.
A (1,3,4,5)	A (1,3,4,5) = d ₁₁ d ₃₃ d ₄₄ d ₅₅ ≥ 0.
A (2,3,4,5)	A (2,3,4,5) = d ₂₂ d ₃₃ d ₄₄ d ₅₅ ≥ 0.

Det (A) = d₁₁d₂₂d₃₃d₄₄d₅₅ ≥ 0. Therefore det (A) = 1 > 0.

Determinants of 4 × 4 submatrices are obtained. i.e., det A (1,2,3,4) = Det (1,2,3,4) = d₁₁[d₂₂ (d₃₃ d₄₄ - x₃₄ x₄₃) - x₂₃ (d₄₄ x₃₂ - x₃₄ x₄₂) + x₂₄ (x₃₂ x₄₃ - d₃₃ x₄₂)]


$$- x_{12} [x_{21} (d_{33} d_{44} - x_{34} x_{43}) - x_{23} (d_{44} x_{31} - x_{34} x_{41}) + x_{24} (x_{31} x_{43} - d_{33} x_{41})]$$

$$+ x_{13} [x_{21} (x_{32} d_{44} - x_{34} x_{42}) - d_{22} (d_{44} x_{31} - x_{34} x_{41}) + x_{24} (x_{31} x_{42} - x_{32} x_{41})]$$

$$- x_{14} [x_{21} (x_{32} x_{43} - d_{33} x_{42}) - d_{22} (x_{43} x_{31} - d_{33} x_{41}) + x_{23} (x_{31} x_{42} - x_{32} x_{41})] \geq 0.$$

since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and a_{ij}=1 then; det A (1,2,3,4) = 1 - 0 + 0 - 0 = 1 > 0. Therefore, all the determinants of 4 × 4 submatrices are non-negative then the partial matrix can be completed into Wss P_o-matrix. Hence it has zero completion into a Wss P_o-matrix.

DiJABB_121114 D of order 5 and 2 arcs: consider a diJABB_121114 D = {(1,1), (2,2), (3,3), (3,4), (4,3), (4,4), (5,5)} with 5 vertices and 2 arcs given by:



The Partial matrix that specifies the diJABB_121114 D is

$$A = \begin{pmatrix} d_{11} & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & d_{22} & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & d_{33} & a_{34} & x_{35} \\ x_{41} & x_{42} & a_{43} & d_{44} & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix}.$$

Figure 3. Digraph D of order 5 and 2 arcs

By definition of partial Wss P_o- matrix d₁ ≥ 0, d₂ ≥ 0, d₃ ≥ 0, d₄ ≥ 0, d₅ ≥ 0.

For d_{ii} = 1, a_{ij} = 1, then the partial matrix becomes A = $\begin{pmatrix} 1 & x_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & 1 & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & 1 & 1 & x_{35} \\ x_{41} & x_{42} & 1 & 1 & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix}$. Next, we

compute the principal minors.

Det (1,2) = d₁₁d₂₂ - x₁₂x₂₁, Setting the unspecified entry to zero, i.e., x₁₂x₂₁ = 0, Det (3,4) = d₃₃ d₄₄ - a₃₄ a₄₃ = 1-1 ≥ 0. Similarly;

$$\text{Det (1,3)} = d_{11} d_{33} - x_{13} x_{31}, \text{Det (1,4)} = d_{11} d_{44} - x_{14} x_{41}$$

$$\text{Det (1,5)} = d_{11} d_{55} - x_{15} x_{51}, \text{Det (2,3)} = d_{22} d_{33} - x_{23} x_{32},$$

$$\text{Det (2,4)} = d_{22} d_{44} - x_{24} x_{42}, \text{Det (2,5)} = d_{22} d_{55} - x_{25} x_{52},$$

$$\text{Det (3,5)} = d_{33} d_{55} - x_{35} x_{53}, \text{Det (4,5)} = d_{44} d_{55} - x_{45} x_{54}$$

TABLE 7: Determinants of 2×2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	Det A (1,2) = $d_{11}d_{22} \geq 0$.
A (1,3)	Det A (1,3) = $d_{11}d_{33} \geq 0$.
A (1,4)	Det A (1,4) = $d_{11}d_{44} \geq 0$.
A (1,5)	Det A (1,5) = $d_{11}d_{55} \geq 0$.
A (2,3)	Det A (2,3) = $d_{22}d_{33} \geq 0$.
A (2,4)	Det A (2,4) = $d_{22}d_{44} \geq 0$.
A (2,5)	Det A (2,5) = $d_{22}d_{55} \geq 0$.
A (3,4)	Det A (3,4) = $(d_{33}d_{44}) - (a_{34}a_{43}) \geq 0$
A (3,5)	Det A (3,5) = $d_{33}d_{55} \geq 0$.
A (4,5)	Det A (4,5) = $d_{44}d_{55} \geq 0$.

By definition of completion, determinants of 2×2 submatrices are obtained. i.e. $\text{Det} (1,2) = d_{11}d_{22} - x_{12}x_{21} \geq 0$, $\text{Det} (3,4) = d_{33}d_{44} - a_{34}a_{43} \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2) = 1 - 0 = 1 > 0$, $\text{Det} (3,4) = 1 - 1 \geq 0$. Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 8: Determinants of 3×3 sub-matrices

Principal submatrix	Principal minor
A (1,2,3)	Det A (1,2,3) = $d_{11}d_{22}d_{33} \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{11}d_{22}d_{44} \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{11}d_{22}d_{55} \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{55}(d_{33}d_{44}) - (a_{34}a_{43}) \geq 0$

Determinants of 3×3 submatrices are obtained. i.e., $\text{Det} (3,4,5) = d_{33}(d_{44}d_{55} - x_{45}x_{54}) - a_{34}(a_{43}d_{55} - x_{45}x_{53}) + x_{35}(a_{43}x_{54} - d_{44}x_{53}) \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (3,4,5) = 1 - 1 = 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 9: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A (1,2,3,4)	Det (1,2,3,4) = $d_{11}d_{22}d_{33}d_{44} \geq 0$.
A (1,2,3,5)	Det (1,2,3,5) = $d_{11}d_{22}d_{33}d_{55} \geq 0$.
A (1,2,4,5)	Det (1,2,4,5) = $d_{11}d_{22}d_{44}d_{55} \geq 0$.
A (1,3,4,5)	Det (1,3,4,5) = $d_{11}d_{55}(d_{33}d_{44}) - (a_{34}a_{43}) \geq 0$.
A (2,3,4,5)	Det (2,3,4,5) = $d_{22}d_{33}d_{44}d_{55} \geq 0$.

$\text{Det} (A) = d_{11}d_{22}d_{33}d_{44}d_{55} - (a_{34}a_{43}) \geq 0$. therefore $\text{Det} (A) = 1 - 1 = 0$.

Determinants of 4×4 submatrices are obtained. i.e., $\text{Det}(1,3,4,5) = d_{11} [d_{33} (d_{44} d_{55} - x_{45} x_{54}) - a_{34} (d_5 a_{43} - x_{45} x_{53}) + x_{35} (a_{43} x_{54} - d_{44} x_{53})]$


$$- x_{13} [x_{31} (d_{44} d_{55} - x_{45} x_{54}) - a_{34} (d_5 x_{41} - x_{45} x_{51}) + x_{35} (x_{41} x_{54} - d_{44} x_{51})]$$

$$+ x_{14} [x_{31} (d_{55} a_{43} - x_{45} x_{53}) - d_{33} (d_5 x_{41} - x_{45} x_{51}) + x_{35} (x_{41} x_{53} - a_{43} x_{51})]$$

$$- x_{15} [x_{31} (a_{43} x_{54} - d_{44} x_{53}) - d_{33} (x_{41} x_{54} - d_{44} x_{51}) + a_{34} (x_{41} x_{53} - x_{51} a_{43})] \geq 0.$$

Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij} = 1$ then; $\det A(1,3,4,5) = 0 - 0 + 0 - 0 = 0$. Therefore, all the determinants of 4×4 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

DiJABB_121114 D of order 5 and 3 arcs; consider a diJABB_121114 $D = \{(1,1), (1,2), (1,3), (2,1), (2,2), (3,3), (4,4), (5,5)\}$ with 5 vertices and 3 arcs given by:



The partial matrix that specifies the diJABB_121114 D is $A =$

$$\begin{pmatrix} d_{11} & a_{12} & a_{13} & x_{14} & x_{15} \\ a_{21} & d_{22} & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & d_{33} & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & d_{44} & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix}$$

By definition of partial Wss P_0 - matrix $d_1 \geq 0, d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0$.

For $d_{ii} = 1, a_{ij} = 1$, then the partial matrix becomes $A = \begin{pmatrix} 1 & 1 & 1 & x_{14} & x_{15} \\ 1 & 1 & x_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & 1 & x_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & 1 & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix}$. Next, we

compute the principal minors.

$\text{Det}(1,2) = d_{11}d_{22} - a_{12}a_{21}$. Setting unspecified entries to zero, $d_{ii} = 1, a_{ij} = 1$, then we have; $1 - 1 \geq 0$.

$\text{Det}(1,3) = d_{11}d_{33} - a_{13}x_{31} = 1 - x_{31} = 1 - 0 = 1 \geq 0$. Similarly

$\text{Det}(1,2) = d_{11}d_{22} - a_{12}a_{21}$, $\text{Det}(1,4) = d_{11}d_{44} - x_{14}x_{41}$,

$\text{Det}(1,5) = d_{11}d_{55} - x_{15}x_{51}$, $\text{Det}(2,3) = d_{22}d_{33} - x_{23}x_{32}$, $\text{Det}(2,4) = d_{22}d_{44} - x_{24}x_{42}$

$\text{Det}(2,5) = d_{22}d_{55} - x_{25}x_{52}$, $\text{Det}(3,4) = d_{33}d_{44} - x_{34}x_{43}$, $\text{Det}(3,5) = d_{33}d_{55} - x_{35}x_{53}$

$\text{Det}(4,5) = d_{44}d_{55} - x_{45}x_{54}$

TABLE 10: Determinants of 2×2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	$\text{Det} A(1,2) = (d_{11}d_{22}) - (a_{12}a_{21}) \geq 0$
A (1,3)	$\text{Det} A(1,3) = d_{11}d_{33} \geq 0$.
A (1,4)	$\text{Det} A(1,4) = d_{11}d_{44} \geq 0$.
A (1,5)	$\text{Det} A(1,5) = d_{11}d_{55} \geq 0$.
A (2,3)	$\text{Det} A(2,3) = d_{22}d_{33} \geq 0$.
A (2,4)	$\text{Det} A(2,4) = d_{22}d_{44} \geq 0$.
A (2,5)	$\text{Det} A(2,5) = d_{22}d_{55} \geq 0$.

A (3,4)	Det A (3,4) = $d_{33}d_{44} \geq 0$.
A (3,5)	Det A (3,5) = $d_{33}d_{55} \geq 0$.
A (4,5)	Det A (4,5) = $d_{44}d_{55} \geq 0$.

By definition of completion, determinants of 2×2 submatrices are obtained. i.e. Det (1,2) = $d_{11}d_{22} - a_{12}a_{21} \geq 0$. Det (1,3) = $d_{11}d_{33} - a_{13}x_{31} \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; Det (1,2) = $1-1 \geq 0$, det A (1,3) = $1-0 = 1 > 0$. Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 11: Determinants of 3×3 sub-matrices

Principal submatrix	Principal minor
A (1,2,3)	Det A (1,2,3) = $d_{33} (d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{44}(d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{55}(d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{33}d_{44}d_{55} \geq 0$.

Determinants of 3×3 submatrices are obtained. i.e., Det (3,4,5) = $d_{11} (d_{22}d_{33} - x_{23} x_{32}) - a_{12} (a_{21} d_{33} - x_{23} x_{31}) + a_{13} (a_{21} x_{32} - d_{22} x_{31}) \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; det A (3,4,5) = $1-1+0 = 1 > 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 12: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A (1,2,3,4)	Det A (1,2,3,4) = $d_{33}d_{44} (d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,2,3,5)	Det A (1,2,3,5) = $d_{33}d_{55}(d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,2,4,5)	Det A (1,2,4,5) = $d_{33}d_{55}(d_{11}d_{22} - a_{12}a_{21}) \geq 0$.
A (1,3,4,5)	Det A (1,3,4,5) = $d_{11}d_{33}d_{44}d_{55} \geq 0$.
A (2,3,4,5)	Det A (2,3,4,5) = $d_{22}d_{33}d_{44}d_{55} \geq 0$.

Det (A) = $d_{11}d_{22}d_{33}d_{44}d_{55} - (a_{12}a_{21}d_{33}d_{44}d_{55}) \geq 0$. Therefore, Det (A) = $1 - 1 = 0$.

Determinants of 4×4 submatrices are obtained. i.e., Det (1,2,3,4) = $d_{11}[d_{22} (d_{33} d_{44} - x_{34} x_{43}) - x_{23} (d_{44} x_{32} - x_{34} x_{42}) + x_{24} (x_{32} x_{43} - d_{33} x_{42})]$


$$\begin{aligned}
& - a_{12} [a_{21} (d_{33} d_{44} - x_{34} x_{43}) - x_{23} (d_{44} x_{31} - x_{34} x_{41}) + x_{24} (x_{31} x_{43} - d_{33} x_{41})] \\
& + a_{13} [a_{21} (x_{32} d_{44} - x_{34} x_{42}) - d_{22} (d_{44} x_{31} - x_{34} x_{41}) + x_{24} (x_{31} x_{42} - x_{32} x_{41})] \\
& - x_{14} [a_{21} (x_{32} x_{43} - d_{33} x_{42}) - d_{22}(x_{43} x_{31} - d_{33} x_{41}) + x_{23} (x_{31} x_{42} - x_{32} x_{41})] \geq 0.
\end{aligned}$$

Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; det A (1,2,3,4) = $1 - 1 + 0 - 0 = 0$. Therefore, all the determinants of 4×4

submatrices are non-negative then the partial matrix can be completed into Wss P_o-matrix. Hence it has zero completion into a Wss P_o-matrix.

DiJABB_121114 D of order 5 and 4 arcs; consider a diJABB_121114 D = {(1,1), (1,2), (2,2), (2,3), (3,3), (3,4), (3,5), (4,4), (5,5)} with 5 vertices and 4 arcs given by:

Figure 5
Digraph
D of order
5 and 4
arcs



The Partial matrix that specifies the diJABB_121114 D is $A =$

$$\begin{pmatrix} d_{11} & a_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & d_{22} & a_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & d_{33} & a_{34} & a_{35} \\ x_{41} & x_{42} & x_{43} & d_{44} & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix} \text{By definition of partial Wss P}_o\text{- matrix } d_i \geq 0, \\ d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0.$$

For $d_{ii} = 1, a_{ij} = 1,$ then the partial matrix becomes $A =$

$$\begin{pmatrix} 1 & 1 & x_{13} & x_{14} & x_{15} \\ x_{21} & 1 & 1 & x_{24} & x_{25} \\ x_{31} & x_{32} & 1 & 1 & 1 \\ x_{41} & x_{42} & x_{43} & 1 & x_{45} \\ x_{51} & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix} \text{Next, we compute the principal minors.}$$

Det (1,2) = $d_{11} d_{22} - a_{12} x_{21}$ Setting unspecified entries to zero, $d_{ii} = 1, a_{ij} = 1,$ then we have; $1 - 0 = 0$. Similarly; Det (1,3) = $d_{11} d_{33} - x_{13} x_{31},$ Det (1,4) = $d_{11} d_{44} - x_{14} x_{41},$ Det (1,5) = $d_{11} d_{55} - x_{15} x_{51},$ Det (2,3) = $d_{22} d_{33} - a_{23} x_{32},$ Det (2,4) = $d_{22} d_{44} - x_{24} x_{42},$ Det (2,5) = $d_{22} d_{55} - x_{25} x_{52},$ Det (3,4) = $d_{33} d_{44} - a_{34} x_{43},$ Det (3,5) = $d_{33} d_{55} - a_{35} x_{53}$

Det (4 5) = $d_{44} d_{55} - x_{45} x_{54}.$

TABLE 13: Determinants of 2x2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	Det A (1,2) = $d_{11}d_{22} \geq 0.$
A (1,3)	Det A (1,3) = $d_{11}d_{33} \geq 0.$
A (1,4)	Det A (1,4) = $d_{11}d_{44} \geq 0.$
A (1,5)	Det A (1,5) = $d_{11}d_{55} \geq 0.$
A (2,3)	Det A (2,3) = $d_{22}d_{33} \geq 0.$
A (2,4)	Det A (2,4) = $d_{22}d_{44} \geq 0.$
A (2,5)	Det A (2,5) = $d_{22}d_{55} \geq 0.$
A (3,4)	Det A (3,4) = $d_{33}d_{44} \geq 0.$
A (3,5)	Det A (3,5) = $d_{33}d_{55} \geq 0.$
A (4,5)	Det A (4,5) = $d_{44}d_{55} \geq 0.$

By definition of completion, determinants of 2×2 submatrices are obtained. i.e. Det (1,2) = $d_{11}d_{22} - a_{12}x_{21} \geq 0.$ Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij} = 1$ then; $\det A (1,2) = 1 - 0 = 1 > 0.$ Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_o-matrix. Hence it has zero completion into a Wss P_o-matrix.

TABLE 14: Determinants of 3x3 sub-matrices

Principal submatrix	Principal minor
---------------------	-----------------

A (1,2,3)	Det A (1,2,3) = $d_{11}d_{22}d_{33} \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{11}d_{22}d_{44} \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{11}d_{22}d_{55} \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{33}d_{44}d_{55} \geq 0$.

Determinants of 3×3 submatrices are obtained. i.e., $\text{Det} (1,2,3) = d_{11} (d_{22}d_{33} - a_{23} x_{32}) - a_{12} (a_{21} d_{33} - a_{23} x_{31}) + x_{13} (x_{21} x_{32} - d_{22} x_{31}) \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2,3) = 1 - 1 + 0 = 1 > 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_o -matrix. Hence it has zero completion into a Wss P_o -matrix.

TABLE 15: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A (1,2,3,4)	Det A (1,2,3,4) = $d_{11}d_{22}d_{33}d_{44} \geq 0$.
A (1,2,3,5)	Det A (1,2,3,5) = $d_{11}d_{22}d_{33}d_{55} \geq 0$.
A (1,2,4,5)	Det A (1,2,4,5) = $d_{11}d_{22}d_{44}d_{55} \geq 0$.
A (1,3,4,5)	Det A (1,3,4,5) = $d_{11}d_{33}d_{44}d_{55} \geq 0$.
A (2,3,4,5)	Det A (2,3,4,5) = $d_{22}d_{33}d_{44}d_{55} \geq 0$.

$\text{Det} (A) = d_{11}d_{22}d_{33}d_{44}d_{55} \geq 0$. Therefore $\text{Det} (A) = 1 > 0$.

Determinants of 4×4 submatrices are obtained. i.e., $\text{Det} (1,2,3,4) = d_{11}[d_{22} (d_{33} d_{44} - a_{34} x_{43}) - a_{23} (d_{44} x_{32} - a_{34} x_{42}) + x_{24} (x_{32} x_{43} - d_{33} x_{42})]$


$$- a_{12} [x_{21} (d_{33} d_{44} - a_{34} x_{43}) - a_{23} (d_{44} x_{31} - a_{34} x_{41}) + x_{24} (x_{31} x_{43} - d_{33} x_{41})]$$

$$+ x_{13} [x_{21} (x_{32} d_{44} - a_{34} x_{42}) - d_{22} (d_{44} x_{31} - a_{34} x_{41}) + x_{24} (x_{31} x_{42} - x_{32} x_{41})]$$

$$- x_{14} [x_{21} (x_{32} x_{43} - d_{33} x_{42}) - d_{22}(x_{43} x_{31} - d_{33} x_{41}) + a_{23} (x_{31} x_{42} - x_{32} x_{41})] \geq 0$$

Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2,3,4) = 1 - 0 + 0 - 0 = 1 > 0$. Therefore, all the determinants of 4×4 submatrices are non-negative then the partial matrix can be completed into Wss P_o -matrix. Hence it has zero completion into a Wss P_o -matrix.

DiJABB_121114 D of order 5 and 5 arcs: consider a diJABB_121114 $D = \{(1,1), (1,2), (2,2), (2,3), (3,3), (3,4), (4,4), (4,5), (5,1), (5,5)\}$ with 5 vertices and 5 arcs given by:



The partial matrix that specifies the diJABB_121114 D is

$$A = \begin{pmatrix} d_{11} & a_{12} & x_{13} & x_{14} & x_{15} \\ x_{21} & d_{22} & a_{23} & x_{24} & x_{25} \\ x_{31} & x_{32} & d_{33} & a_{34} & x_{35} \\ x_{41} & x_{42} & x_{43} & d_{44} & a_{45} \\ a_{51} & x_{52} & x_{53} & x_{54} & d_{55} \end{pmatrix}.$$

Figure 6
Digraph
D of
order 5
and 5
arcs

By definition of partial Wss P_o - matrix $d_1 \geq 0, d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0$.
By definition of partial Wss P_o - matrix $d_1 \geq 0, d_2 \geq 0, d_3 \geq 0, d_4 \geq 0, d_5 \geq 0$.

For $d_{ii} = 1, a_{ij} = 1$, then the partial matrix becomes $A = \begin{pmatrix} 1 & 1 & x_{13} & x_{14} & x_{15} \\ x_{21} & 1 & 1 & x_{24} & x_{25} \\ x_{31} & x_{32} & 1 & 1 & x_{35} \\ x_{41} & x_{42} & x_{43} & 1 & 1 \\ 1 & x_{52} & x_{53} & x_{54} & 1 \end{pmatrix}$. Next, we

compute the principal minors.

$\text{Det}(1,2) = d_{11}d_{22} - a_{12}x_{21}$, $\text{Det}(1,2) = d_{11}d_{22} - a_{12}x_{21}$ Setting unspecified entries to zero, $d_{ii} = 1, a_{ij} = 1$, then we have; $1 - 0 = 1 > 0$. similarly; $\text{Det}(1,3) = d_{11}d_{33} - x_{13}x_{31}$, $\text{Det}(1,4) = d_{11}d_{44} - x_{14}x_{41}$

$\text{Det}(1,5) = d_{11}d_{55} - x_{15}a_{51}$, $\text{Det}(2,3) = d_{22}d_{33} - a_{23}x_{32}$, $\text{Det}(2,4) = d_{22}d_{44} - x_{24}x_{42}$

$\text{Det}(2,5) = d_{22}d_{55} - x_{25}x_{52}$, $\text{Det}(3,4) = d_{33}d_{44} - a_{34}x_{43}$, $\text{Det}(3,5) = d_{33}d_{55} - x_{35}x_{53}$

$\text{Det}(4,5) = d_{44}d_{55} - a_{45}x_{54}$.

TABLE 16: Determinants of 2x2 sub-matrices

Principal submatrix	Principal minor
A (1,2)	$\text{Det} A(1,2) = d_{11}d_{22} \geq 0$.
A (1,3)	$\text{Det} A(1,3) = d_{11}d_{33} \geq 0$.
A (1,4)	$\text{Det} A(1,4) = d_{11}d_{44} \geq 0$.
A (1,5)	$\text{Det} A(1,5) = d_{11}d_{55} \geq 0$.
A (2,3)	$\text{Det} A(2,3) = d_{22}d_{33} \geq 0$.
A (2,4)	$\text{Det} A(2,4) = d_{22}d_{44} \geq 0$.
A (2,5)	$\text{Det} A(2,5) = d_{22}d_{55} \geq 0$.
A (3,4)	$\text{Det} A(3,4) = d_{33}d_{44} \geq 0$.
A (3,5)	$\text{Det} A(3,5) = d_{33}d_{55} \geq 0$.
A (4,5)	$\text{Det} A(4,5) = d_{44}d_{55} \geq 0$.

Determinants of 2×2 submatrices are obtained. i.e. $\text{Det}(1,2) = d_{11}d_{22} - a_{12}x_{21} \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij} = 1$ then; $\text{det} A(1,2) = 1 - 0 = 1 > 0$. Therefore, all the determinants of 2×2 submatrices are non-negative then the partial matrix can be completed into Wss P_o -matrix. Hence it has zero completion into a Wss P_o -matrix.

TABLE 17: Determinants of 3x3 sub-matrices

Principal submatrix	Principal minor
A (1,2,3)	Det A (1,2,3) = $d_{11}d_{22}d_{33} \geq 0$.
A (1,2,4)	Det A (1,2,4) = $d_{11}d_{22}d_{44} \geq 0$.
A (1,2,5)	Det A (1,2,5) = $d_{11}d_{22}d_{55} \geq 0$.
A (1,3,4)	Det A (1,3,4) = $d_{11}d_{33}d_{44} \geq 0$.
A (1,3,5)	Det A (1,3,5) = $d_{11}d_{33}d_{55} \geq 0$.
A (1,4,5)	Det A (1,4,5) = $d_{11}d_{44}d_{55} \geq 0$.
A (2,3,4)	Det A (2,3,4) = $d_{22}d_{33}d_{44} \geq 0$.
A (2,3,5)	Det A (2,3,5) = $d_{22}d_{33}d_{55} \geq 0$.
A (2,4,5)	Det A (2,4,5) = $d_{22}d_{44}d_{55} \geq 0$.
A (3,4,5)	Det A (3,4,5) = $d_{33}d_{44}d_{55} \geq 0$.

Determinants of 3×3 submatrices are obtained. i.e., $\text{Det} (1,2,3) = d_{11} (d_{22}d_{33} - a_{23} x_{32}) - a_{12} (x_{21} d_{33} - a_{23} x_{31}) + x_{13} (x_{21} x_{32} - d_{22} x_{31}) \geq 0$. Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2,3) = 1 - 0 + 0 = 1 > 0$. Therefore, all the determinants of 3×3 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

TABLE 18: Determinants of 4×4 sub-matrices

Principal sub-matrix	Principal minor
A (1,2,3,4)	Det A (1,2,3,4) = $d_{11}d_{22}d_{33}d_{44} \geq 0$.
A (1,2,3,5)	Det A (1,2,3,5) = $d_{11}d_{22}d_{33}d_{55} \geq 0$.
A (1,2,4,5)	Det A (1,2,4,5) = $d_{11}d_{22}d_{44}d_{55} \geq 0$.
A (1,3,4,5)	Det A (1,3,4,5) = $d_{11}d_{33}d_{44}d_{55} \geq 0$.
A (2,3,4,5)	Det A (2,3,4,5) = $d_{22}d_{33}d_{44}d_{55} \geq 0$.

$\text{Det} (A) = (d_{11}d_{22}d_{33}d_{44}d_{55}) - (a_{12}a_{23}a_{34}a_{45}a_{51}) \geq 0$. Therefore $\text{Det} (A) = 1 - 1 = 0$. By definition of completion, determinants of 4×4 submatrices are obtained. i.e., $\text{Det} (1,2,3,4) = d_{11}[d_{22} (d_{33} d_{44} - a_{34} x_{43}) - a_{23} (d_{44} x_{32} - a_{34} x_{42}) + x_{24} (x_{32} x_{43} - d_{33} x_{42})]$

$$\begin{aligned}
& - a_{12} [x_{21} (d_{33} d_{44} - a_{34} x_{43}) - a_{23} (d_{44} x_{31} - a_{34} x_{41}) + x_{24} (x_{31} x_{43} - d_{33} x_{41})] \\
& + x_{13} [x_{21} (x_{32} d_{44} - a_{34} x_{42}) - d_{22} (d_{44} x_{31} - a_{34} x_{41}) + x_{24} (x_{31} x_{42} - x_{32} x_{41})] \\
& - x_{14} [x_{21} (x_{32} x_{43} - d_{33} x_{42}) - d_{22}(x_{43} x_{31} - d_{33} x_{41}) + a_{23} (x_{31} x_{42} - x_{32} x_{41})] \geq 0.
\end{aligned}$$

Since diagonal entries is given to be 1 or larger than other entries while setting unspecified entries to zero and $a_{ij}=1$ then; $\det A (1,2,3,4) = 1 - 0 + 0 - 0 = 1 > 0$. Therefore, all the determinants of 4×4 submatrices are non-negative then the partial matrix can be completed into Wss P_0 -matrix. Hence it has zero completion into a Wss P_0 -matrix.

All the diJABB_121114s mentioned above are not isomorphic to each other.

4.0 CONCLUSION

Upon comprehensive examination and analysis of various diJABB_121114 patterns, the research arrived at significant conclusions. We used the zero-completion method where unspecified entries in partial Wss P_0 - matrices are set to zero to evaluate principal minors. We concluded that a null JABB_121114 of order 5 has Wss P_0 - completion. Additionally, we demonstrated that diJABB_121114s of order 5 with one to five arcs whether cyclic, acyclic, or with cliques, can be completed into a Wss P_0 - matrix. Therefore, we found that all partial Wss P_0 - matrices have completion into a Wss P_0 - matrix. Conversely, no partial Wss P_0 - matrix formed by

diJABB_121114s of order 5 with up to 5 arcs lacks the ability to be completed into a Wss P₀-matrix. Insights gained from these studies could be applied to fill gaps in data from retail surveys, aiding in the prediction of market trends.

REFERENCES

- [1] Tomno, V. (2018). On Completion Problems for Various Subclasses of. *Annals of Pure and Applied Mathematics*, 18(2), 207-212.
- [2] Tomno, V., Nyamwala, F., & Kamaku, W. (2018). The Wss Po Matrix Completion Problem for Symmetric Patterns of Acyclic DiJABB_121114s of Order Four. *International Journal of Sciences: Basic and Applied Research (IJSBAR)*, 37(1), pp. 112-121.
- [3] Bowers, J., Evers, J., Hogben, L., Shaner, S., Snider, K., & Wangsness, A. (2006). On completion problems for various classes of P-matrices. *Linear algebra and its applications*, 413(2-3), 342-354.
- [4] Choi, J. Y., DeAlba, L., Hogben, L., Kivunge, B., Nordstrom, S., & Shedenhelm, M. (2003). The nonnegative P₀-matrix completion problem.
- [5] Choi, J. Y., DeAlba, L., Hogben, L., Maxwell, M., & Wangsness, A. (2002). The P₀-matrix completion problem. *Electronic Journal of linear Algebra*, 9.
- [6] DeAlba, L., Hardy, T., Hogben, L., & Wangsness, A. (2003). The (weakly) sign symmetric P-matrix completion problems. *The Electronic Journal of Linear Algebra*, 10, 257-271.
- [7] Harary, F. (2018). *JABB_121114 theory (on Demand Printing of 02787)*. CRC Press.
- [8] Paula, M., Kamakub, W., & Nyaga, L. (2021). The non-negative P₀-matrix completion problem for 5× 5 matrices specifying diJABB_121114s with 5 vertices and 4 arcs for acyclic diJABB_121114s.
- [9] Mutembei, J., Kamaku, W., & Kivunge, B. (2015). Positive **P^{0,1}** Matrix Completion Problem for DiJABB_121114s of Order Three with Zero, One, Two and Three Arcs.
- [10] Entner, H. (2020). Matrix Completion Problems. University of Innsbruck. https://www.uibk.ac.at/mathematik/algebra/media/teaching/matrix-completion-problems_entner.pdf.
- [11] Fallat, S. M., Johnson, C. R., Torregrosa, J. R., & Urbano, A. M. (2000). P-matrix completions under weak symmetry assumptions. *Linear Algebra and Its Applications*, 312(1-3), 73-91.
- [12] <https://www.academia.edu/98565036/on> completion problems for various subclasses.
- [13] Munyiri J. et al, IJSBAR (2014) Vol 15, No. 1, pp 379-385]