

Original Research Article

Efficacy of Various Ready-mix Insecticides against *Spodopteralitura* (Fab.) in Groundnut

ABSTRACT

Field experiment was carried out to study the efficacy of various ready-mix insecticides against *Spodopteralitura*(Fab.) in groundnut at Main Oilseeds Research Station, JAU, Junagadh during *kharif*, 2023. Out of the seven ready-mix insecticidal treatments evaluated, chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% found most effective against *S. litura* after two sprays. It recorded the lowest larval population (0.60 larvae/plant), highest pod and haulm yields (1584 and 3286 kg/ha, respectively), highest yield increase over the untreated control (69.95% for pods and 64.66% for haulms) and the highest net realization (₹48178/ha). The treatments of novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% and chlorpyrifos 50 + cypermethrin 5 EC 0.11% were proved next best treatments over untreated control. Among the various ready-mix insecticidal treatments, chlorpyrifos 50 + cypermethrin 5 EC 0.11% exhibited the highest ICBR of 1:14.3, followed by chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% (1:11.4). However, the treatments viz., profenophos 40 + cypermethrin 4 EC 0.088% (1:4.4), novaluron 5.25 + indoxacarb 4.5 SC 0.016% (1:4.3) and emamectin benzoate 5 + lufenuron 40 WG 0.009% (1:2.8) exhibited lower ICBR.

Keywords: Efficacy, Groundnut, *S. litura*, Ready-mix insecticides

1. INTRODUCTION

Groundnut (*Arachis hypogaea* L.) is a leguminous oilseed crop native to South America. As the king of oilseeds, it is the fourth most important oilseed in the world. It is the largest source of edible oil and ranks 13th among food crops in the world[11]. India comes ranked second to China in terms of groundnut production. Of all the groundnut growing states in India, Gujarat has the largest share in terms of area and production. In Gujarat, the area under groundnut cultivation is 17.09 lakh hectares, with an annual production of 28.14 lakh tonnes and a productivity of 1647 kg/ha[4].

Groundnut yield is affected by direct pest damage or by pests such as disease vectors. More than a hundred insect species have been reported on groundnut in India[2]. Among the various insect pests infesting this crop in Gujarat, the tobacco caterpillar *S. litura* is considered the most important pest due to its polyphagous nature and prevailing favorable climate, it occurs throughout the year. The newly hatched and early instar larvae of *S. litura* feed together on the underside of the leaf, resulting in leaf skeletonization and severe leaf destruction in later stages, leaving only petioles and branches, scraping chlorophyll and causing total yield loss up to 15-30 per cent[8]; more than 180 crops[9].

Comment [SC1]: World

Comment [SC2]: Mention the production of India.

Comment [SC3]: Mention the full genus name.

It is a fact that farmers are depended upon synthetic insecticides for the control of pests and due to injudicious use of chemical insecticides, the pest has developed considerable level of resistance to conventional insecticides including synthetic pyrethroid[5]. Therefore, it is now high time to use certain other pesticides like ready-mix insecticides for its control as well as lesser chance of development of resistance against them.

Comment [SC4]: Need to explain more regarding the objective of the study. Why the ready-mix insecticides were preferred ?

2. MATERIAL AND METHODS

To study the efficacy of various ready-mix insecticides against *Spodopteralitura*(Fab.) in groundnut, the variety, GJG-9 was sown at Main Oilseeds Research Station, Junagadh Agricultural University, Junagadh during the *kharif*, 2023. The experiment was laid out in a Randomized Block Design (RBD) with eight treatments (Table 1) and four replications. Each plot had a gross size of 5 m x 2.7 m and a net size of 4 m x 1.8 m with spacing of 45 cm between rows and 10 cm between plants.

Table 1. Treatment details of insecticides

Sr. No.	Treatment detail	Concentration (%)	Dose (ml or gm/10 l)
1.	Chlorantraniliprole 10 + Lambda-cyhalothrin 5 ZC	0.006	4
2.	Novaluron 5.25 + Emamectin benzoate 0.9 SC	0.009	15
3.	Novaluron 5.25 + Indoxacarb 4.5 SC	0.015	15
4.	Profenofos 40 + Cypermethrin 4 EC	0.088	20
5.	Chlorpyrifos 50 + Cypermethrin 5 EC	0.11	20
6.	Thiamethoxam 12.6 + Lambda-cyhalothrin 9.5 ZC	0.007	3
7.	Emamectin benzoate 5 + Lufenuron 40 WG	0.009	2
8.	Untreated control	-	-

Comment [SC5]: Make a column of trade name of each formulations.

Comment [SC6]: Replace sr. no. with treatments

2.1 Applications of insecticides

All ready-mix insecticides were applied in the form of foliar spray with help of knapsack sprayer. For deciding quantity of spray fluid required per plot, the control plots were spray with water and determined the required spray fluid. Spray fluid was prepared by mixing measured quantity of water and ready-mix insecticide. The necessary care was taken to prevent the drift of insecticides to reach the adjacent plots. The first spray was done at initiation of pest population and need base subsequent application was given at 15 days interval.

Comment [SC7]: Is this the common practice in the field?

2.2 Method of recording observations

To evaluate the efficacy of the ready-mix insecticides, observations of the larval population of *S. litura* were recorded on five randomly selected plants of each treatment before and 5, 10 and 14 days after spraying. Further, obtained data was converted into per cent reduction of *S. litura* population over control through following formula (Abbott, 1925)[1].

$$\text{Per cent reduction over control} = \frac{C - T}{C} \times 100$$

Where,

T = Population of *S. litura* (per plant) from treated plot

C = Population of *S. litura* (per plant) from control plot

Comment [SC8]: How observation is taken from infested leaves? Mention clearly

Comment [SC9]: No need to mention the formula.

2.3 Yield and Economics

With a view to ascertain the effect of various ready-mix insecticides on the groundnut pod and haulm yield, the crop was harvested from each net plot and weigh separately. The harvested pod and haulm yield were converted into kg per hectare. The percent increase yield over control was also calculated by using the following formula mentioned below given by Pradhan (1969)[10].

$$\text{Yield increased(\%)} = \frac{T - C}{C} \times 100$$

Where,

T = Yield of respective treatment (kg/ha)

C = Yield from control (kg/ha)

Economics of all the treatments were worked out by considering the price of groundnut pod and haulm, cost of insecticides used and labour charges for spraying of ready-mix insecticides. Incremental Cost benefit ratio (ICBR) was also work out to compare the economics of various treatments.

$$\text{Incremental Cost Benefit Ratio (ICBR)} = \frac{\text{Net realization(₹/ha)}}{\text{Total cost of insecticide (₹/ha)}}$$

Comment [SC10]: No need to mention the formula.

Comment [SC11]: No need to mention the formula.

3. RESULTS AND DISCUSSION

3.1 First spray

S. litura populations appeared similar in all treatments before spraying, while treatment differences were not significant during the experiment (Table 2). The larval population before spraying ranged from 2.52 to 2.74 larvae per plant. The pooled over period of 5, 10 and 14 days at first spray exhibited that the treatment with chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% was found to be the most effective (0.82 larvae/plant) which was at par with novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% (0.94 larvae/plant). The treatments viz., chlorpyrifos 50 + cypermethrin 5 EC 0.11% and emamectin benzoate 5 + lufenuron 40WG 0.009% were moderately effective, recording larval populations of 1.02 and 1.22 larvae per plant, respectively. While treatment with novaluron 5.25 + indoxacarb 4.5 SC 0.015%, thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007%) and profenofos 40 + cypermethrin 4 EC 0.088%) were found comparatively less effective against *S. litura* as they recorded the larval population of 1.33, 1.39 and 1.45 larvae per plant, respectively. While, the control plot had the highest larval population of 2.33 larvae per plant. The result on interaction effect suggests that there was a significant difference in efficacy of treatment over period due to variable efficacy at different interval tested.

Per cent reduction over untreated control of *S. litura* after the first spray showed that the highest reduction in *S. litura* population compared to the untreated control was chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% (64.95%), followed by novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% and chlorpyrifos 50 + cypermethrin 5 EC 0.11%, with reductions of 59.65 per cent and 56.03 per cent, respectively. In subsequent order, treatments with emamectin benzoate 5 + lufenuron 40 WG 0.009% and novaluron 5.25 + indoxacarb 4.5 SC 0.015% resulted in reductions of 47.55 per cent and 42.70 per cent over the untreated control. The treatment with thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% and profenofos 40 + cypermethrin 4 EC 0.088% recorded the lowest reduction over untreated control (40.11% and 37.63%, respectively).

Table 2. Efficacy of various ready-mix insecticides against *S. litura* in groundnut (First spray)

Sr. No.	Treatment	Concentration (%)	Mean number of larvae per plant				Per cent reduction over untreated control	
			Before spray	Days after spraying				
				5	10	14		
1.	Chlorantraniliprole 10 + Lambda-cyhalothrin 5 ZC	0.006	1.61 (2.58)	0.93 (0.86)	0.77 (0.59)	1.02 (1.03)	0.90 (0.82)	64.95
2.	Novaluron 5.25 + Emamectin benzoate 0.9 SC	0.009	1.60 (2.57)	1.00 (1.00)	0.85 (0.72)	1.06 (1.12)	0.97 (0.94)	59.65
3.	Novaluron 5.25 + Indoxacarb 4.5 SC	0.015	1.64 (2.69)	1.18 (1.38)	1.06 (1.12)	1.23 (1.51)	1.16 (1.33)	42.70
4.	Profenofos 40 + Cypermethrin 4 EC	0.088	1.62 (2.61)	1.22 (1.49)	1.13 (1.27)	1.27 (1.60)	1.21 (1.45)	37.63
5.	Chlorpyrifos 50 + Cypermethrin 5 EC	0.11	1.63 (2.67)	1.04 (1.08)	0.92 (0.84)	1.08 (1.17)	1.01 (1.02)	56.03
6.	Thiamethoxam 12.6 + Lambda-cyhalothrin 9.5 ZC	0.007	1.59 (2.52)	1.20 (1.43)	1.09 (1.19)	1.25 (1.57)	1.18 (1.39)	40.11
7.	Emamectin benzoate 5 + Lufenuron 40 WG	0.009	1.63 (2.65)	1.11 (1.22)	0.99 (1.00)	1.21 (1.45)	1.11 (1.22)	47.55
8.	Untreated control	-	1.66 (2.74)	1.50 (2.26)	1.52 (2.31)	1.56 (2.42)	1.53 (2.33)	-
S. Em. ±			T	0.07	0.05	0.05	0.05	0.03
			P	-	-	-	-	0.02
			T x P	-	-	-	-	0.05
C.D. at 5%			T	NS	0.16	0.15	0.16	0.09
			P	-	-	-	-	0.05
			T x P	-	-	-	-	0.15
C.V. (%)			-	9.05	9.45	10.03	8.74	9.36

Figures in parenthesis are original values, while outsidess are square root transformed values.
NS: Non-significant

3.2 Second spray

S. litura populations appeared similar in all treatments before spraying, while significant differences among treatments were observed during the experiment (Table 3). The larval population before spraying ranged from 1.03 to 2.42 larvae per plant. The pooled over period of 5, 10 and 14 days at second spray, chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% was the most effective treatment (0.60 larvae/plant), which was at par with novaluron 5.25 + emamectin benzoate 0.9 SC 0.009%(0.73 larvae/plant). Moderately effective treatments included chlorpyrifos 50 + cypermethrin 5 EC 0.11% and emamectin benzoate 5 + lufenuron 40 WG 0.009%, which recorded larval populations of 0.84 and 0.95 larvae per plant, respectively. Comparatively less effective treatments against *S. litura* included novaluron 5.25 + indoxacarb 4.5 SC 0.015%, thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% and profenophos 40 + cypermethrin 4 EC 0.088%, with larval populations of 1.03, 1.14, and 1.21 larvae per plant, respectively. The control plot had the highest larval population of 2.54 larvae per plant. The result on interaction effect suggests that there was a significant difference in efficacy of treatment over period due to variable efficacy at different interval tested.

Comment [SC12]: Which treatment is used as the standard check? Where is data regarding per cent reduction over standard check?

Comment [SC13]: Replace with treatment no.

Per cent reduction over untreated control of *S. litura* after the second spray showed that the highest reduction in *S. litura* population compared to the untreated control was chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% (76.42%), followed by novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% and chlorpyrifos 50 + cypermethrin 5 EC 0.11%, with reductions of 71.29 per cent and 67.12 per cent, respectively. In subsequent order, treatments with emamectin benzoate 5 + lufenuron 40 WG 0.009% and novaluron 5.25 + indoxacarb 4.5 SC 0.015% resulted in reductions of 62.73 per cent and 59.33 per cent over the untreated control. The treatment with thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% and profenophos 40 + cypermethrin 4 EC 0.088% recorded the lowest reduction over untreated control (55.02% and 52.46%, respectively).

According to, Gadhiya *et al.* (2014)[7] chlorantraniliprole (0.006%) and emamectin benzoate (0.002%) were noticed higher effective in protecting the groundnut from the infestation of *S. litura*. Anon. (2022b)[3] concluded that the treatments chlorantraniliprole 10 + lambda cyhalothrin 5 15 ZC 0.006% and novaluron 5.25 + emamectin benzoate 0.9 6.15 SC 0.009% were found to be the most effective and economical ready-mix insecticides for management of *S. litura* in groundnut. Bhut *et al.* (2023)[6] indicated that lowest number of larvae/plant was recorded in the treatment of chlorpyrifos 50 + cypermethrin 5 EC 625 + 62.5 a.i./ha which was statistically at par chlorpyrifos 50 + cypermethrin 5 EC 500+50 a.i./ha. Verma *et al.* (2024)[12] revealed that after 1st and 2nd spray, the treatment chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 250 ml/ha was most effective (0.61 larvae/plant and 0.57 larvae/plant) against *S. litura* in cabbage.

Table 3. Efficacy of various ready-mix insecticides against *S. litura* in groundnut (Second spray)

Sr. No.	Treatments	Concentration (%)	Mean number of larvae per plant					Per cent reduction over untreated control	
			Before spray	Days after spraying			Pooled over period		
				5	10	14			
1.	Chlorantraniliprole 10 + Lambda-cyhalothrin 5 ZC	0.006	1.02 (1.03)	0.88 (0.78)	0.76 (0.59)	0.68 (0.46)	0.77 (0.60)	76.42	
2.	Novaluron 5.25 + Emamectin benzoate 0.9 SC	0.009	1.06 (1.12)	0.99 (0.98)	0.82 (0.67)	0.76 (0.57)	0.85 (0.73)	71.29	
3.	Novaluron 5.25 + Indoxacarb 4.5 SC	0.015	1.23 (1.51)	1.10 (1.21)	1.00 (1.00)	0.95 (0.91)	1.02 (1.03)	59.33	
4.	Profenofos 40 + Cypermethrin 4 EC	0.088	1.27 (1.60)	1.19 (1.41)	1.08 (1.16)	1.04 (1.08)	1.10 (1.21)	52.46	
5.	Chlorpyrifos 50 + Cypermethrin 5 EC	0.11	1.08 (1.17)	1.02 (1.05)	0.89 (0.80)	0.83 (0.68)	0.91 (0.84)	67.12	
6.	Thiamethoxam 12.6 + Lambda-cyhalothrin 9.5 ZC	0.007	1.25 (1.57)	1.16 (1.34)	1.05 (1.09)	1.01 (1.02)	1.07 (1.14)	55.02	
7.	Emamectin benzoate 5 + Lufenuron 40 WG	0.009	1.21 (1.45)	1.08 (1.16)	0.96 (0.91)	0.89 (0.79)	0.97 (0.95)	62.73	
8.	Untreated control	-	1.56 (2.42)	1.58 (2.51)	1.60 (2.55)	1.61 (2.58)	1.59 (2.54)	-	
S. Em. ±			T	0.05	0.06	0.05	0.05	0.03	-
			P	-	-	-	-	0.02	-
			T x P	-	-	-	-	0.05	-
C.D. at 5%			T	0.16	0.16	0.15	0.15	0.08	-
			P	-	-	-	-	0.05	-

Comment [SC14]: Replace with treatment no.

	T x P	-	-	-	-	0.15	-
C.V. (%)	-	8.74	9.79	10.05	10.52	10.10	-

Figures in parenthesis are original values, while outsidess are square root transformed values.

3.3 Yield and economics of various ready-mix insecticidal treatments

3.3.1 Yield

The highest pod and haulm yield (1584 and 3286 kg/ha) (Table 4) was obtained from chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% and it was at par with novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% (1451 and 3088 kg/ha). Which was followed by chlorpyriphos 50 + cypermethrin 5 EC 0.11% (1349 and 2951 kg/ha) and emamectin benzoate 5 + lufenuron 40 WG 0.009% (1263 and 2735 kg/ha). Novaluron 5.25 + indoxacarb 4.5 SC 0.015%, thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% and profenophos 40 + cypermethrin 4 EC 0.088% recorded pod yield of 1195, 1119 and 1072 kg/ha, respectively and haulm yield of 2714, 2651 and 2210 kg/ha, respectively, which was notably higher than the pod and haulm yield of the untreated control of 932 and 1996 kg/ha, respectively.

3.3.2 Increase in yield over untreated control

The treatment of chlorantraniliprole 10 + lambda-cyhalothrin 5 ZC 0.006% recorded the highest per cent yield increase (Table 4) for both pods (69.95%) and haulm (64.66%), followed by novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% (55.67 and 54.72%) and chlorpyriphos 50 + cypermethrin 5 EC 0.11% (44.75 and 47.86%). The treatment of emamectin benzoate 5 + lufenuron 40 WG 0.009% noted 35.58 and 37.06 per cent increase in pod and haulm yield over untreated control. Conversely, treatments such as novaluron 5.25 + indoxacarb 4.5 SC 0.015%, thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% and profenophos 40 + cypermethrin 4 EC 0.088% exhibited comparatively lowest per cent increases in both pod and haulm yield over the untreated control.

Table 4. Effectiveness of various ready-mix insecticides on the yield of groundnut pod and haulm

Sr. No.	Treatments	Yield (kg/ha)		Increase in yield over untreated control (kg/ha)		Increase in yield over untreated control (%)	
		Pod	Haulm	Pod	Haulm	Pod	Haulm
1.	Chlorantraniliprole 10 + Lambda-cyhalothrin 5 ZC	1584	3286	652	1291	69.95	64.66
2.	Novaluron 5.25 + Emamectin benzoate 0.9 SC	1451	3088	519	1092	55.67	54.72
3.	Novaluron 5.25 + Indoxacarb 4.5 SC	1195	2714	264	718	28.28	36.00
4.	Profenofos 40 + Cypermethrin 4 EC	1072	2210	140	214	15.03	10.71
5.	Chlorpyriphos 50 + Cypermethrin 5 EC	1349	2951	417	955	44.75	47.86
6.	Thiamethoxam 12.6 + Lambda-cyhalothrin 9.5 ZC	1119	2651	187	656	20.04	32.84
7.	Emamectin benzoate 5 + Lufenuron 40 WG	1263	2735	332	740	35.58	37.06
8.	Untreated control	932	1996	-	-	-	-
	S. Em. ±	66.48	163.03	-	-	-	-
	C.D. at 5%	195.52	479.48	-	-	-	-
	C.V. (%)	10.68	12.06	-	-	-	-

Comment [SC15]: Replace with treatment no.

3.3.3 Economics of various ready-mix insecticides against *S. litura* on groundnut

The economics of various ready-mix insecticides were worked out along with incremental cost benefit ratio (ICBR) (Table 5). The economic analysis of various ready-mix insecticidal treatments showed that chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% recorded the highest net realization of 48,178 ₹/ha followed by novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% (38,676 ₹/ha), chlorpyrifos 50 + cypermethrin 5 EC 0.11% (31,463 ₹/ha). The remaining ready-mix insecticides recorded net realizations ranging from 24879 to 10030 ₹/ha. The plots treated with chlorpyrifos 50 + cypermethrin 5 EC 0.11% showed the highest incremental cost benefit ratio (ICBR) of 1:14.3, followed by chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% (1:11.4), thiamethoxam 12.6 + lambda-cyhalothrin 9.5 ZC 0.007% (1:10.1) and novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% (1:9.7). Rest of the treatments viz., profenophos 40 + cypermethrin 4 EC 0.088% (1:4.4), novaluron 5.25 + indoxacarb 4.5 SC 0.016% (1:4.3) and emamectin benzoate 5 + lufenuron 40 WG 0.009% (1:2.8) exhibited lower ICBR and were deemed less economically viable. According to Anon. (2022b)[3] the treatment of chlorantraniliprole 10 + lambda cyhalothrin 5 15 ZC 0.006% recorded significantly highest pod and haulm yield (1561 and 2688 kg/ha) with highest net return (22288 ₹/ha), while, novaluron 5.25 + emamectin benzoate 0.9 6.15 SC 0.009% and chlorpyrifos 50 + cypermethrin 5 55 EC 0.11% were the next best treatments. Whereas, in point of ICBR treatment chlorpyrifos 50 + cypermethrin 5 55 EC 0.11% noted highest ICBR i.e. (1:6.34) followed by the treatment chlorantraniliprole 10 + lambda cyhalothrin 5 15 ZC 0.006% (1:5.89) and novaluron 5.25 + emamectin benzoate 0.9 6.15 SC 0.009% (1:4.79). Thus, the results of the present findings are more or less close agreement with earlier findings.

Table 5. Economics of various ready-mix insecticides against *S. litura* on groundnut

Tr. No.	Quantity of insecticide required for two spray (l or kg/ha)	Cost of insecticide for 2 spray (₹/ha)	Total cost of treatment (₹/ha)	Yield (kg/ha)		Gross realization (₹/ha)	Net realization (₹/ha)	ICBR
				Pod	Haulm			
T ₁	0.40	3240	4240	1584	3286	117806	48178	1:11.4
T ₂	1.50	3000	4000	1451	3088	108304	38676	1:9.7
T ₃	1.50	3750	4750	1195	2714	90050	20422	1:4.3
T ₄	2.00	1300	2300	1072	2210	79658	10030	1:4.4
T ₅	2.00	1200	2200	1349	2951	101091	31463	1:14.3
T ₆	0.30	510	1510	1119	2651	84871	15243	1:10.1
T ₇	0.20	7995	8995	1263	2735	94507	24879	1:2.8
T ₈	-	-	-	932	1996	69628	-	-

Labour charge @ 500 ₹/spray/ha

Market value of ground pod @ 64 ₹/kg and market value of groundnut haulm @ 5 ₹/kg

4. CONCLUSION

Based on the present investigation, it can be concluded that chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% proved to be the most effective against *S. litura*. It exhibited the lowest larval population, maximum pod and haulm yield, highest pod and haulm yield increase over the untreated control and the highest net realization. While, the treatments of

novaluron 5.25 + emamectin benzoate 0.9 SC 0.009% and chlorpyrifos 50 + cypermethrin 5 EC 0.11% were proved next best treatments over untreated control. Among the various ready-mix insecticidal treatments, chlorpyrifos 50 + cypermethrin 5 EC 0.11% obtained the highest ICBR of 1:14.3, followed by chlorantraniliprole 10 + lambda cyhalothrin 5 ZC 0.006% with ICBR of 1:11.4.

Comment [SC16]: What could be your advise to groundnut growers?

REFERENCES

1. Abbott WS. A method of computing the effectiveness of insecticides. *Journal of Economic Entomology*. 1925;18(2):265-267.
2. Amin PW. Insect and mite pests and their control. In P. S. Reddy (ed.) *Groundnut*, I.A.R.C., New Delhi. 1988;393-452.
3. Anonymous. Annual Research Report of Oilseed crops, Main Oilseeds Research Station, Junagadh Agricultural University, Junagadh, 2022b;234-240.
4. Anonymous. Groundnut crop survey report. 2022a. Accessed 9 December 2023. Available at [https://apeda.gov.in/apedawebsite/HACCP/2022_Groundnut_Survey_Report.pdf].
5. Armes NJ, Bond GS and Cooter RJ. The Laboratory culture and development of *Helicoverpa armigera*. *Natural Resources Institute Bulletin No. 57*. Natural Resources Institute, Chatham, UK. 1992;15.
6. Bhut JB, Bharadiya AM and Madariya RB. Evaluation of chlorpyrifos 50% + cypermethrin 5% EC against lepidopteran defoliators *Helicoverpa armigera* L. and *Spodoptera litura* Fab. on soybean. *Journal of Oilseeds Research*. 2023;40:336-337.
7. Gadhya HA, Borad PK and Bhut JB. Effectiveness of synthetic insecticides against *Helicoverpa armigera* (Hubner) Hardwick and *Spodoptera litura* (Fabricius) infesting groundnut. *The Bioscan*. 2014;9(1):23-26.
8. Ghewande MP and Nandagopal V. Integrated pest management in groundnut *A. hypogaea* L. in India. *Integrated Pest Management Reviews*. 1997;2:1-15.
9. Isman MB, Machial CM, Miresmaili S and Bainard LD. Essential oil based pesticides: new insights from old chemistry. In: *Pesticide chemistry*, H. Ohkawa and H. Miyagawa, (Eds): Wiley, Weinheim. 2007;113.
10. Pradhan S. *Insect pests of crops*. National Book Trust, New Delhi, India. 1969;80.
11. Ramanathan T. *Genetic improvement of groundnut*. Associated Publishing Company, XI. 2001;260.
12. Verma RK, Singh SK, Sharma KR, Chandra U and Yadav PK. Bio-efficacy of chlorantraniliprole 10% + lambda-cyhalothrin 5% ZC and some sole insecticides against lepidopteran borers in cabbage. *Annals of Plant and Soil Research*. 2024;26(1):136-142.