

Yield of lettuce fertilized with hairy woodrose (*Merremiaegyptia* L.) plus poultry manure.

ABSTRACT

The use of plant resources available in agricultural areas is extremely important for farmers who work in this activity, contributing to an economic return in lettuce production. In view of the above, the objective was to study the yield of lettuce fertilized with hairy woodrose (*Merremiaegyptia* L.) plus poultry manure. The experiment was carried out in a completely randomized design (D.I.C) with six treatments and four replications. The treatments consisted of six amounts of the hairy woodrose (*Merremiaegyptia* L.) mixture plus poultry manure (0; 1.4; 2.8; 4.2; 5.6 and 7.0 kg m⁻²). For the lettuce crop, the cultivar "Babá de Verão". Twenty-two days after transplanting, the experiment was harvested where the plants were transported to the Vegetable Post-Harvest Laboratory of the Department of Agronomic and Forestry Sciences at UFERSA, where the following characteristics were analyzed: plant height (performed from a sample of four plants plot⁻¹, measuring the height from the base to the inflection of the leaves using a millimetric ruler, expressed in cm plant⁻¹); number of leaves per plant (determined in a sample of ten plants, counting the number of leaves per plant over five centimeters, expressed in plant⁻¹ units) fresh lettuce mass (measured by the weight of all plants in the plot on an electronic scale with an accuracy of 1.0 g, expressed in g plant⁻¹); lettuce dry mass (was carried out by weighing four plants on an electronic scale with an accuracy of 1.0 g, which were then placed in a heating oven with forced air at 65 °C, until constant mass). The amount of 7.0 kg m⁻² of the mixture of hairy woodrose (*Merremiaegyptia* L.) plus poultry manure was what promoted the greatest increase in lettuce cultivation, with a maximum value of 237.97 g plant⁻¹. The mixture of alternative fertilizer sources is extremely important in the production of leafy vegetables.

Keywords: Leafy vegetable; organic fertilizer and monoculture.

1. INTRODUCTION

Vegetable production is intense in semi-arid regions, where manure (bovine, goat and poultry) is used exclusively. However, the farmer does not always have this resource available on his property, which increases the cost of vegetable production [1], [2]. Among the vegetable crops produced in the region of Mossoró, RN, Brazil, lettuce (*Lactuca sativa*) stands out [3]. This type of vegetable is the most sold and consumed in Brazil, with low caloric value and as a source of minerals [4].

In this context, the use of a mixture of green manures with sources of fertilizers of animal origin is extremely important, which provides greater availability of nutrients in the soil, conditioning the soil environment for the cultivation of vegetable crops [3].

In the semi-arid region of Brazil, the presence of species with the potential to be used as green manure is quite prominent, notably hairy woodrose (*Merremiaegyptia* L.), which produces green and dry biomass in the order of 40,000 and 6,000 kg ha⁻¹, respectively, with of nitrogen concentration of 22.4 g kg⁻¹ in dry matter and a carbon-nitrogen ratio of 17/1, at the phenological stage of 126 days after emergence [5].

This species has been used as green fertilizer in the development of leafy and root vegetables, as a source of nitrogen in arugula and coriander [6].

Given the importance of using resources present within agricultural areas, as a source of nutrients in the fertilization of vegetables, the objective was to study the yield of lettuce fertilized with hairy woodrose (*Merremiaegyptia* L.) plus poultry manure.

2. MATERIAL AND METHODS

2.1 Location of the experiment installation

The experiment was carried out in a greenhouse in the teaching garden of the Department of Agricultural and Forestry Sciences at the Universidade Federal Rural do Semi-árido (UFERSA), Mossoró, RN, Brazil, from December 2022 to January 2023, in classified soil as Red Yellow OxisolArgissolic sandy loam [5]. According to [6] and the Köppen classification, the local climate is BSw^h, dry and very hot, with a dry season, often from June to January, and a rainy season from February to May, average annual precipitation of 673.9 mm and average relative humidity of 68.9%.

Before setting up the experiment, soil samples were collected from the 0-20 cm arable layer, homogenized, and sent to the soil chemistry and fertility laboratory for the following analyses: pH (water) = 6.8; EC = 0.30 dS m⁻¹; O.M. = 3.6 g kg⁻¹; N = 0.75 g kg⁻¹; P = 37.81 mg dm⁻³; K = 69.45 mg dm⁻³; Ca = 47.65 cmolc dm⁻³; Mg = 5.14 cmolc dm⁻³; Na = 1.93 mg dm⁻³; Cu = 0.45 mg dm⁻³; Fe = 2.26 mg dm⁻³; Mn = 8.94 mg dm⁻³; and Zn = 3.24 mg dm⁻³.

2.2 Experimental Design

The experiment was carried out in a completely randomized design (D.I.C) with six treatments and four replications. The treatments consisted of six amounts of the hairy woodrose (*Merremiaegyptia* L.) mixture plus poultry manure (0; 1.4; 2.8; 4.2; 5.6 and 7.0 kg m⁻²). For the lettuce crop, the cultivar "Babá de Verão" (Incomplete sentence).

Four holes were opened in each pot, and the plants were placed in the experimental plots. During the period in which the cowpea crop was in the field, weeding was carried out to remove invasive plants that compete for water and nutrients during the crop cycle. Irrigation was carried out by drip, with a daily irrigation shift divided into two applications (morning and afternoon).

To compose the mixture of fertilizers in the research hairy woodrose (*Merremiaegyptia* L.) was used, a spontaneous species from the semi-arid region with production of green and dry phytomass of the order of 42,000 kg ha⁻¹ and 6,000 kg ha⁻¹, respectively, with nitrogen content of 24.7 g kg⁻¹ at 104 days after emergence (LINHARES et al., 2021).

The hairy woodrose (*Merremiaegyptia* L.) was harvested in an area of semi-arid vegetation adjacent to the Federal Rural University of the Semi-arid 100 days after emergence, being crushed in forage into segments of 2.0 to 3.0 centimeters (Figure 1). The material was then dried in the sun for a period of 80 hours until the moisture content was 15%, after which samples were taken and sent to the

soil fertility and plant nutrition laboratory at the Center for Agricultural Sciences at UFERSA for carbon analysis. (W); nitrogen (N); phosphorus (P); potassium (K⁺); calcium (Ca²⁺); magnesium (Mg²⁺) and carbon/nitrogen ratio, whose values were: 535 g kg⁻¹ C, 23.5 g kg⁻¹ N, 10.8 g kg⁻¹ P, 15.4 g kg⁻¹ K, 9, 7 g kg⁻¹ Ca, 11.7 g kg⁻¹ Mg and a nitrogen/carbon ratio of 23/1.

Poultry manure was collected from the poultry sector of the Department of Animal Sciences at UFERSA, from laying poultry farming and sent to the soil fertility and plant nutrition laboratory at the Center for Agricultural Sciences at UFERSA for carbon (C) analysis. ; nitrogen (N); phosphorus (P); potassium (K⁺); calcium (Ca²⁺); magnesium (Mg²⁺) and carbon/nitrogen ratio, whose values were: 440 g kg⁻¹ C, 28.7 g kg⁻¹ N, 12.6 g kg⁻¹ P, 17.3 g kg⁻¹ K, 16, 9 g kg⁻¹ Ca, 13.2 g kg⁻¹ Mg and a carbon/nitrogen ratio of 15/1.

2.3 Agronomic characteristics of lettuce crop

Twenty-two days after transplanting, the experiment (Use appropriate word like crop) was harvested where the plants were transported to the Vegetable Post-Harvest Laboratory of the Department of Agronomic and Forestry Sciences at UFERSA, where the following characteristics were analyzed: plant height (performed from a sample of four plants plot⁻¹, measuring the height from the base to the inflection of the leaves using a millimetric ruler, expressed in cm plant⁻¹); fresh lettuce mass (measured by the weight of all plants in the plot on an electronic scale with an accuracy of 1.0 g, expressed in g plant⁻¹); lettuce dry mass (was carried out by weighing four plants on an electronic scale with an accuracy of 1.0 g, which were then placed in a heating oven with forced air at 65 °C, until constant mass).

Statistical analysis

Statistical analysis was performed according to conventional methods of analysis of variance [7], using ESTAT statistical software [8]. The response curve fitting procedure was performed using the ESTAT Software [8], applying regression analysis and conducting hypothesis testing that helps the researcher accept or reject a statistical hypothesis based on experimental results [9], [10].

3. RESULTS AND DISCUSSION

It was observed that there was a significant effect on the characteristics of plant height, number of leaves, green and dry mass of lettuce depending on the amounts of hairy woodrose (*Merremia aegyptia* L.) plus poultry manure (Figure 2 to 5). The mixture of alternative fertilizer sources is extremely important for soil fertilization in vegetable cultivation (LINHARES et al., 2024).

For plant height, there was a maximum point with a value of 14.63 cm plant in the amount of 5.8 kg m⁻² of the mixture of hairy woodrose (*Merremiaaegyptia* L.) plus poultry manure (Figure 2). It is worth

noting that, as it is a hardwood, it is greatly influenced by the addition of organic material rich in nitrogen. Linhares et al. (2024) studying Productivity of lettuce with different amounts of the mixture of scarlet starglory (*Merremiaegyptia* L.) with rooster tree (*Calotropis procera*) applied in soil cover found a plant height of 15.43 cm/plant with an application of 3.5 kg m⁻² of the mixture of scarlet starglory and rooster tree, higher than the aforementioned research. Bezerra Neto et al. (2011), studying different quantities and decomposition times of scarlet starglory (*Merremiaegyptia* L.) in lettuce, found an average height of 21.14 cm plant⁻¹, using spacing between plants greater than the aforementioned research.




Figure 2. Lettuce plant height as a function of different amounts of *Merremiaegyptia* mixture with poultry manure.

In relation to the number of leaves, the different amounts of the mixture of hairy woodrose (*Merremiaegyptia* L.) plus poultry manure had a significant influence, with a maximum value of 8.21 plant units, in the amount of 4.2 kg m (Figure 3). According to Linhares et al. (2024) the number of leaves is extremely important, being the vegetative part where photosynthesis takes place. Moura et al. (2020), studying the agronomic development of lettuce crops fertilized with organic compost, obtained a maximum number of 11.3 leaves, lower than the aforementioned research.




Figure 3. Number of leaves as a function of different amounts of *Merremiaegyptia* mixture with poultry manure.

Regarding the fresh and dry mass of lettuce, the observed data fit an increasing linear equation, with maximum values of 237.97 and 19.10 g plant⁻¹ in the amount of 7.0 kg m⁻², respectively (Figure 4 and

5). Peixoto Filho [et al.](#) (2013), studying lettuce productivity with doses of chicken, cattle and sheep manure in successive crops, found a maximum lettuce weight of 296.7 g plant⁻¹, different from the present research. Santos [et al.](#) (2021) studying the productivity of *Lactuca sativa*, cultivated under doses of organic compost and biofertilizer, found a maximum weight of fresh lettuce mass of 71.72 g plant⁻¹ at a dose of 9.0 kg m⁻² of biofertilizer lower than the [aforementioned](#) research. Linhares [et al.](#) (2011), evaluating different periods of incorporation of hairy woodrose (*Merremiaegyptia* L.) in lettuce, found a maximum value of 64.7 g plant, lower than the aforementioned research. According to Silva [et al.](#) (2011) organic fertilizer increases productivity, in addition to producing plants with better qualitative characteristics to the detriment of plants grown exclusively with mineral fertilizers. For Sediyaama [et al.](#) (2016) organic fertilizer is a viable alternative, with great potential for lettuce cultivation.




Figure 4. Lettuce green mass as a function of different amounts of *Merremiaegyptia* mixture with poultry manure.




Figure 5. Lettuce dry mass as a function of different amounts of *Merremiaegyptia* mixture with poultry manure.

CONCLUSION

The amount of 7.0 kg m⁻² of the mixture of hairy woodrose (*Merremiaegyptia* L.) plus poultry manure was [what](#) promoted the greatest increase in lettuce cultivation, with a maximum value of 237.97 g plant⁻¹.

The mixture of alternative fertilizer sources is extremely important in the production of leafy vegetables.

REFERENCES

1. Linhares PCF, Assis JP, Sousa RP, Sá JR, Pereira MFS, Ramalho WB, Silva RIG, Silva RA, Pereira KLV (2018). Optimized amount of hairy woodrose (*Merremiaegyptia* L.) in the productivity of coriander cultivars. *Bulgarian Journal of Agricultural Science*. 24(4): 654-659. <https://www.agrojournal.org/24/04-17.pdf>
2. Linhares PCF, Maracajá PBM, Pereira FS, Assis JP and Sousa RP. Rooster tree (*Calotropis procera*) under different amounts and periods of incorporation on yield of coriander. *Green Magazine of Agroecology and Sustainable Development*, 2014; 9(3): 07-12. <https://www.gvaa.com.br/revista/index.php/RVADS/article/view/2779>
3. Linhares PCF, Sousa JDAS, Maracajá PB, De Medeiros AC, Alves LDeS, Da Silva UL, Carlos KGDaS, De Souza Júnior DS (2023). Coriander yield as a function of green manure incorporation of hairy woodrose (*Merremiaegyptia* L.), rooster tree (*Calotropis procera*) and kills pasture (*Senna uniflora* L.) in a semiarid region of Brazil. *DELLOS: DESARROLLO LOCAL SOSTENIBLE*, 16 (46), 2370–2385. <https://ojs.revistadelos.com/ojs/index.php/dellos/article/view/977>
4. Figueira F.A.R. 2013. Vegetable farming manual: Modern agrotechnology in the production and marketing of vegetables. UFV: Viçosa. 402p.
5. Linhares PCF, Maracajá PB, Liberalino Filho J, Assis JP, Sousa RP, Medeiros AC. Hairy woodrose (*Merremiaegyptia* L. Urban) [electronic book]: Potential use as a spontaneous species in the semi-arid region in the green management of vegetables In: Linhares PCF, Cunha LMM, Silva NV, Neves AM, Medeiros BBM and Paiva B.C. Green and dry phytomass, levels and accumulation of macronutrients in hairy woodrose (*Merremiaegyptia* L. Urban) at different phenological stages – Nova Xavantina, MT: Ed. Pantanal. 96p. 2021; Chapter 2, p.24-45. <https://doi.org/10.46420/9786588319901>
6. Linhares PCF (2009). Spontaneous vegetation as green manure in the agroeconomic performance of leafy vegetables. Department of Agricultural and Forestry Sciences at the Federal Rural University of Semi-Árido (Thesis), Mossoró. 109p.
7. BRAZILIAN AGRICULTURAL RESEARCH COMPANY – EMBRAPA (2018). Brazilian system of soil classification. 2nd ed. Rio de Janeiro: Embrapa, 306p.
8. Banzatto DA, Kronka SN. Agricultural experimentation. 3rd ed. Jaboticabal: FUNEP, 1995:245p.
9. Barbosa JC, Malheiros EB, Banzatto D. A. ESTAT: A system for statistical analysis of agronomic trials. Jaboticabal: Unesp, Version 2.0. 1992.
10. Assis JP, Sousa RP, & Linhares PCF (2020). Statistical hypothesis testing. EdUFERSA. <https://livraria.ufersa.edu.br/wp-content/uploads/sites/165/2020/08/testes-de-hipoteses-estatisticas-edufersa.pdf>

11. Assis JP (2013). Simple linear regression, simple linear correlation, multiple linear regression and multiple linear correlation. EdUFERSA, 310p.
<https://livraria.ufersa.edu.br/regressao-e-correlacao-linear-simples-e-multipla/>
12. Linhares PCF, Sousa JDAS, Maracajá PB, De Medeiros AC, Alves LDeS, Da Silva UL, CARLOS, K. G. DA S.; DE SOUZA JÚNIOR, D. S. Coriander yield as a function of green manure incorporation of hairy woodrose (*Merremiaegyptia* L.), rooster tree (*Calotropis procera*) and kills pasture (*Senna uniflora* L.) in a semiarid region of Brazil. *DELOS: DESARROLLO LOCAL SOSTENIBLE*, 16 (46), 2370–2385.2023. <https://ojs.revistadelos.com/ojs/index.php/delos/article/view/977>
13. Linhares PCF, Silva MLdaS, Silva JdosS, Holanda ARde, Silva ULda (2009). InfluenceAgronomic performance of lettuce in different quantities and decomposition times of green jitrana. *Brazilian Journal of Agricultural Sciences*, 6(2): 236-242.
15. Moura AQ, Correa EB, Fernandes JD, Monteiro Filho AF, Leão AC, Boava LP (2020). Agronomic efficiency of lettuce fertilized with different organic compounds. *Brazilian Journal of Sustainable Agriculture*, 10(1): 155-163.
16. Peixoto Filho JU, Freire MBGdosS, Freire FJ, Miranda MFA, Pessoa LGM, Kamimiera KM (2013). Lettuce productivity with doses of chicken, cattle and sheep manure in successive crops. *Brazilian Journal of Agricultural and Environmental Engineering*, 17(4): 419-424.
17. Santos AGdos, Mesquita JB, Pessoa AMdosS, Silva LMda (2021). Productivity of *Lactuca sativa* L. grown under doses of organic compost and biofertilizer. *Brazilian Journal of Development*, 7(1), 8481–8496. <https://doi.org/10.34117/bjdv7n1-575>
18. Silva EMNCP, Ferreira RLF, Araújo Neto SE, Tavella LB, Solino AJS (2011). Quality of curly lettuce grown in organic, conventional and hydroponic systems. *Brazilian Horticulture*, 29(1): 242-245.
19. Sediya MAN, Magalhães IPB, Vidigal SM, Pinto CLO, Cardoso DSCP, Fonseca MCM, Carvalho IPL (2016). Use of organic fertilizers in the cultivation of iceberg lettuce (*Lactuca sativa* L.) 'KAISER'. *Brazilian Journal of Sustainable Agriculture*, 6(2): 66-74.