

Original Research Article

Biology of Muga Uzi Fly: The Menace threatening the Muga Industry

ABSTRACT

The experiment was carried out with the purpose to thoroughly study the biology of a very serious endo-larval parasitoid of muga silkworm, *Blepharipa sp.* (Walker). The study was done in experiment field where the muga silkworms were reared under net condition and were allowed to infest by the uzi fly, whereas the study of biology was carried out in laboratory condition at room temperature and humidity. The study regarding biology revealed that the life cycle of *Blepharipa sp.* (Walker) is completed in 30 days on average for male and 33 days for female fly which consist an average of 2.40 days of incubation period, 6.20 days larval period, 12.40 days pupal period and adult longevity was 10.80 days in case of male and 12.00 days in case of female. Other aspects of their life cycle regarding peak infestation season, site of oviposition were also studied. The mortality due to uzi fly infestation was 100% as the worms died either before spinning stage or died inside their cocoons and regarding the cocoons they produced, almost all were defective, not suitable for reeling. Since this is the most destructive among all insect pests of muga silkworm causing not only economic losses but also destroying farmers motivation, a thorough study of the biology was of utmost importance.

Keywords: *Antheraea assamensis*, *Blepharipa sp.*, Major pest, Pest biology

Commented [A1]: Remove this and add only to

Commented [A2]: Start as a new sentence

1. Introduction

India's diverse natural environment, conducive to the production of all four types of commercial silk, bestows upon it a distinctive presence in sericulture. Of them is muga silkworm which, being multivoltine and endemic to Assam and some other north-eastern states, are typically reared outdoors, with cocoon spinning to seed production occurring indoors, rendering them vulnerable to pest infestations. Among these pests, the uzi fly poses a significant threat, leading to yield reduction, particularly during the peak infestation period i.e. the chotua crop, a pivotal commercial seed crop within the muga cultivation. They causes loss of silkworm crops which hampers the generation of future seed crops hence disrupts the commercial seed crop production which ultimately causes loss in silk production (Goswami et al., 2013). The uzi fly is a member of the family Tachinidae, most of which are exclusive parasitoids, primarily targeting Lepidoptera but also known to attack different stages of other insects such as Heteroptera, Coleoptera and Symphyta of Hymenoptera (Dai et al., 2022). The muga silkworm are known to be parasitized by two Tachinid species viz. *Exorista sorbillans* and *Blepharipa sp.*. Among them, *B. sp.* is identified as the primary Tachinid fly that preys on muga silkworms where the peak period of attack occurs between December and April, with reported losses ranging from 48.7% to 80% in various seed areas across Assam (Choudhury et al., 2014). The first report of *Blepharipa sp.* as a serious pest of muga silkworm was reported in the year 1989 by Goswami and Barah. The infestation of uzi fly is recognizable by the black sear where the attack occurred, and maggots emerging from the cocoon are evident through holes in it making them inappropriate for reeling. Muga silkworms, having six crops annually, are raised year-round. Due to their close synchronization with the uzi fly life cycle, it's challenging to rear them without infestation. Additionally, outdoor rearing increases the risk of crop loss due to other pest and diseases, compounded by different climatic conditions throughout the year. Managing the infestation involves employing nylon nets to cover the host plants, which is difficult because farmers typically don't prune their fields, resulting in large trees that cannot be completely covered by nets. The current study aimed to investigate the biology and damage symptoms by the pest as well as to promote simple yet effective cultural and hygienic practices for its management.

2. Materials and methods

The study of uzi fly life cycle was conducted in field condition at experimental field of Department of Sericulture, AAU, Jorhat, Assam (Lat 26.721544°, Long 94.196678°) with temperature, humidity and rainfall levels ranging from 13.48°C-26.73°C, 77%-78.5% and 0mm-86mm. To calculate the oviposition and fecundity of the uzi fly, muga silkworm larvae were infested artificially by keeping them with mated female uzi fly in field and were properly covered with nylon net to prevent the uzi fly from flying out. Later part of the study regarding pupal period and adult emergence were done in laboratory condition at room temperature and humidity by keeping the cocoons of infested muga silkworm each separately in different plastic containers and was made sure to carry out everyday observation. The measurements of dimensions of egg, maggot, pupa and adult were carried out with the help of a binocular stereoscope.

3. Result

The female fly after copulation search for the suitable host for oviposition which apparently found to be mostly the late instars of the muga silkworm i.e. 4th and 5th instars. They attach the eggs to the exposed integument of silkworm larvae mostly in the intersegmental region, although they were also found to lay eggs on exposed sides of prolegs, near spiracles etc. With an average oviposition rate of 35.20 numbers per day (table 3) during the experimental period, the female fly was spotted to lay 7-16 eggs per silkworm. The eggs were found to have hatched through chorion on the antero-ventral surface, with an observed hatching percentage of 84.48%, following an incubation period of 2-3 days (table 2). The measurements of egg, maggot, pupa and adult of *B. sp.* are depicted in table 1.

3.1. Egg stage of *Blepharipa sp.*

It was found that the eggs were ovoid shaped and of macro type (fig.1). The surface of the egg was relatively flat ventrally with an opaque chorion and convex dorsally. Although the gravid adult female uzi flies appeared to lay their eggs anywhere on the larval body of muga silkworm, it became apparent that the lateral region was the most desired location, subsequently followed by both ventral and dorsal site with percent infestation of 50.83%, 25.42% and 23.97% respectively (fig. 2-a, b, c) (table 4).

3.2. Larval stage of *Blepharipa sp.*

With a nearly spherical posterior region and a tapered anterior region, the apodous and fusiform morphology of the maggots were typified by their curved hooked mouth part displaying a total of 12 body segmentation. The third instar matured maggots emerged from the host body to undergo pupation meanwhile completing their all three larval instars inside the host body.

First instar maggot: The initial instar maggots (fig. 3a) were minuscule, off white in appearance, and possessed two tiny, dot-like spiracles at posterior body region. They were linked to the respiratory funnel (fig. 8) that had developed

Commented [A3]: cause

Commented [A4]: can mention this in abstract

Commented [A5]: fig 8 missing check

Commented [A6]: all their

at the puncture site and did not move much, staying close to it and they do it by gathering a mass of fat bodies near the entrance point inside the infested silkworm larvae (fig. 3b)

Second instar maggot: The maggots in their second instar (fig. 4) had a gentle white appearance with two light brown spiracles at the posterior end. Additionally, they also exhibited sedentary behavior and remained close to the puncture site in the host larval body.

Third instar maggot: The maggots in their third instar were dirty white to yellowish white in colour (fig. 5), having two noticeable spiracles with sclerotized peritreme. The third instar uzi maggots go throughout the entire host body, eating voraciously on the fat content of the host larva. They exited the host larval body by piercing the dead host's body wall at maturation.

3.3. Pupal stage of *Blepharipa* sp.

After emerging from the silkworm larval body, the mature uzi maggots searched for a suitable location to pupate. The maggots preferred to pupate in soil although they typically hunt for fissures and crevices in their surroundings. The freshly produced pupae had a light brown colour (fig 6a) that eventually grew darker with time (fig. 6b). These coarctate pupae comprised of 11 body segments. After the pupal stage, which lasted an average of 12.40 days (table 2), the adult fly emerged from the pupa by breaking the predetermined line of weakness at the anterior region of the pupa. The adults appeared to emerge most often in the morning, when the temperature appeared to be at its best. Temperature and rainfall had a significant impact on adult emergence; greater emergence was detected following rainy days and a greater infestation percentage was found in Chotua crop of muga silkworm (Feb-Mar). With an average emergence percentage of 74.00, the emergence persisted for three to four days (table 2).

3.4. Adult stage of *Blepharipa* sp.

The adult uzi flies had four longitudinal lines of bristles on the thoracic notum and long lateral bristles on the lateral abdominal area. They were blackish in appearance with an orange tinge at the dorso-lateral region of the abdomen. It was evident that the female was a bit bigger over the male. Mating mostly took place on the day of emergence (fig. 7-a,b).

4. Discussion

Blepharipa sp. is without a doubt the most destructive pest resulting in significant losses in the muga industry among all the insect pests of muga silkworm in Assam and Meghalaya states of India (Kabiraj *et al.*, 2022). Despite being most destructive, there is little information accessible regarding the biology of the muga uzi fly. Reddy and Ranjan (2011) reported higher infestation in Chotua crop (43.0%) than Jarua crop (19.0%) of muga silkworm in upper Assam. According to Manjunatha and Puttaraju (1993), the eggs of uzi fly were also of macrotype, dull white in colour, almost oval in shape with a slightly pointed proximal region. Patil and Savanurmath (1989) reported that usually the eggs of *B. sp.* hatched 3 days after oviposition and the hatching percentage was as high as 97.9% when observed under lab condition. They also revealed that the *B. sp.* preferred late instar worms of *Antheraea mylitta* D., mainly the 4th instar worms over the early instar worms. *B. sp.* were reported to infest hosts on the basis of host-stage, vigour and health of the silkworm. Ramprakash and Kshirsagar (2019) found that the maggot period of *Exorista bombycis* was found to be 5-7 days. The size of uzi flies attacking muga silkworm and mulberry silkworm also varies. Maggots of uzi flies targeting mulberry silkworms were smaller compared to those attacking muga silkworms and as reported these variations may stem from the larger body size of muga silkworm (Thangavelu and Sahu, 1986). Thangavelu and Sahu (1986) also stated that the pupa of uzi fly were photonegative and undergo pupation mostly in soil but sometimes the pupation may occur inside the host larval cocoon. Goswami *et al.* (2013) reported that the maggot of uzi fly infesting muga silkworm after maturation and coming out of the silkworm body pupate within 7-8 hours and the pupal period was found to be 10 days. Negi *et al.* (1993) stated that the total life span of *B. sp.* was 48.1-64.1 days in an alternate host, *Cricula trifenestrata*.

5. Conclusion

Various reports regarding the infestation of a very destructive Tachinid pest of muga silkworm were recorded in Assam causing devastating economical loss. This study was done to provide a thorough study and familiarize the biology of the pest i.e. uzi fly. The uzi flies were found to mostly attack the muga silkworm during Chotua crop (Feb-Mar) although the infestation occurs throughout the year. Starting from egg stage to adult, the life cycle of the male uzi fly is 30 days long while the female has longer life span of 33 days on average. The adult female finds the lateral region of muga larva to be most favorable for egg laying which hatched within an incubation period of 2-3 days (table 2). The hatching of egg is characterized by a black scar forming near the egg which occurred due to the puncture made by the maggot while entering the host larval body (fig. 9). The shorter egg incubation & larval period which definitely the destructive period makes it very difficult to control which is aided by the fact that they are endo-larval parasitoid,

Commented [A7]: remove this

Commented [A8]: remove this –don't give in conclusion

Commented [A9]: remove

Commented [A10]: remove here and add in result fig 9, 10.

hence once the host is infested, they will surely die. Even if the host larvae managed to survive till cocoon spinning, those cocoons will be either flimsy, small, deformed or not fully formed (fig. 10). The resulting losses sometimes include the whole crop loss which renders the production of next generation progeny.

Commented [A11]: remove from here and add in result

References

1. Goswami, N. K.; Nath, P. and Saharia, D (2013). Uzi Fly Infestation Severity in Muga Seed Cocoons, *Antheraea assamensis* Helfer and Crop Loss During Chotua Crop in Assam. *Indian Journal of Applied Research*.3(10): 1-4.
2. Kabiraj, D.; Chetia, H.; Nath, A.; Sharma, P.; Mosahari, P.; Singh, D.; Dutta, P.; Neog, K. and Bora, U. (2022). Mitogenome-wise codon usage pattern from comparative analysis of the first mitogenome of *Blepharipa* sp. (Muga uzifly) with other Oestroid flies. *Scientific Reports*. 12(1): 7028.
3. Dai, M.; Yang, J.; Liu, X.; Gu, H.; Li, F.; Li, B. and Wei, J. (2022). Parasitism by the tachinid parasitoid *Exorista japonica* leads to suppression of basal metabolism and activation of immune response in the host *Bombyx mori*. *Insects*. 13: 792.
4. Choudhury, B. and Kumar, R.; Chutia, P. and Rajkhowa, G. (2014). Host infestation Potentiality of *N. thymus* to Control the Uzifly of muga silkworm, *Antheraea assamensis* (Helfer) - A Bio-control Tool for Controlling Uzifly. *Biological Forum – An International Journal*,6(1): 1-4.
5. Goswami, M. C. and Barah, A. (1989). Report of *Blepharipa* sp. Walker (Tachinidae) as a serious pest of muga silkworm, *Antheraea assama* Westwood (Saturniidae). *Current Science*,58 (5): 267-268.
6. Reddy, S. G. E. and Ranjan, R. K. (2011). present status of uzi fly, *Exorista bombycis* (Louis) (Diptera: Tachinidae) incidence on muga silkworm, *Antheraea assamensis* Helfer (Lepidoptera: Saturniidae) in upper Assam. *Munis Entomology & Zoology*.6(2): 856-858.
7. Manjunatha, H. and Puttaraju, H. P. (1993). The Egg of Uzi Fly, *Exorista sorbillans* (? *E. bombycis* Louis) (Diptera: Tachinidae). *Appl. Entomol. Zool.*,28(4): 574-577.
8. Patil, G. M. and Savanurmath, C. J. (1989). Oviposition behavior and egg hatchability in tasar uzi fly *Blepharipasp.* (Walker). *J. Bomb. Nat. Hist.*, 86: 472-473.
9. Ramprakash and Kshirsagar, R. V. (2019). Eco-friendly management of uzifly (*Exorista bombycis*) for reduction to loss of cocoon productivity in Pune division of Maharashtra state. *International Journal of Innovation Scientific Research and Review*,1(1): 007-008.
10. Thangavelu, K. and Sahu, A. K (1986). Some studies on the bionomics of *Exorista sorbillans* (Wied.) from North Eastern India. *Sericologia*, 26(1): 77-82.
11. Negi, B. and Barah, A.; Siddiqui, A. A. and Sengupta, A. K. (1993). *Criculatrifenestrata* (Lepidoptera: Saturniidae)- a new alternate host of *Blepharipa* sp. (Diptera: Tachinidae). *Recent advances in uzi fly research: Proceedings of the National Seminar on Uzi Fly and Its Control*, 269-271.

Commented [A12]: space

Commented [A13]: space

Table 1 Measurement of different life stages of *Blepharipa sp.* (Walker)

Sl. No.	Life stages	Length(mm)	Range	Width(mm)	Range
		Mean (±) SEM		Mean (±) SEM	
1	Egg	0.80±0.09	0.7-0.9	0.38±0.07	0.3-0.5
2	Maggot				
	First instar	1.18±0.19	1-1.5	0.46±0.08	0.4-0.6
	Second instar	6.18±0.68	5-7	2.42±0.38	2-3
	Third instar	17.52±0.34	17-18	5.48±0.34	5-6
3	Pupa	11.32±0.68	10-12	5.34±0.68	4-6
4	Adult				
	Male	13.12±0.68	12-14	5.74±0.38	5-6
	Female	11.24±0.67	10-12	5.72±0.37	5-6
5	Wing span				
	Male	9.48±0.40	9-10	3.70±0.36	3-4
	Female	8.50±0.35	8-9	3.34±0.42	3-4

Table 2 Developmental parameters of uzi fly, *Blepharipa sp.* Walker in muga silkworm

Sl. No.	Parameter	Mean (\pm) SEM	Range
1	Incubation period (days)	2.40 \pm 0.48	2-3
2	Hatching percentage (%)	84.48 \pm 2.08	81.25-87.50
3	Larval period (days)	6.20 \pm 0.73	5-7
4	Pupal period (days)	12.40 \pm 0.78	11-13
5	Adult emergence (%)	74.00 \pm 2.45	70.00-76.67
6	Sex ratio (female:male)	2.84:1	-
7	Adult longevity (days)		
	Male	10.80 \pm 0.96	9-12
	Female	12.00 \pm 0.88	11-13

Table 3 Reproductive parameters of uzi fly, *Blepharipa sp.* infesting muga silkworm

Sl. No.	Parameter	Mean (\pm) SEM	Range
1	Pre-oviposition period (hours)	7.04 \pm 0.652	6-8
2	Oviposition period (days)	6.40 \pm 0.784	5-7
3	Rate of oviposition (no. of eggs/female/day)	35.20 \pm 4.487	27-41
4	Fecundity	219.00 \pm 17.188	186-237
5	Post-oviposition period (days)	4.40 \pm 0.48	4-5

Table 4 Site of oviposition of *Blepharipa* sp. on host body (*Antheraea assamensis* Helfer)

Site of oviposition on The host body	Mean (\pm) SEM	Percent infestation (%)
Dorsal	11.60 \pm 0.78	23.97
Lateral	24.60 \pm 0.99	50.83
Ventral	12.20 \pm 0.73	25.42

UNDER PEER REVIEW



Fig. 1 Egg of Muga uzifly



a



b



c

Fig. 2 Site of oviposition of Muga uzifly a. dorsal, b. lateral, c. ventral



Fig. 3a 1st instar maggot of Muga uzifly



Fig. 3b Fat mass of host larval body enclosing 1st instar maggot of Muga uzifly



Fig. 4 2nd instar maggot of Muga uzifly



Fig. 5 3rd instar maggot of Muga uzifly



Fig. 6a Freshly formed Muga uzifly pupa

Fig. 6b Muga uzifly pupa after they darkens in colour

Commented [A14]: dark in colour



Fig. 7 Adult Muga uzifly




Fig. 8 Respiratory funnel formed by 1st instar maggot of Muga uzifly

Commented [A15]: include in result



Fig. 9 Black scar formed after maggot penetration to larval body of muga

UNDER PEER REVIEW






Fig. 10 Defective cocoon due to uzi infestation

UNDER PEER REVIEW

