

Original Research Article

Biology of Muga Uzi Fly: The Menace threatening the Muga Industry

[However, if you're looking for an alternative that might capture attention more dynamically, here are a couple of suggestions:](#)

1. [“Unveiling the Uzi Fly: Understanding Its Impact on the Muga Silkworm Industry”](#)
2. [“The Uzi Fly Challenge: Biological Insights and Threats to Muga Silkworms”](#)

ABSTRACT

The experiment was carried out with the purpose to thoroughly study the biology of a very serious endo-larval parasitoid of muga silkworm, *Blepharipa sp.* (Walker). The study was done in experiment field where the muga silkworms were reared under net condition and were allowed to infest by the uzi fly, whereas the study of biology was carried out in laboratory condition at room temperature and humidity. The study regarding biology revealed that the life cycle of *Blepharipasp.* (Walker) is completed in 30 days on average for male and 33 days for female fly which consist an average of 2.40 days of incubation period, 6.20 days larval period, 12.40 days pupal period and adult longevity was 10.80 days in case of male and 12.00 days in case of female. Other aspects of their life cycle regarding peak infestation season, site of oviposition were also studied. The mortality due to uzi fly infestation was 100% as the worms died either before spinning stage or died inside their cocoons and regarding the cocoons they produced, almost all were defective, not suitable for reeling. Since this is the most destructive among all insect pests of muga silkworm causing not only economic losses but also destroying farmers motivation, a thorough study of the biology was of utmost importance.

Keywords: *Antheraea assamensis*, *Blepharipa sp.*, Major pest, Pest biology

[Revised abstract:](#)

ABSTRACT

[This study aims to thoroughly investigate the biology of a significant endo-larval parasitoid of muga silkworms, *Blepharipa sp.* \(Walker\). The research was conducted both in the field, where muga silkworms were reared under net conditions and allowed to be infested by the uzi fly, and in the laboratory at room temperature and humidity. The findings revealed that the life cycle of *Blepharipa sp.* is completed in an average of 30 days for males and 33 days for females, with an incubation period of 2.40 days, a larval period of 6.20 days, a pupal period of 12.40 days, and adult longevity of 10.80 days for males and 12.00 days for females. Additional aspects such as peak infestation season and oviposition sites were also studied. The mortality rate due to uzi fly infestation was 100%, with worms dying either before the spinning stage or inside their cocoons, resulting in defective cocoons unsuitable for reeling. Given the severe economic losses and](#)

demotivation caused to farmers by this pest, a thorough understanding of its biology is of utmost importance.

UNDER PEER REVIEW

1. Introduction

India's diverse natural environment, conducive to the production of all four types of commercial silk, bestows upon it a distinctive presence in sericulture. Of them is the muga silkworm which, being multivoltine and endemic to Assam and some other north-eastern states, are typically reared outdoors, with cocoon spinning to seed production occurring indoors, rendering them vulnerable to pest infestations. Among these pests, the uzi fly poses a significant threat, leading to yield reduction, particularly during the peak infestation period i.e. the chotua crop, a pivotal commercial seed crop within the muga cultivation. They cause as loss of silkworm crops which hampers the generation of future seed crops and hence disrupts the commercial seed crop production which ultimately causes loss in silk production (Goswami *et al.*, 2013). The uzi fly is a member of the family Tachinidae, most of which are exclusive parasitoids, primarily targeting Lepidoptera but also known to attack different stages of other insects such as Heteroptera, Coleoptera, and Symphyta of Hymenoptera (Dai *et al.*, 2022). The muga silkworm are known to be parasitized by two Tachinid species viz. *Exorista sorbillans* and *Blepharipa sp.* Among them, *B. sp.* is identified as the primary Tachinid fly that preys on muga silkworms where the peak period of attack occurs between December and April, with reported losses ranging from 48.7% to 80% in various seed areas across Assam (Choudhury *et al.*, 2014). The first report of *Blepharipasp.* as a serious pest of muga silkworm was reported in the year 1989 by Goswami and Barah. The infestation of the uzi fly is recognizable by the black scar where the attack occurred, and maggots emerging from the cocoon are evident through holes in it making them inappropriate for reeling. Muga silkworms, having six crops annually, are raised year-round. Due to their close synchronization with the uzi fly life cycle, it's challenging to rear them without infestation. Additionally, outdoor rearing increases the risk of crop loss due to other pests and diseases, compounded by different climatic conditions throughout the year. Managing the infestation involves employing nylon nets to cover the host plants, which is difficult because farmers typically don't prune their fields, resulting in large trees that cannot be completely covered by nets. The current study aimed to investigate the biology and damage symptoms by of the pest as well as to promote simple yet effective cultural and hygienic practices for its management.

2. Materials and methods

The study of the uzi fly life cycle was conducted in field conditions at the experimental field of Department the Department of Sericulture, AAU, Jorhat, Assam (Lat 26.721544°, Long 94.196678°) with temperature, humidity and rainfall levels ranging from 13.48°C-26.73°C, 77%-78.5% and 0mm-86mm. To calculate the oviposition and fecundity of the uzi fly, muga silkworm larvae were infested artificially by keeping them with mated female uzi fly in the field and were properly covered with a nylon net to prevent the uzi fly from flying out. ~~Later~~ The later part of the study regarding pupal period and adult emergence ~~were was~~ done in laboratory conditions at room temperature and humidity by keeping the cocoons of infested muga silkworm each separately in different plastic containers and was made sure to carry out everyday observation. The measurements of the dimensions of the egg, maggot, pupa and adult were carried out with the help of a binocular stereoscope.

3. Result

The female fly after copulation searches for the suitable host for oviposition which apparently found to be mostly the late instars of the muga silkworm i.e. 4th and 5th instars. They attach the eggs to the exposed integument of silkworm larvae mostly in the intersegmental region, although they were also found to lay eggs on exposed sides of prolegs, near spiracles, etc. With an average oviposition rate of 35.20 numbers per day (table 3) during the experimental period, the female fly was spotted to lay 7-16 eggs per silkworm. The eggs were found to have hatched through chorion on the antero-ventral surface, with an observed hatching percentage of 84.48%, following an incubation period of 2-3 days (table 2). The measurements of the egg, maggot, pupa, and adult of *B. sp.* are depicted in table Table 1.

3.1. Egg stage of *Blepharipasp.*

It was found that the eggs were ovoid-shaped and of macro type (fig.1). The surface of the egg was relatively flat ventrally with an opaque chorion and convex dorsally. Although the gravid adult female uzi flies appeared to lay their eggs anywhere on the larval body of the muga silkworm, it became apparent that the lateral region was the most desired location, subsequently followed by both ventral and dorsal site with a percent infestation of 50.83%, 25.42% and 23.97% respectively (fig. 2-a, b, c) (table 4).

3.2. Larval stage of *Blepharipa sp.*

With a nearly spherical posterior region and a tapered anterior region, the apodous and fusiform morphology of the maggots were typified by their curved hooked mouth part displaying a total of 12 body segmentations. The third instar matured maggots emerged from the host body to undergo pupation meanwhile completing their all three larval instars inside the host body.

Formatted: Font: Italic

First instar maggot: The initial instar maggots (fig. 3a) were minuscule, off white in appearance, and possessed two tiny, dot-like spiracles at posterior body region. They were linked to the respiratory funnel (fig. 8) that had developed at the puncture site and did not move much, staying close to it and they do it by gathering a mass of fat bodies near the entrance point inside the infested silkworm larvae (fig. 3b)

Second instar maggot: The maggots in their second instar (fig. 4) had a gentle white appearance with two light brown spiracles at the posterior end. Additionally, they also exhibited sedentary behavior and remained close to the puncture site in the host larval body.

Third instar maggot: The maggots in their third instar were dirty white to yellowish white in colour (fig. 5), having two noticeable spiracles with sclerotized peritreme. The third instar uzi maggots go throughout the entire host body, eating voraciously on the fat content of the host larva. They exited the host larval body by piercing the dead host's body wall at maturation.

3.3. Pupal stage of *Blepharipasp.*

After emerging from the silkworm larval body, the mature uzi maggots searched for a suitable location to pupate. The maggots preferred to pupate in soil although they typically hunt for fissures and crevices in their surroundings. The freshly produced pupae had a light brown colour (fig 6a) that eventually grew darker with time (fig. 6b). These coarctate pupae comprised of 11 body segments. After the pupal stage, which lasted an average of 12.40 days (table 2), the adult fly emerged from the pupa by breaking the predetermined line of weakness at the anterior region of the pupa. The adults appeared to emerge most often in the morning, when the temperature appeared to be at its best. Temperature and rainfall had a significant impact on adult emergence; greater emergence was detected following rainy days and a greater infestation percentage was found in the Chotua crop of muga silkworm (Feb-Mar). With an average emergence percentage of 74.00, the emergence persisted for three to four days (table Table 2).

3.4. Adult stage of *Blepharipasp.*

The adult uzi flies had four longitudinal lines of bristles on the thoracic notum and long lateral bristles on the lateral abdominal area. They were blackish in appearance with an orange tinge at the dorso-lateral region of the abdomen. It was evident that the female was a bit bigger ~~over than~~ the male. Mating mostly took place on the day of emergence (fig. 7-a,b).

4. Discussion

Blepharipa sp. is without a doubt the most destructive pest resulting in significant losses in the muga industry among all the insect pests of muga silkworm in Assam and Meghalaya states of India (Kabiraj *et al.*, 2022). Despite being the most destructive, there is little information accessible regarding the biology of the muga uzi fly. Reddy and Ranjan (2011) reported higher infestation in the Chotua crop (43.0%) and the Jarua crop (19.0%) of muga silkworm in upper Assam. According to Manjunatha and Puttaraju (1993), the eggs of the uzi fly were also of macrotype, dull white in colour, almost oval in shape with a slightly pointed proximal region. Patil and Savanurmah (1989) reported that usually the eggs of *B. sp.* hatched 3 days after oviposition and the hatching percentage was as high as 97.9% when observed under lab conditions. They also revealed that the *B. sp.* preferred late instar worms of *Antheraea mylitta* D., mainly the 4th instar worms over the early instar worms. *B. sp.* were ~~was~~ reported to infest hosts on the basis of based on the host-stage, vigour, and health of the silkworm. Ramprakash and Kshirsagar (2019) found that the maggot period of *Exorista bombycis* was found to be 5-7 days. The size of uzi flies attacking muga silkworms and mulberry silkworms also varies. Maggots of uzi flies targeting mulberry silkworms were smaller compared to those attacking muga silkworms and as reported these variations may stem from the larger body size of muga silkworm (Thangavelu and Sahu, 1986). Thangavelu and Sahu (1986) also stated that the pupa of the uzi fly were photonegative and undergo pupation mostly in soil but sometimes the pupation may occur inside the host larval cocoon. Goswami *et al.* (2013) reported that the maggot of uzi fly infesting muga silkworm after maturation and coming out of the silkworm body pupates within 7-8 hours and the pupal period was found to be 10 days. Negi *et al.* (1993) stated that the total life span of *B. sp.* was 48.1-64.1 days in an alternate host, *Cricula trifenestrata*.

5. Conclusion

Various reports regarding the infestation of a very destructive Tachinid pest of muga silkworm were recorded in Assam causing devastating economic loss. This study was done to provide a thorough study and familiarize the biology of the pest i.e. uzi fly. The uzi flies were found to mostly attack the muga silkworm during the Chotua crop (Feb-Mar) although the infestation occurs throughout the year. Starting from the egg stage to adult, the life cycle of the male uzi fly is 30 days long while the female has a longer life span of 33 days on average. The adult female finds the lateral region of the muga larva to be most favorable for egg laying which hatched within an incubation period of

2-3 days (table 2). The hatching of the egg is characterized by a black scar forming near the egg which occurred due to the puncture made by the maggot while entering the host larval body (fig. 9). The shorter egg incubation & larval period which definitely is the destructive period makes it very difficult to control which is aided by the fact that they are endo-larval parasitoid, hence once the host is infested, they will surely die. Even if the host larvae managed to survive till cocoon spinning, those cocoons will be either flimsy, small, deformed or not fully formed (fig. 10). The resulting losses sometimes include the whole crop loss which renders the production of next generation progeny.

References

1. Goswami, N. K.; Nath, P. and Saharia, D (2013). Uzi Fly Infestation Severity in Muga Seed Cocoons, *Antheraea assamensis* Helfer and Crop Loss During Chotua Crop in Assam. *Indian Journal of Applied Research*, **3**(10): 1-4.
2. Kabiraj, D.; Chetia, H.; Nath, A.; Sharma, P.; Mosahari, P.; Singh, D.; Dutta, P.; Neog, K. and Bora, U. (2022). Mitogenome-wise codon usage pattern from comparative analysis of the first mitogenome of *Blepharipa* sp. (Muga uzifly) with other Oestroid flies. *Scientific Reports*. **12**(1): 7028.
3. Dai, M.; Yang, J.; Liu, X.; Gu, H.; Li, F.; Li, B. and Wei, J. (2022). Parasitism by the tachinid parasitoid *Exorista japonica* leads to suppression of basal metabolism and activation of immune response in the host *Bombyx mori*. *Insects*, **13**: 792.
4. Choudhury, B. and Kumar, R.; Chutia, P. and Rajkhowa, G. (2014). Host infestation Potentiality of *N. thymus* to Control the Uzifly of muga silkworm, *Antheraea assamensis* (Helfer) - A Bio-control Tool for Controlling Uzifly. *Biological Forum – An International Journal*, **6**(1): 1-4.
5. Goswami, M. C. and Barah, A. (1989). Report of *Blepharipa* sp. (Walker) (Tachinidae) as a serious pest of muga silkworm, *Antheraea assama* Westwood (Saturniidae). *Current Science*, **58** (5): 267-268.
6. Reddy, S. G. E. and Ranjan, R. K. (2011). present status of uzi fly, *Exorista bombycis* (Louis) (Diptera: Tachinidae) incidence on muga silkworm, *Antheraea assamensis* Helfer (Lepidoptera: Saturniidae) in upper Assam. *Munis Entomology & Zoology*, **6**(2): 856-858.
7. Manjunatha, H. and Puttaraju, H. P. (1993). The Egg of Uzi Fly, *Exorista sorbillans* (? *E. bombycis* Louis) (Diptera: Tachinidae). *Appl. Entomol. Zool.*, **28**(4): 574-577.
8. Patil, G. M. and Savanurmath, C. J. (1989). Oviposition behavior and egg hatchability in tasar uzi fly *Blepharipasp.* (Walker). *J. Bomb. Nat. Hist.*, **86**: 472-473.
9. Ramprakash and Kshirsagar, R. V. (2019). Eco-friendly management of uzifly (*Exorista bombycis*) for reduction to loss of cocoon productivity in Pune division of Maharashtra state. *International Journal of Innovation Scientific Research and Review*, **1**(1): 007-008.
10. Thangavelu, K. and Sahu, A. K (1986). Some studies on the bionomics of *Exorista sorbillans* (Wied.) from North Eastern India. *Sericologia*, **26**(1): 77-82.
11. Negi, B. K.; Barah, A.; Siddiqui, A. A. and Sengupta, A. K. (1993). *Criculatrifenestrata* (Lepidoptera: Saturniidae)- a new alternate host of *Blepharipa* sp. (Diptera: Tachinidae). *Recent advances in uzi fly research: Proceedings of the National Seminar on Uzi Fly and Its Control*, 269-271.

Table 1 Measurement of different life stages of *Blepharipasp.* (Walker)

Sl. No.	Life stages	Length(mm)	Range	Width(mm)	Range
		Mean (\pm) SEM		Mean (\pm) SEM	
1	Egg	0.80 \pm 0.09	0.7-0.9	0.38 \pm 0.07	0.3-0.5
2	Maggot				
	First instar	1.18 \pm 0.19	1-1.5	0.46 \pm 0.08	0.4-0.6
	Second instar	6.18 \pm 0.68	5-7	2.42 \pm 0.38	2-3
	Third instar	17.52 \pm 0.34	17-18	5.48 \pm 0.34	5-6
3	Pupa	11.32 \pm 0.68	10-12	5.34 \pm 0.68	4-6
4	Adult				
	Male	13.12 \pm 0.68	12-14	5.74 \pm 0.38	5-6
	Female	11.24 \pm 0.67	10-12	5.72 \pm 0.37	5-6
5	Wing span				
	Male	9.48 \pm 0.40	9-10	3.70 \pm 0.36	3-4
	Female	8.50 \pm 0.35	8-9	3.34 \pm 0.42	3-4

Table 2 Developmental parameters of uzi fly, *Blepharipasp.* Walker in muga silkworm

Sl. No.	Parameter	Mean (\pm) SEM	Range
1	Incubation period (days)	2.40 \pm 0.48	2-3
2	Hatching percentage (%)	84.48 \pm 2.08	81.25-87.50
3	Larval period (days)	6.20 \pm 0.73	5-7
4	Pupal period (days)	12.40 \pm 0.78	11-13
5	Adult emergence (%)	74.00 \pm 2.45	70.00-76.67
6	Sex ratio (female:male)	2.84:1	-
7	Adult longevity (days)		
	Male	10.80 \pm 0.96	9-12
	Female	12.00 \pm 0.88	11-13

Table 3 Reproductive parameters of uzi fly, *Blepharipasp.* infesting muga silkworm

Sl. No.	Parameter	Mean (\pm) SEM	Range
1	Pre-oviposition period (hours)	7.04 \pm 0.652	6-8
2	Oviposition period (days)	6.40 \pm 0.784	5-7
3	Rate of oviposition (no. of eggs/female/day)	35.20 \pm 4.487	27-41
4	Fecundity	219.00 \pm 17.188	186-237
5	Post-oviposition period (days)	4.40 \pm 0.48	4-5

Table 4 Site of oviposition of *Blepharipasp.* on host body (*Antheraea assamensis* Helfer)

Site of oviposition on The host body	Mean (\pm) SEM	Percent infestation (%)
Dorsal	11.60 \pm 0.78	23.97
Lateral	24.60 \pm 0.99	50.83
Ventral	12.20 \pm 0.73	25.42

UNDER PEER REVIEW



Fig. 1 Egg of *Muga uzifly*




a



b

REVIEW





c

Fig. 2 Site of oviposition of *Muga uziflya*. dorsal, b. lateral, c. ventral



Fig. 3a 1st instar maggot of *Muga uzifly*



Fig. 3b Fat mass of host larval body enclosing 1st instar maggot of *Muga uzifly*



Fig. 4 2nd instar maggot of Muga uzifly



Fig. 5 3rd instar maggot of Muga uzifly



Fig. 6a Freshly formed Muga uzifly pupa

Fig. 6b Muga uzifly pupa after they darkens in colour



Fig. 7 Adult Muga uzifly



Fig. 8 Respiratory funnel formed by 1st instar maggot of Muga uzifly



Fig. 9 Black scar formed after maggot penetration to larval body of muga

UNDER PEER REVIEW



Fig. 10 Defective cocoon due to uzi infestation

UNDER PEER REVIEW