

Prevalence and Factors Associated with Non-adherence to Iron and Folic Acid Supplementation among Women for Antenatal Care at Mwembeladu Hospital, Zanzibar

ABSTRACT

BACKGROUND: Daily iron and folic acid supplementation (IFAS) during pregnancy reduces the risk of all types of maternal anaemia by 70% and iron deficiency anaemia by 57% at term; despite the WHO recommendations, pregnant women are still vulnerable because the use of Iron and Folic Acid Supplementation is still low in many countries including Tanzania. Therefore, the study aims to comprehensively understand the prevalence and factors associated with non-adherence to Iron and Folic Acid Supplementation among pregnant women attending Antenatal Care at Mwembeladu Hospital. Achieving this objective will contribute to developing effective strategies to improve IFAS adherence.

METHODOLOGY: The study was cross-sectional and ANC-based, using questionnaires. Questionnaires were distributed to the 260 pregnant women attending ANC at Mwembeladu Hospital. Data were analyzed using SPSS computer software version 22.

RESULTS: Our study showed that 52.7% had more than 4 ANC visits. Based on self-reported adherence, about (61.9%) of pregnant women were taking IFAS supplementation, and (38.1%) were not taking IFAS completely. Among those taking IFAS, (60.2%) out of 161 pregnant women were taking four tablets per week as recommended by WHO, and (39.8%) out of 161 were not following the WHO recommendations. Therefore, out of 260, zero adherence was 38.1%, poor adherence was 24.6%, and good adherence was 37.3%. Hence, 62.7% were non-adherent to IFAS supplementation as recommended by WHO. Also, the result showed that the major

factors militating against pregnant women taking IFAS are lack of knowledge of IFAS (22.1%), side effects of IFAS (19%), forgetfulness (18.4%), and lack of understanding of anaemia (12.3%).

CONCLUSION: Overall, the adherence to IFAS among pregnant women was low and did not meet the WHO recommendations for preventing and treating anaemia during pregnancy. Socio-demographic factors, including occupation and education level, maternal characteristics, parity, and gestation age, are all significantly associated with adherence to IFAS. Factors associated with poor adherence to IFAS include side effects, failure to access IFAS, forgetfulness, and knowledge about anaemia and IFAS. Healthcare facilities and providers should strengthen the system to create community awareness of IFAS, its benefits, and the side effects of IFAS, as well as knowledge of how they contribute to increasing the rate of anaemia in pregnant women.

Keywords: adherence, iron with folic acid supplementation, Zanzibar

Introduction

Anaemia is a global public health problem affecting both developed and developing countries. It significantly impacts human health, as well as social and economic health. It is associated with increased morbidity and mortality in women and children, poor birth outcomes such as preterm delivery, low birth weight, and perinatal mortality, and decreased productivity of workforces in adults [1]. In 2019, the global prevalence of anaemia was 29.9% in women of reproductive age, equivalent to over half a billion women aged 15 – 49 years. Update global anaemia, 2021; the Prevalence of anaemia was 29.6% in non-pregnant women of reproductive age and 36.5% in pregnant women; hence, pregnant women are at high risk of anaemia [2]. Also, according to the data provided by WHO, the prevalence in Africa is estimated to be 37.8% and 46.3%, respectively. In Tanzania, the prevalence of anaemia in reproductive age between 15 – 49 years is 57%. Also, 50% of all anaemia is estimated to be due to iron deficiency. Hence, anaemia in pregnant remains persistently high. In Zanzibar, 2018, the reported overall prevalence of anaemia was 80.8% [3,4]

Anaemia refers to a haemoglobin (Hb) level of less than 11g/dl for mild anaemia, between 7-9.9g/dl for moderate anaemia, and less than 7g/dl considered severe anaemia. It affects all age groups, but pregnant women are more vulnerable. [5]

The most common cause of anaemia in pregnancy is iron and folic acid deficiency due to physiological changes during pregnancy, foetal growth, and development that increase the need for iron and folic acid. The increased demand for these nutrients cannot be met by diet alone as the availability of nutrients in pregnant women declines. Three factors that may be associated with anaemia among pregnant women are reduced intake of red meat and green vegetables and low MUAC [6]. Folic acid deficiency can occur if food is not replaced with iron and folic acid tablets during pregnancy.

According to WHO, daily oral iron and folic acid supplementation are recommended for antenatal care to reduce the risk of low birth weight, maternal anaemia, and iron deficiency. The suggested schemes for daily iron and folic acid supplementation in pregnant women are [7] :

Table 1. Suggested schemes for daily iron and folic acid supplementation in pregnant women

Supplement composition	Iron: 30-60mg of iron. Folic acid: 0.4mg
Frequency	One supplement daily
Duration	Throughout pregnancy and 3 months post-delivery.
Target group	All pregnant adolescent and adult women
Settings	All setting

Despite this recommendation, some pregnant women are non-compliant with iron and folic acid supplements. Several factors contribute to non-adherence to IFAS, and different studies highlight them. A study concluded that factors contributing to low compliance with Iron and Folic Acid supplements were no formal education, Poor knowledge about

anaemia, Poor families (in economic status), forgetfulness, attending health information about IFAS and Developing, and other problems during the current pregnancy [8]

Women who failed to consume the IFAS tablet, according to the recommendation, would experience iron and folic acid deficiency and associated adverse effects on mothers and newborns. [8]. Anaemia is reported to have adverse maternal and child health effects and increase the risk of maternal and perinatal mortality. The adverse health effects for the mother include fatigue, poor work capacity, impaired immune function, increased risk of cardiac diseases, and mortality. Anaemia in pregnancy is associated with an increased risk of preterm birth and LBW babies postpartum haemorrhage and poor birth outcomes. Preterm and LBW are still the leading causes of neonatal death in developing countries like Tanzania, contributing to 30% of deaths. [9]

Broad objective

OBJECTIVE: Our study aimed to comprehensively understand the prevalence and factors associated with non-adherence to Iron and Folic Acid Supplementation among pregnant women attending Antenatal Care at Mwembeladu Hospital. By achieving this objective, we hope to contribute to developing effective strategies to improve IFAS adherence.

Specific objectives

1. To determine the social-demographic characteristics associated with non-adherence to IFAS among pregnant women attending Antenatal Care.
2. To identify the factors associated with non-adherence to Iron and Folic Acid Supplementation among pregnant women attending Antenatal Care.

Methodology

Description of the study area

Mwembeladu Maternity Hospital is about a 10-minute drive from Mnazi Mmoja Referral Hospital (MMH) in Zanzibar. The hospital has 36 beds and conducts an average of 550 monthly births. Only low-risk expecting mothers deliver at the hospital. High-risk and complicated cases are referred to MMH.

Target population

Pregnant women at Mwembeladu Hospital from January 2023 to May 2023.

Study population

The study population included pregnant women attending Antenatal Care at Mwembeladu Hospital.

Study design

A descriptive cross-sectional design was used in this study to determine the proportion of non-adherence to IFAS among pregnant women at ANC.

Sampling technique.

Simple random sampling (SRS) was used to get respondents for the study. In this method, each member of the study population had an equal chance of being chosen as part of the sample. Simple random sampling was used to select pregnant women attending ANC.

Sample size determination

The sample size was determined by using the formula shown below

The sample size was as follows;

$$N = Z^2 \times P \times q / E^2$$

Where= is the minimum desired sample size

Z = the standard average deviation of 1.96, corresponding to a 95% confidence interval.

P = Proportion of the target population estimated from previous studies is 20.3%, as shown by Winfrida B. Lyoba et al. in Tanzania, 2020.

E= expected margin of error is 5%

$$q = (1-P)$$

$$N = Z^2 p (1-P) / e^2$$

$$N = (1.96)^2 \times 0.203 \times (1-0.203) / (0.05)^2$$

$$N = 249$$

Therefore, the minimum sample size was 249.

For convenience and to accommodate the non-response, there were 260 samples.

Inclusion and exclusion criteria.

All Pregnant women attending a prenatal clinic at MLH were included. The Non-pregnant women, pregnant women who had a mental illness, and pregnant women who were severely ill and could not respond well were excluded.

Study variable

Dependent variables:

- Number of pregnant women taking IFAS

Independent variables:

- **Social demographic factors:** Age, Educational status, marital status, and occupation status.
- **Obstetric and Health-related factors:** gravidity, number of children, time initiating ANC, knowledge of anemia, and IFAS.
- **Risk factors for non-adherence such as** availability of IFAS, side effects and socio-cultural feeding practice of pregnant women, hospital-related factors, and family history of anaemia.

Definitions

Good adherence to IFAS: pregnant women who were taking more than four tablets per week

Poor adherence: pregnant women who were taking less than four tablets per week

Data collection technique

Primary and secondary data collection techniques and methods were used.

Primary data collection techniques/methods.

This study collected primary data through face-to-face interviews using an open questionnaire, group discussion with the target group, and personal observation.

Secondary data collection techniques/methods.

Secondary data was collected from up-to-date documentary sources, i.e., published and unpublished books and journal articles.

Ethical approval and participant consent:

The Zanzibar Health Research Institute, ZAHREC, approved the ethical clearance for conducting health research with **REF NO. ZAHREC/05/MARCH/2023/39**. The collected data were used for this research purpose only and kept completely confidential. Verbal informed consent was obtained from the study participants, and personal identifiers were excluded during the data collection to ensure confidentiality.

RESULTS

socio-demographic characteristics

The study consisted of 260 pregnant women attending ANC at Mwembeladu Hospital Zanzibar. Regarding age, About 136 (52.3%) were between the age of 23-32 years, followed by 94 (36.2%) between the age of 33-42 years, 15 (5.8%) were between the age of 13-22 years, 15 (5.8%) were between the age of greater than 42 years. About 150 (57.7%) had secondary education, followed by 54 (20.8%) who attained primary education, then 44 (16.9%) had a university education level and 12(4.6%) never go to school. Relating to marital status, about 212 (81.5%) were married, 23 (8.8%) were single, 18 (6.9%) were widowed, and 7 (2.7%) were divorced. In this group of pregnant women, 139 (53.5%) were homemakers, followed by 88 (33.8%) who were daily laboured, and the remaining 33 (12.7%) were government employees. (Table 2)

Adherence of IFAS on pregnant women and associated factors

Among the pregnant women, 161(61.9%) were taking IFAS, although 99 (38.1%) were not taking the supplementation (graph 1). Of the 161 gravitas who had taken IFAS, 97 (60.2%) followed the recommendation of more than four tablets per week (good adherence), and 64 (39.8%) did not follow the recommendation of less than four tablets per week (poor adherence), (table 3)

Out of 260 participants, 163 had no adherence to IFAS (99 who had not taken IFAS plus 64 who had taken less than four tablets). The factors that contributed to them not taking IFAS were; 3 (1.8%) failed access IFAS, 30 (18.4%) forgetfulness, 11 (6.7%) late initiated ANC, 5 (3.1%) long distance to health facilities, 31 (19%) side effects of IFAS, 8

(4.9%) tiredness, 20 (12.3%) lack of knowledge of anaemia, 36 (22.1%) lack of knowledge of IFAS, 10 (6.1 %) lack of support from partner, 9 (5.5%) others.

Factors associated with poor adherence to treatment

Within the sociodemographic factors, there was no relationship between marital status, level of education, or occupation with treatment adherence. Pregnant women with a large family had 1.3 times more adherence to IFAS than those with a small family OR 1.349 CI 0.630- 0.91 $p < 0.05$. Gestational age and time of attendance at the prenatal clinic were related to treatment adherence.

Prior knowledge about anaemia or having previously suffered from anaemia was unrelated to adherence. Not hearing about IFAS correlated with poor treatment adherence: Pearson Chi-Square 17.45 $p < 0.00$. Likewise, not knowing that IFAS is beneficial was associated with poor adherence: Pearson Chi-Square 22.41 $p < 0.000$.

Table 2: Socio-demographic characteristics data

Social demographic data	Frequency	Percentages (%)
Ages(years)		
13-22	15	5.8
23-32	136	52.3
33-42	94	36.2
Greater than 42	15	5.8
TOTAL	260	100.0
EDUCATIONAL STATUS		
Primary	54	20.8
Secondary	150	57.7

University	44	16.9
Never go to school	12	4.6
TOTAL	260	100
MARITAL STATUS		
Single	23	8.8
Married	212	81.5
Widowed	18	6.9
Divorced	7	2.7
TOTAL	260	100
OCCUPATIONAL STATUS		
Housewife	139	53.3
Daily laboured	88	33.8
Government employee	33	12.7
TOTAL	260	100

Table 3: Prevalence of adherence of IFAS on pregnant Women.

Number of women who are taking IFAS	Frequency	Percentage (%)
Yes	161	61.9
No	99	38.1
TOTAL	260	100
Adhere to IFAS as recommended among those who taking IFAS (n=161).		
Less than 4 tabs per week	64	39.8
More than 4 tabs per week	97	60.2

TOTAL	161	100.0
-------	-----	-------

Table 4. Factors that are associated with non-adherence of IFAS as recommended on pregnant women. (n=163)

	frequency	Percentage (%)	Valid percent	Cumulative percent
failed access IFAS	3	1.8	1.8	1.8
Forgetfulness	30	18.4	18.4	20.2
Late initiated ANC	11	6.7	6.7	26.9
Long distance to health facilities	5	3.1	3.1	30.0
Side effects of IFAS	31	19.0	19.0	49.0
Tiredness	8	4.9	4.9	53.9
Lack of knowledge of anaemia	20	12.3	12.3	66.2
Lack of knowledge of IFAS	36	22.1	22.1	88.3
Lack of support from partner	10	6.1	6.1	94.5
Others	9	5.5	5.5	100
TOTAL		100.0	100.0	

Table 5. Factors associated to adherence to IFAS among pregnant women at Mwembeladu Maternity Hospital, Zanzibar

(n =260) Factors	Good adherence to IFAS (yes)	Poor adherence to IFAS (no)	OR	
			OR	95% CI
Marital status				

Married	144 (55.4%)	91 (35%)	0.989	9.13-1.07
Single	16 (6.2%)	9 (3.4%)	1.11	0.51-2.41
Education of pregnant woman				
Educated	154 (59.2%)	94 (36.2%)	1.024	0.966-1.085
Never go to school	6 (2.30%)	6 (2.30%)	0.625	207-1.884
Occupation				
House wife	78 (30%)	61 (23.5%)	0.779	0.639-999
Worker	82 (31.5%)	39 (15%)	1.314	0.95-1.753
Family size				
Small	78 (30%)	62 (23.85 %)	0.786	0.350-.970
Médium and large	82 (31.5%)	38 (14.61%)	1.349	.630-0.91
History of anemia				
Yes	83 (31.92%)	34 (13.07%)	1.526	1.11-2.08
No	77 (29.61%)	66 (25.38)	.729	.589-903
Heard about IFAS				
Yes	134 (51.53%)	62 (23.84%)	1.368	1.157-1.161
No	24 (9.23%)	38 (14.61%)	0.040	2.56-6.24*
Is IFAS beneficial				
Yes	143 (55%)	64 (24.62%)	1.396	1.194-1.633
No	17 (6.54%)	36 (13.84 %)	.0295	0.175-0.496**

***Pearson Chi-Square 17.45 p .000**

**** Pearson Chi-Square 22.41 p .000**

Discussion

The study consisted of 260 pregnant women attending ANC at Mwembeladu Hospital. Based on socio-demographical characteristics, women's educational level and occupational status were significantly associated with good adherence. This is in contrast with studies done in Ethiopia [10, 11], which showed that women's educational level was associated with poor adherence and the association of anemia with intestinal parasites. The study recorded an interesting result on the area of adherence; although more than half of the participants reported taking IFAS, majority of the participants showed a no adherence to the WHO's recommendations by either

not taking at all or taking less than four times per week. The findings align with a study conducted by Kasulu Hospital in Tanzania [12], whereby only 20.3% of pregnant women adhered to IFAS as recommended. A study in Aykel town, Ethiopia [13], showed that only 47.6% adhered to IFAS as recommended. However, this figure contrasted with other studies [14,15], which show that about 56.0% adhered to IFAS. Another study was conducted in Assefa, Ethiopia, showing that 59.8% adhered to IFAS. The result also showed that the major factors militating against pregnant women taking IFAS are lack of knowledge of IFAS (22.1%), side effects of IFAS (19%), forgetfulness(18.4%), and lack of understanding of anaemia (12.3%). These results correspond to the study done at Kasulu Hospital, which showed that the major contributing factors to non-adherence were a lack of knowledge of anaemia and IFAS. This also collaborates with the study done at Yaounde Hospital in 2019, which showed that side effects (19.7%) and forgetfulness (70.1%) contributed to non-adherence to IFAS.

Conclusion

Overall, the adherence to IFAS among pregnant women was low and did not meet the WHO recommendations for the prevention and treatment of anaemia during pregnancy. Socio-demographic factors, including occupation and education level, maternal characteristics, parity, and gestation age, are all significantly associated with adherence to IFAS. Factors associated with poor adherence to IFAS include side effects, failure to access IFAS, forgetfulness, and knowledge about anaemia and IFAS

Recommendation

Healthcare facilities and providers should strengthen the system to create community awareness of IFAS, its benefits, and the side effects as well as knowledge of how they contribute to increasing the rate of anaemia in pregnant women. The RCH institution must ensure enough IFAS and that all pregnant

women receive IFAS regardless of the distance from their homes to health facilities. Bringing proper health education to television and radio jingles in the Kiswahili language can help awaken their enthusiasm for this project.

Family support, especially from partners to their pregnant wives, is critical in adhering to IFAS. This will help prevent forgetfulness and tiredness. Overall, family encouragement must be a task to arrest this dangerous trend in maternal health.

References

1. Kare AP, Gujo AB. Anemia among Pregnant Women Attending Ante Natal Care Clinic in Adare General Hospital, Southern Ethiopia: Prevalence and Associated Factors. *Health Serv Insights*. 2021 Jul 29; 14:11786329211036303. doi: 10.1177/11786329211036303. PMID: 34376992; PMCID: PMC8327009..
2. WHO Prevalence of anaemia in pregnant women (aged 15-49) (%) [Internet]. [cited 2024 Jul 24]. Available from: [https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-pregnant-women-\(-\)](https://www.who.int/data/gho/data/indicators/indicator-details/GHO/prevalence-of-anaemia-in-pregnant-women-(-))
3. Gibore NS, Ngowi AF, Munyogwa MJ, Ali MM. Dietary Habits Associated with Anemia in Pregnant Women Attending Antenatal Care Services. *Curr Dev Nutr*. 2020 Dec 11;5(1):nzaa178. doi: 10.1093/cdn/nzaa178. PMID: 33501404; PMCID: PMC7809361.
4. Ngimbudzi EB, Massawe SN, Sunguya BF. The Burden of Anemia in Pregnancy Among Women Attending the Antenatal Clinics in Mkuranga District, Tanzania. *Front*

Public Health. 2021 Dec 2;9:724562. doi: 10.3389/fpubh.2021.724562. PMID: 34926366; PMCID: PMC8674738

5. Solomon Y, Sema A, Menberu T. Adherence and associated factors to iron and folic acid supplementation among pregnant women attending antenatal care in public hospitals of Dire Dawa, Eastern Ethiopia. *Eur J Midwifery*. 2021 Aug 25;5:35. doi: 10.18332/ejm/138595. PMID: 34514359; PMCID: PMC8386124.
6. Osman MO, Nour TY, Bashir HM, Roble AK, Nur AM, Abdilahi AO. Risk Factors for Anemia Among Pregnant Women Attending the Antenatal Care Unit in Selected Jigjiga Public Health Facilities, Somali Region, East Ethiopia 2019: Unmatched Case-Control Study. *J Multidiscip Healthc*. 2020 Aug 10;13:769-777. doi: 10.2147/JMDH.S260398. PMID: 32848406; PMCID: PMC7428401.
7. WHO. Guideline: Daily iron and folic acid supplementation in pregnant women. Geneva, World Health Organization, 2012.
8. Tegodan E, Tura G, Kebede A. Adherence to Iron and Folic Acid Supplements and Associated Factors Among Pregnant Mothers Attending ANC at Gulele Sub-City Government Health Centers in Addis Ababa, Ethiopia. *Patient Prefer Adherence*. 2021 Jun 23;15:1397-1405. doi: 10.2147/PPA.S301630. PMID: 34188456; PMCID: PMC8236246.
9. Stephen G, Mgongo M, Hussein Hashim T, Katanga J, Stray-Pedersen B, Msuya SE. Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania. *Anemia*. 2018 May 2;2018:1846280. doi: 10.1155/2018/1846280. PMID: 29854446; PMCID: PMC5954959.
10. Girma S, Teshome T, Worku M, Solomon T, Kehulu S, Aman R, Bonsa M, Assefa T, Gezahegn H. Anemia and Associated Factors Among Pregnant Women Attending Antenatal Care at Madda Walabu University Goba Referral Hospital, Bale Zone, Southeast Ethiopia. *J Blood Med*. 2020 Dec 22;11:479-485. doi: 10.2147/JBM.S285190. PMID: 33376435; PMCID: PMC7765680.

11. Argaw D, Hussen Kabthymer R, Birhane M. Magnitude of Anemia and Its Associated Factors Among Pregnant Women Attending Antenatal Care in Southern Ethiopia: A Cross-Sectional Study. *J Blood Med.* 2020 Oct 9;11:335-344. doi: 10.2147/JBM.S264369. PMID: 33117019; PMCID: PMC7553252.
12. Lyoba WB, Mwakatoga JD, Festo C, Mrema J, Elisaria E. Adherence to Iron-Folic Acid Supplementation and Associated Factors among Pregnant Women in Kasulu Communities in North-Western Tanzania. *Int J Reprod Med.* 2020 Jun 4;2020:3127245. doi: 10.1155/2020/3127245. PMID: 32566646; PMCID: PMC7293754.
13. Assefa, H., Abebe, S.M. & Sisay, M. Magnitude and factors associated with adherence to Iron and folic acid supplementation among pregnant women in Aykel town, Northwest Ethiopia. *BMC Pregnancy Childbirth* **19**, 296 (2019). <https://doi.org/10.1186/s12884-019-2422-4>
14. Fouelifack FY, Sama JD, Sone CE. Assessment of adherence to iron supplementation among pregnant women in the Yaounde gynaeco-obstetric and paediatric hospital. *The Pan African Medical Journal* [Internet]. 2019 Dec 26 [cited 2024 Jul 25];34(211). Available from: <https://www.panafrican-med-journal.com/content/article/34/211/full>
15. Desta M, Kassie B, Chanie H, Mulugeta H, Yirga T, Temesgen H, Leshargie CT, Merkeb Y. Adherence of iron and folic acid supplementation and determinants among pregnant women in Ethiopia: a systematic review and meta-analysis. *Reprod Health.* 2019 Dec 21;16(1):182. doi: 10.1186/s12978-019-0848-9. PMID: 31864397; PMCID: PMC6925441.

UNDER PEER REVIEW