

Original Research Article

Scenario of Bacterial Leaf Blight of Rice Caused by *Xanthomonas oryzae* pv. *oryzae* in Major Rice Growing Areas of Andhra Pradesh

ABSTRACT

Rice is among the oldest crops of the world which feeds more than half of the world's population. Bacterial leaf blight caused by *Xanthomonas oryzae* pv. *oryzae*, is a most devastating disease which results in yield losses upto 70-90%.

Aim:The current research study was aimed to investigate the occurrence of disease and distribution of pathotypes.

Methodology:A roving survey was conducted over twelve major rice growing districts of Andhra Pradesh viz., East Godavari, West Godavari, Eluru, Kakinada, NTR district, Krishna, Nellore, Chittoor, Tirupati, Kadapa, Kurnool, Nandyal.

Place and Duration of Study: Department of Plant Pathology, S. V. Agricultural College, Tirupati, between November 2023 and February 2024.

Results:The highest disease incidence was found in Putrela village (72.34%) of Krishna district and lowest disease incidence was recorded in the fields of Burugupudi village (21.06%) of Kakinada District.

Conclusion:The results indicate that the disease is prevalent in all rice-growing areas, ranging with varying levels of disease incidence of 20 to 72 per cent.

Key words:Bacterial leaf blight, *Xanthomonas oryzae* pv. *oryzae*, Survey, Disease incidence, Symptomatology

Comment [mm1]: Please add methods, not only material

INTRODUCTION

Rice is the staple food of Asia where about 90% of the world's rice is produced and consumed. It was reported that rice production in the world was 503.27 million tonnes and in India, was to be around 124 million tonnes in 2022-2023. The major constraint in the rice production and productivity of rice was irregular rainfall, pests and diseases (Firdousee *et al.*, 2023).

Among various diseases, the bacterial disease caused by *Xanthomonas oryzae* pv. *oryzae* is the most devastating disease due to its high epidemic potential. Yield loss due to bacterial blight can reach up to 70% when susceptible varieties are cultivated and can be as high as 100% under severe conditions in environments conducive to the disease.

The disease was first time reported by farmers of Fukuoka area of Japan in 1884-1885 (Yamanukiet *al.*, 1962, Tagami and Mizukami, 1962). In India, it was first reported during the 1950s by Srinivasan *et al.* (1959). Earlier the disease occurred in an endemic form in Maharashtra. Later it broke out in epidemic form in the Shahabad district of Bihar in 1963 (Srivastava and Rao, 1963) since then, has become one of the important diseases of rice plants.


Bacterial leaf blight can affect rice crops both in the highlands and lowlands, but particularly in lowland rice, and often appears during the rainy season (Shankara *et al.*, 2017). *Xanthomonas oryzae* pv. *oryzae* enters through hydathodes at the leaf tip and leaf margin, multiplies in the intercellular spaces, migrates to the xylem vessels and causes systemic infection (Banerjee *et al.*, 2018).

Comment [mm2]: Alphabetic

Surveying the disease over time reveals how severely it impacts production and quality, in addition to revealing the presence of different races in various agro climatic zones. Therefore, this is an effort to know the disease status of bacterial leaf blight in major rice growing areas.

The current study aims to obtain a more detailed and accurate estimate of bacterial leaf blight occurrence and to know the prevalence and distribution of *Xanthomonas oryzae* pv. *oryzae* pathotypes in Andhra Pradesh. As the new pathotypes continue to emerge, majority of them varied in their virulence depending upon areas and fields within the area. Continuous monitoring of

the virulence structure of pathogen in an area is a prerequisite to make resistance breeding more durable and sustainable.



Formatted: Space After: 10 pt

Formatted: Justified, Indent: First line: 0.5", Line spacing: single, Pattern: Clear (White)

Figure 1. Map showing survey locations of bacterial leaf blight from major rice growing areas of Andhra Pradesh.

Comment [mm3]: ?

Comment [mm4]: Figure 1 move to results

MATERIALS AND METHODS

Aroving survey was conducted during *latekharif*, 2023-2024 to determine the prevalence of bacterial leaf blight disease in twelve major rice growing districts of Andhra Pradesh viz., West Godavari, East Godavari, NTR district, Krishna, Kakinada, Eluru, Nellore, Tirupati, Chittoor, Kurnool, Nandyal, Kadapa. Survey depended on the field diagnosis of the disease based on the plant symptoms and further confirmation in the laboratory by pathogenicity tests.

Diseased samples showing characteristic symptoms of bacterial leaf blight were collected, properly labelled with date of collection, variety name, place along with GPS location etc. and sealed in polythene bags. Additional information on the parameters like variety name, place, crop growth stage, crops in neighbouring fields and plant parts affected were recorded. Random sampling was conducted from each field and per cent disease incidence was calculated using the following formula.

$$\text{Disease Incidence} = \frac{\text{No. of diseased plants}}{\text{Total no. of plants observed}} \times 100$$

Pathogenicity test was performed to confirm the presence of bacterial leaf blight infection from randomly collected diseased samples. For this, 72hr old isolates collected from each district and multiplied on nutrient broth is clip inoculated onto 40day old seedlings of susceptible cultivar TN₁ using sterilized scissors (Kauffman *et al.*, 1973).

RESULTS AND DISCUSSION

The survey conducted in this study revealed that the average disease incidence ranged from 20 -72%. The lowest disease incidence was recorded in Burugupudivillage of Kakinada district (21.02%), Kalaparru village of Elurudistrict (25.6%) and Chinnadandluru village (28.6%) of Kadapa district. Highest incidence was observed in the fields Putrela village (72.34%) of Krishna district, Research plots of ARS, Nellore (59.25%), Davuluru village (54.5%) of NTR district, Perumalapalli village (49.12%) of Tirupati district (Table 1 and Fig. 1) as there were optimum environmental conditions for the disease development.

Xanthomonas oryzae pv. *oryzae* bacteria infects rice crop at all growth stages. Infected plants display symptoms like greyish leaf tip discoloration, dried leaves, water-soaked lesions that enlarge with wavy margins, and straw-yellow colour across the entire leaf. In susceptible varieties such as TN₁ and BPT-5204, symptoms become systemic, resembling burning. Infection during the flowering stage in the NLR 3228 variety, observed in Tirupati district, leads to partial or incomplete grain filling. Water-soaked lesions and leaf blight symptoms are found across Andhra Pradesh, regardless of growth stage.

Younger plants exhibit Kresiek (seedling blight or wilt phase) which is prevalent in Putrela village of Krishna district, which is also seen in West Godavari. Older plants commonly show water-soaked, translucent lesions turning straw yellow with wavy margins. Severe conditions can cause leaf blight and seedling wilt in Yerlampalli village of Chittoor district, and Burugupudi village of Kakinada district. Bacterial ooze on young lesions is noted in Nellore and Tirupati districts in the early morning. Kresiek is the most damaging symptom of BLB, while blight symptoms are the most common.

The above variation in the symptomology may be because of variable temperature and rainfall, as these two factors play important role in survival, disease development and spread. Similar information was reported by Garret *et al.* (2008) who suggested that climate change affects the level of disease through genomic, cellular, physiological levels of plant and pathogen.

As there is irregular increase in the temperatures and uneven rainfall over past ten years there is a sporadic outbreak of the disease. The data analysis on BB incidence with the weather parameters of rice growing seasons from 2015 to 2017 revealed that the rainfall distribution varied greatly within rice growing seasons over the years (Haque *et al.*, 2022).

Local climatic conditions, variability of the strain, rice variety, cultural practices adopted by the farmer can also be attributed for variation in symptomology and disease incidence which are in accordance with the results presented by Lubis *et al.* (2020). Climatic conditions play a vital role in the

Comment [mm5]: On methods: add references on all parameters

Comment [mm6]: Add references

Comment [mm7]: On discussion: Please add discussion for all parameters. Add references, explain the results from the others study, then compares with your study. Then placed discussion before conclusion

infection cycle, influencing the dispersal of the pathogen and its transmission between host plants (Naqvi, 2019).

Kresek symptoms in rice caused by *Xanthomonas oryzae* pv. *oryzae* occurs at 25-30°C which is optimum temperature for disease development. Delayed sowings and unseasonal rainfall were responsible for creating favourable conditions for BLB disease. Amin *et al.* (2022) also highlighted the significant impact of epidemiological factors like temperature, relative humidity, rainfall and sunshine hours etc. on the occurrence and spread of bacterial leaf blight caused by *Xanthomonas oryzae* pv. *oryzae* (Xoo). It is important to understand these epidemiological parameters to manage the disease efficiently.

Adhikari *et al.* (1994) found a strong correlation between the progression of bacterial leaf blight and environmental factors like rainfall and relative humidity. Additionally, Kapoor *et al.* (2004) equally suggested significant variations in rainfall and its distribution within growing seasons from 1979 to 1999. Understanding these parameters is crucial for effective disease management.

Formatted: Space Before: 0 pt



Figure 2. Blighted appearance of the field infected with BLB



3a. Kresek phase



3b. yellowish stripes at the margins



3c. straw yellow wavy margins

Figure 3. Symptoms at different growth stages of rice crop



Figure 4. Leaf inoculation of bacteria by leaf clipping method for pathogenicity test




Figure 5. Prevalence of BLB disease incidence across twelve districts of Andhra Pradesh



Figure 6. Rice bacterial blight disease scoring on 0–9 scale based on lesion length on the clip inoculated leaves.

CONCLUSION

The findings indicated that the disease is prevalent in all rice-growing areas, with varying levels of disease incidence ranged from 20 to 72 per cent and diverse pathogenic variations existed in the *Xanthomonas oryzae* pv. *oryzae* population. Further studies on pathotype identification and its distribution in these districts will solve the problems encountered in developing rice cultivars with specific and broad-spectrum resistance against *Xanthomonas oryzae* pv. *oryzae* population. This would aid in deploying better strategies for planning and incorporation of different Xa gene combinations for effective management against this pathogen.

Table 1: Disease incidence of bacterial leaf blight in major rice growing districts of Andhra Pradesh.

S. No.	District	Village	Variety name	Crop stage	Surrounding crop	Disease incidence	Pathogenicity test	Symptoms in field
1	West Godavari	Maruteru	MTU 2077	Kresek	Rice	40.84%	+ve	Kresek
2	East Godavari	Peravali	MTU 1318	Tillering stage	Rice	44.9%	+ve	Wavy margins
3	Eluru	Kalaparru	MTU 1121	Reproductive stage	Coconut orchad	25.6%	+ve	Yellowish wavy margins
4	Kakinada	Burugupudi	MTU 1064	Panicle initiation	Black gram	21.02%	+ve	Water soaked lesions
5	NTR district	Davuluru	MTU 1061	Vegetative stage	Banana orchad	54.5%	+ve	Wilted seedlings with straw coloured leaves
6	Krishna	Putrela	MTU 1010	Kresek	Rice	72.34%	+ve	Kresek stage
7	Nellore	ARS, Nellore	Bpt 5204	Tillering stage	Rice	59.25%	+ve	Ooze on leaf tips
8	Tirupati	Perumalapalli	NLR 3228	Panicle initiation	Maize	49.12%	+ve	Straw coloured wilted leaves
9	Chittoor	Yerlampalli	RNR 15048	Vegetative stage	Guava orchad	39.7%	+ve	Water soaked lesions
10	Kadapa	Chinnadandluru	IR 93	Vegetative stage	Tomato	28.6%	+ve	Water soaked lesions with yellow wavy margins
11	Nandyal	Bukkapuram	Nandyalson a	Vegetative stage	Rice	44.4%	+ve	Yellowish wavy margins
12	Kurnool	Veldurthi	Jilakarason a	Tillering stage	Groundnut	40.8%	+ve	Water soaked yellow stripes

References

- Firdouse, M.F., Sudha, A., Johnson, I., Anandham, R., Saranya, N., Amala, A.S and Arunprakash, S., 2023. Survey and Isolation of Rice Bacterial Leaf Blight Caused by *Xanthomonas oryzae* pv. *oryzae*. *International Journal of Plant & Soil Science*. 35(19):1755-1760.
- Yamanuki, S., Nakamura, K., Kayano, M., Narita, T and Iwata, T. 1962. First occurrence of bacterial leaf blight of rice in Hokkaido. *Annual Phytopathology*.264.
- Tagami, Y and Mizukami, T. 1962. Review of the researches on bank roll leaf blight of rice caused by *Xanthomonas oryzae*. Special report of the plant disease and insect pests forecasting service no. 10. Plant Prot. Div. Ministry Agric. For. Tokyo Japan.112.
- Srinivasan, M. C., Thirumulachar, M.J., Patel, M. K. 1959. Bacterial blight of rice. *Current Science*. (28):469-470
- Srivastava, D.N and Rao, Y.P., 1963. Epidemic of bacterial blight disease of Rice in North India. 16(4):393-394.
- Shankara, K., Patil, M. B., Pramesh, D., Sunkad, G., Yenjerappa, S. T., Ibrahim, M., Rajesh, N, L and Chikkannaswamy. 2017. Characterization of *Xanthomonas oryzae* pv. *oryzae* Isolates from Rice Growing Regions of Southern India. *International Journal of Pure and Applied Biosciences*. 5(4):452-461.
- Banerjee, A., Roy, S., Bag, M.K., Bhagat, S., Kar, M.K., Mandal, N.P., Mukherjee, A.K. and Maiti, D.2018. A survey of bacterial blight (*Xanthomonas oryzae* pv. *oryzae*) resistance in rice germplasm from eastern and northeastern India using molecular markers. *Crop protection*. (112):168-176.
- Kauffman, H. E., A. Reddy, S. P. Y. Hsieh and S.D. Merca. 1973. An improved technique for evaluating resistance of varieties to *Xanthomonas oryzae* pv. *oryzae*. *Plant Dis. Rep*. (57):537- 541.
- Garrett, K.A., Dendy, S.P., Frank, E.E., Rouse, M.N and Travers, S.E., 2006. Climate change effects on plant disease: genomes to ecosystems. *Annu. Rev. Phytopathol*.44(1): 489-509.
- Haque, Md. Mahbulul & Mita, Mamuna Mahjabin & Bashar, Samrin & Masud, Md Mostafa & Protic, Ismam Ahmed & Mozumdar, Uttam & Islam, Md. (2022). Weather parameters largely regulate the outbreak of rice bacterial blight: an endemic disease poses a threat for food security in Bangladesh. *Asian-Australasian Journal of Food Safety and Security*. (6):35-47.
- Lubis, K., Maathuis, F.J.Mand Safni, I., 2020. Mapping bacterial leaf blight disease of rice (*Xanthomonas oryzae* pv *oryzae*) in North Sumatra. In *IOP Conference Series: Earth and Environmental Science*(454)1:012160. IOP Publishing.
- Naqvi SAH, 2019. Bacterial leaf blight of rice: an overview of epidemiology and management with special reference to Indian sub-continent. *Pak. J. AgricRes*. 32:359.
- Amin, T., Vishal, G., Aarushi, S and Saima Khushboo, S.2022. Effect of Weather Parameters on the Severity of Bacterial Leaf Blight of Rice (cv. Basmati-370). *Biological Forum – An International Journal*. 2249-3239

Comment [mm8]: On references: add or change references using limited 5 years ago

Adhikari TB, TW Mew and PS Teng, 1994. Phenotypic diversity of *Xanthomonas oryzae* pv. *oryzae* in Nepal. *Plant Dis.* 78:68–72.

Kapoor S, NSK Harsh and SK Sharma, 2004. A new wilt disease of *Acacia nilotica* caused by *Fusarium oxysporum*. *J. Trop. For. Sci.* 16:453-462.

UNDER PEER REVIEW

