

2 **MANAGEMENT OF SUNFLOWER POWDERY MILDEW AND *Alternariaster* LEAF**
3 **SPOT: HARNESSING THE POTENTIAL OF COMBINED NEW FUNGICIDE**
4 **MOLECULES**

5
6 **ABSTRACT**

7 The study investigates the efficacy of seed treatment with Fluxapyroxad FS @ 1.5 g/kg
8 seed followed by foliar sprays of various fungicide combinations in managing powdery mildew
9 and *Alternariaster* leaf spot diseases in sunflower. Conducted over three Kharif seasons (2021-
10 2023), the trials consistently demonstrated the superior performance of Fluopyram 17.7% +
11 Tebuconazole 17.7% (400 SC) @ 1 ml/L in reducing disease incidence and severity. In Kharif
12 2021, this treatment recorded the lowest incidence of *Alternariaster* leaf spot (7.36%) and
13 powdery mildew (4.52%), outperforming other treatments. Similar trends were observed in
14 Kharif 2022 and 2023, with Fluopyram + Tebuconazole maintaining the lowest disease severities
15 (7.39% and 2.95% for powdery mildew, respectively). Notably, the study found no significant
16 differences in yield components across treatments, though the highest yields were consistently
17 recorded in the treatment combining Fluxapyroxad seed treatment with Fluopyram +
18 Tebuconazole foliar spray, achieving yields of 2124.67 kg/ha in Kharif 2022 and 2157.33 kg/ha
19 in Kharif 2023. These results indicate that integrated disease management using these fungicide
20 combinations can effectively control both diseases in sunflower, particularly under humid
21 weather conditions, without adversely affecting crop yield.

22 **Keywords:** Sunflower, *Alternariaster* leaf spot, Powderymildew, Disease severity, Fungicides

23 **INTRODUCTION**

24 Sunflower is an important oilseed crop as it contains good quality oil with pleasant
25 flavour and excellent keeping quality when refined. The oil is rich in Linoleic acid (62.5%)
26 and is recommended as dietary constituent to heart patients. Sunflower oil is also used in
27 industrial production of soaps, cosmetics, paints, lubricants, plastics, etc. Its seed cake is an

28 important by product which is used as concentrates for cattle, pigs, sheep, goat and poultry as
29 it is a protein rich feed (40-50%). Thus, sunflower being a multipurpose oil seed crop,
30 naturally the demand for crop is increasing tremendously. These demands resulted into
31 extensive cultivation of sunflower over large areas leading to development of diseases in the
32 crop which has considerable impact on grain yield of sunflower.

33 “Diseases are one of the major constraints in successful sunflower cultivation”. [1]Over
34 80 pathogens occur on sunflower. In Karnataka, the major diseases of sunflower are necrosis
35 virus, *Alternaria* blight, powdery mildew, rust, collar rot and downy mildew. Among these
36 *Alternaria* blight caused by *Alternariahelianthi* (Hansf.) [2] has been considered as a
37 potentially destructive disease in many parts of sunflower growing countries [3] and also in
38 some regions of Karnataka [4]. In India, powderymildew caused by
39 (*Erysiphecichocearum*) was first reported on sunflower in 2008 [5]. “A field survey on
40 powdery mildew in seven districts of Karnataka recorded 30–74 % disease severity”[6-7].

41 *Alternaria* leaf blight is recognized as a major disease in more humid areas. Favorable
42 weather may result in yield losses up to 62 per cent which occurred during 1988 in
43 Karnataka(Anon, 2007). [8]. The disease occurred with incidence as high as 95-100 per cent
44 due to heavy rains during flowering and grain filling stage of the crop [8 &9]. It appears on
45 all parts of the crop severely during favourable weather conditions.Powderymildew is
46 worldwide in distribution, but greatest severity is observed in the tropics, where it appears
47 during advanced senescence at the floweringor post floweringstages [10]. The disease
48 primarily affects the leaves, flowers, and head. It appears as a white, powdery fungal growth
49 on the affected parts, leading to reduced photosynthetic area and reduced oil content. The
50 disease thrives in warm, humid conditions and can cause considerable yield losses if not
51 managed properly through cultural practices and fungicide applications. Disease losses due to
52 powdery mildew was reported as much as 70 per cent with significant reduction in oil
53 quantity (17%) and quality (33%) [11].

54 In both the diseases high inoculum in the field coinciding with favourable
55 environmental conditions lead to early infections causing severe losses. Regarding
56 symptoms, the disease appears in the form of whitish powdery patches in Powdery mildew
57 and as dark necrotic spots in *Alternaria*,both appear on leaves, stem, petioles and also on

68 capitulum. The use of fungicides has become an inevitable method for controlling of the
69 disease in the absence of resistant cultivars over large areas. Often, commonly used fungicides
60 fail to give satisfactory control of the diseases under field conditions, particularly during the
61 humid weather conditions. Therefore, it is essential to find out potent fungicides, which
62 would be effective in controlling the disease. In the present study, combination fungicide
63 products were used in the field to test the efficacy under natural weather conditions against
64 powdery mildew and *Alternaria* leaf spot of Sunflower. Loss of powdery mildew mentioned
65 in above with ref 11

66 MATERIAL AND METHODS

67 Management of the diseases was carried out in the experimental plot located at Zonal Agriculture
68 Research Station, GKVK, University of Agriculture Sciences, Bangalore, Karnataka during the
69 late *Kharif* (July – August) season. The trials were carried out for three seasons 2021,2022 and
70 2023. Seven treatments were laid out with three replications. The trial was carried out in RCBD
71 design with plot size of 4.2X3.0M². The susceptible Hybrid KBSH 44 was sown after seed
72 treatment with fluxapyroxad FS @ 1.5g/kg (except for untreated check), 1kg of seeds was taken
73 in seed pan to which 1.5ml of the suspension was added completely and the pan was rotated
74 around so the seeds are uniformly covered with the chemical. The treated seeds were shade dried
75 for one hour and then sown and spray was taken during the initiation of disease symptoms. Two
76 sprays were given at fortnight interval for the effective disease reduction. Observations were
77 carried out on the per cent disease severity (Incidence is literal, scientific research and recording
78 the diseases fall under severity), seed yield and B:C ratio. Disease incidence was rated based on
79 the scale provided (IIOR, Hyderabad technical compendium). Periodically observations were
80 recorded based on the lesion symptoms and graded following the standard scale. Disease scoring
81 (0-9 scale) for *Alternaria* leaf spot, similarly Powdery mildew severity is recorded by scoring
82 the per cent leaf area covered by the disease on leaves and on upper portion of leaves through
83 visual observation using a rating scale of 0-9 scale (Table 1 and Table 2). Seed yield for each
84 treatment (three replications) was averaged and converted to per hectare. Benefit cost ratio was
85 calculated based on the excess cost incurred due to spraying and more seed yield reflected in
86 terms of benefit.

87

88

89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118

RESULT

Seed treatment with Fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with different combined fungicides was carried out to study their effect on powdery mildew and *Alternariaster* leaf spot disease.

During *Kharif* 2021 seed treatment with T_1 fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with Fluopyram 17.7% + Tebuconazole 17.7% (400 SC) @ 1 ml/L (T_1) recorded lowest incidence of *Alternariaster* leaf spot disease (7.36%) followed by T_2 - Seed treatment with Fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with Azoxystrobin 18.2% + Difenconazole 11.4% SC @ 1ml /L – 11.23% (**Table 1**). Powdery mildew was also lowest in T_1 (4.52%) followed by T_6 (7.19%) - spray of myclobutanil 10% WP @ 0.5 g /L

The trial conducted during *Kharif* 2022, spraying with Fluopyram 17.7% + Tebuconazole 17.7% (400 SC) @ 1 ml/L (T_1) was the best treatment revealing good management of the powdery mildew disease with PDI of 7.39% followed by spray with Myclobutanil 10% WP @ 0.5 g /L (9.49%) (**Table 4**). Whereas, T_1 was also effective against *Alternaria* leaf spot with reduced disease severity of 4.28 per cent and followed by the next best treatment, spray with Azoxystrobin 18.2% + Difenconazole 11.4% SC @ 1ml /L (9.32%).

Trial conducted during *Kharif* 2023 recorded lowest disease severity of leaf spot (7.66%) and powdery mildew (2.95%) in seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Fluopyram 17.7% + Tebuconazole 17.7% (400 SC) @ 1 ml/L. This was followed by seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Myclobutanil 10% WP @ 0.5 g /L (4.69%) for powdery mildew and Azoxystrobin 18.2% + Difenconazole 11.4% SC @ 1ml /L (9.47%) for leaf spot (**Table 3**).

The trial conducted for three seasons revealed the efficient management of the severely emerging diseases in sunflower using the new fungicide combined molecules. Fluopyram 17.7% + Tebuconazole 17.7% (400 SC) recorded the lowest severity of both the diseases over three season trials. Two sprays of the fungicide molecules at fortnight interval were sufficient to manage both the diseases in the dryland crop during humid weather conditions.

119 Significant difference was not noticed in yield component among the different treatments. Yield
120 was higher in the T₁ but statistically not much difference between treatments compared to control
121 which was statistically lower yield compared to T₁. But highest yield was recorded in T₁ - Seed
122 treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with fluopyram 17.7%+
123 tebuconazole 17.7% (400 SC) @ 1 ml/L (2124.67 kg/ha) during *Kharif* 2022 and *Kharif* 2021
124 (2153 kg/ha) compared to control. Similarly, during *Kharif* 2023 not much difference among the
125 treatments was recorded in seed yield and highest was in T₁ (2157.33kg/ha)(Table 3).

126

127 DISCUSSION

128 “The fungicide management of foliar diseases for the last decade, has been based on the
129 use of demethylation inhibitors (DMI, Group 3), Triazole fungicides, and quinone outside
130 inhibitors (QoI or Strobilurins, Group 11 fungicides)”. [12].

131 This shows that fluxapyroxad was an effective fungicide for seed treatment in sunflower
132 by protecting it upto 30-45 days after sowing from the fungal diseases. The chemical protects the
133 crop upto 30 days effectively for the disease that appears 30 DAS spray has to be taken after the
134 disease appearance. Fluopyram, a new succinate dehydrogenase inhibiting fungicide (SDHI),
135 registered for use in pistachio for the control of *Alternaria* late blight (ALB) as products
136 premixed with demethylation inhibitor (DMI) tebuconazole was effective against strains showing
137 Boscalid resistance [13]. “*Alternaria* leaf spot, maximum per cent disease control occurred when
138 the apple crop was sprayed with Fluopyram 200 + Tebuconazole 200 SC at a concentration of
139 0.0625 per cent (94.55 %) followed by Fluopyram 200 + Tebuconazole 200 SC at 0.05 per cent
140 (89.42%) concentration and 0.0375 per cent (86.55 %), respectively” [14]. There is no use of
141 combination products in managing fungal diseases in sunflower or other oilseed crop.

142

143 Fluopyram 200 + Tebuconazole 200 – 400 SC at 750 ml/ha concentration were found to
144 be significantly superior over control with incidence of powdery mildew (25.10%) and
145 postharvest disease incidence (15.00%) as compared to other chemical treatments in mango crop.
146 The efficacy of the product, Fluopyram 200 + Tebuconazole 200 SC (Luna Experience), against
147 powdery mildew in pepper was found to be 89 per cent in Timorex Gold assay efficacy against
148 *Leveillulataurica*, and it was safe for use against powdery mildew [15-16]. The fungicide product
149 Fluopyram + Tebuconazole (400 SC) tested in many commercial crops like Mango, apple,

150 pistachio, tomato, grapes, gherkin etc were highly efficient and had safety. Hence, it can be used
151 against Alternaria leaf spot and powdery mildew in sunflower with high disease management
152 efficiency giving good results on yield factor also. Use of such product in large scale under
153 dryland cultivation of food crops in India paves way for ease in disease management at single go
154 for the long term. This is not seed treatment studies it is combination of both seed treatment and
155 spray to manage the disease in the entire crop period.

156

157 CONCLUSION

158 The study demonstrates that seed treatment with Fluxapyroxad FS @ 1.5 g/kg followed
159 by foliar sprays with Fluopyram 17.7% + Tebuconazole 17.7% (400 SC) @ 1 ml/L effectively
160 manages powdery mildew and *Alternariaster* leaf spot diseases in sunflower across multiple
161 Kharif seasons. This treatment consistently recorded the lowest disease incidence and severity,
162 outperforming other fungicide combinations. Additionally, the highest seed yields were obtained
163 with this treatment, indicating its efficiency in disease control without negatively impacting crop
164 yield or quality. Two foliar sprays at a fortnight interval were sufficient to manage these diseases
165 under humid weather conditions with high efficiency. Further research could explore the long-
166 term sustainability and environmental impact of these fungicide treatments. Whereas, resistant
167 sunflower sources against the diseases could enhance the overall disease management and reduce
168 dependency on chemical fungicides which could make way for pathogen resistance against
169 fungicides.

170 This is just a hypothesis statement made not tested and laid out rule. Phytotoxicity was
171 not observed in the sunflower field or other crops tested by me for the dosage given in the
172 treatments. This is just testing not recommending to the farmers as recommendation has to go
173 long process through OFT, FLD and FD etc. extension procedures

174 ETHICAL STATEMENT

175 All the experimental procedures involving only on plant species were conducted
176 following the University of Agricultural Science, Bangalore institutional guidelines. There are no
177 human and animal subjects/trials conducted in this article and informed consent is not applicable.

178

179 **Disclaimer (Artificial intelligence)**

180 Option 1:

181 Author(s) hereby declare that NO generative AI technologies such as Large Language Models
182 (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or
183 editing of manuscripts.

184 Option 2:

185 Author(s) hereby declare that generative AI technologies such as Large Language Models, etc
186 have been used during writing or editing of manuscripts. This explanation will include the name,
187 version, model, and source of the generative AI technology and as well as all input prompts
188 provided to the generative AI technology

189 Details of the AI usage are given below:

190 1.

191 2.

192 3.

193

194

195 **REFERENCES**

196 1. Gulya TJ, Rashid KY, Masirevic SM. Sunflower diseases, Sunflower Technology
197 and Production. Agron. Monogr.35, ASA, CSSA, and SSSA, Madison, WI, 1997;
198 21–65.

199 2. Tubaki K, Nishihara N. *Alternariahelianthi* (Hansf.) comb. nov. Trans Br Mycol
200 Soc, 1969; 53: 147–149.

201 3. Allen SJ, Brown JF, Kochman JK. Effect of temperature, dew period and light on
202 the growth and development of *Alternariahelianthi*. Phytopathol, 1983; 73: 893-
203 896.

204 4. Shankergoud I, Shadakshari YG, Parameshwarappa KG, Chandranath HT, Katti
205 P, Mesta RK. Sunflower and castor research in Karnataka-an overview.

- 206 University of Agricultural Sciences, Dharwad, 2006; 41.
- 207 5. Baiswar P, Kumar R, Chandra S, Ngachan SV. First report of powdery mildew on
208 Mexican sunflower in India. 2009.
- 209 6. Rao SC, Sujatha M, Karuna K, Varaprasad K. Powdery mildew disease in
210 sunflower: A review. The Indian Society of Oilseeds Research, 2006; p. 111.
- 211 7. Dinesh B, Kulkarni S, Harlapur SI, Benagi VI, Mallapur CP. Karnataka J Agric.
212 Sci., 2010; 23: 521–523.
- 213 8. Karuna K, Jagadish KS, Geetha KN, Shadakshari YG. Evaluation of efficacy of
214 chemical fungicides and a plant product for the management of Alternaria blight
215 of sunflower. Indian Phytopath, 2012; 65: 305-306.
- 216 9. Reddy KP, Rao SC, Kirti PB, Sujatha M. Development of a scoring scale for
217 powdery mildew (*Golovinomyces cichoracearum* (DC.) VP Heluta) disease and
218 identification of resistance sources in cultivated and wild sunflowers. Euphytica,
219 2013; 190: 385-399.
- 220 10. Zimmer DE, Hoes JA. Diseases. In: Carter JF (ed) Sunflower Science and
221 Technology. Agronomy Monograph, 1978; 19: 225-262. ASA, CSSA, and SSSA,
222 Madison, WI.
- 223 11. Kolte SJ. Sunflower diseases, Diseases of Annual Edible Oilseeds Crops, III,
224 CRC Press, Florida, 1985; 194 pp.
- 225 12. Vagelas I, Cavalari C, Karapetsi L, Koukidis C, Servis D, Madesis P. Protective
226 effects of Systiva® seed treatment fungicide for the control of winter wheat foliar
227 diseases caused at early stages due to climate change. Agronomy, 2022; 12(9):
228 2000.
- 229 13. Avenot HF, van den Biggelaar H, Morgan DP, Michailides TJ. Fungicidal activity
230 of fluopyram for suppression of Alternaria species pathogenic on California
231 pistachio. Resistant Pest Management Newsletter, Fall 2012; 22(1): 10-14.
- 232 14. Verma S, Khosla K. Chemical management of marssonina leaf blotch and
233 alternaria leaf spot diseases of apple in Himachal Pradesh. International Journal of
234 Chemical Studies, 2018; 6(4): 3316-3319.

- 235 15. Patil PD, Raut RA, Karande RA, Haldavanekar PC. Bio-efficacy and
236 phytotoxicity study of fungicide fluopyram 200+ tebuconazole 200-400SC
237 against powdery mildew and postharvest diseases of mango. *Int J Chem Stud*,
238 2017; 5: 667-670.
- 239 16. Arici ŞE, Özkaya R. Evaluation of biofungicides in the control against powdery
240 mildew disease [*Leveillulataurica* (Lev.) Arm.] in pepper. *KahramanmaraşSütçü*
241 *İmam ÜniversitesiTarımveDoğaDergisi*, 2022; 25(2): 274-281.
- 242 17. Anonymous 2007. Annual Progress Report of AICRP on Sunflower (2006-07).
243 Directorate of Oilseeds Research, ICAR, Hyderabad, 230 pp.

244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261

UNDER PEER REVIEW

262

263

264 **Table 1:** Severity of *Alternaria* leaf spot Disease Measured by Percentage Leaf Area
265 Coverage and Visual Rating Scale (0-9) on Upper Portion of Leaves

Score	Reaction	Leafareacovered
0	Immune	Nosymptomsontheleaf
1	Highlyresistant (HR)	Small,circular,scatteredbrownspotscovering1%orless oftheleafarea
3	Resistant(R)	Spotsenlarging,darkbrownincolourcovering1-10%of leafarea
5	Moderately resistant/ Moderatelysusceptible (MR/MS)	Spotsenlarging,darkbrownincolour,targetlikeappearance covering11-25%of leafarea
7	Susceptible(S)	Spotsdarkbrown,coalescing,withtargets p o t - likeappearance covering26-50%of leafarea
9	Highlysusceptible (HS)	Spotsuniformlydarkbrown,coalescingcovering51%or aboveofleaf area

266 **Table 2:** Powdery Mildew Severity Scoring System Based on Leaf Area Coverage and Disease
267 Reaction

Score	Reaction	Leafareacovered
0	Immune	Nosymptomsontheleaf
1	Highlyresistant (HR)	Powderymildewspeckscovering1%or lessleafarea.
3	Resistant(R)	Powderylesionscovering1-10%of leaf area.
5	Moderately resistant/ Moderatelysusceptible (MR/MS)	Enlargedpowderylesionscovering11-25%ofleaf area.
7	Susceptible(S)	Powderylesionscoalescetoformbigpatchescovering 26-50%ofleaf area.
9	Highlysusceptible (HS)	Powderypatchescovering:51%ormoreofleafarea anddefoliationoccur

268

269

270

271

272

UNDER PEER REVIEW

Table 3: Effect of Different Fungicide Treatments on *Alternariaster* Leaf Spot, Powdery Mildew, Seed Yield, and Benefit-Cost Ratio in Sunflower (During *Kharif*2021)

Trt. No.	Treatment	<i>Alternariaster</i> leaf spot (%)	Powdery mildew (%)	Seed yield (Kg/ha)	B:C ratio	Increment B:C ratio (IBCR)
T ₁	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with fluopyram 17.7%+ tebuconazole17.7% (400 SC) @ 1 ml/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	7.36 (15.74)	4.52 (12.27)	2153 (46.41)	0.74	628.38
T ₂	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with azoxystrobin 18.2% + difenconazole 11.4% SC @1ml /L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	11.23 (19.58)	16.8 (24.20)	2113 (45.97)	0.73	582.19
T ₃	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with boscalid 25.2% + pyraclostrobin 12.8% WG @ 1g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	12.29 (20.52)	9.42 (17.87)	2010 (44.84)	0.70	460.00
T ₄	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with dimethomorph 12% + pyraclostrobin 6.7% @1.5	18.75 (25.66)	11.52 (19.84)	2197 (46.88)	0.68	748.53

	g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)					
T ₅	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with picoxystrobin 7.05%+ propiconazole 11.71% SC @ 2 g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	14.68 (22.53)	10.1 (18.53)	1973 (44.42)	0.71	401.41
T ₆	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with myclobutanil 10% WP @ 0.5 g /L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	20.42 (26.86)	7.19 (15.55)	1865 (43.19)	0.68	260.29
T ₇	Control	23.98 (29.32)	22.12 (28.06)	1688 (41.09)	-	
	CV(%)	5.66	2.19	12.56		
	SEm+/-	0.98	0.73	0.745		
	CD@5%	0.507	9.93	1.439		

Table 4: Effect of Different Fungicide Treatments on *Alternariaster* Leaf Spot, Powdery Mildew, Seed Yield, and Benefit-Cost Ratio in Sunflower (During *Kharif* 2022)

Treatments		Powdery mildew (%)	<i>Alternariaster</i> leaf spot (%)	Seed yield (kg/ha)	B:C ratio	Incr B:C (IBC)
T ₁	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Fluopyram 17.7%+ Tebuconazole17.7% (400 SC) @ 1 ml/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	7.39 (15.68)	4.28 (11.89)	2124.67 (46.10)	0.85	
T ₂	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Azoxystrobin 18.2% + Difenconazole 11.4% SC @1ml /L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	14.72 (22.48)	9.32 (17.75)	2096.33 (45.79)	0.81	
T ₃	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Boscalid 25.2% + Pyraclostrobin 12.8% WG @ 1g/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	21.19 (27.40)	18.60 (25.54)	2036.67 (45.13)	0.64	
T ₄	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar spray with Dimethomorph 12% + Pyraclostrobin 6.7% @1.5 g/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	20.80 (27.12)	19.64 (26.31)	2011.33 (44.85)	0.68	202.
T ₅	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar spray with Picoxystrobin 7.05%+ Propiconazole 11.71% SC @ 2 g/L (first spray at the onset of incidence or 45 days after	15.75 (23.35)	16.97 (24.31)	2066.67 (45.47)	0.80	

	sowing and second spray 15 days after 1st spray)					
T ₆	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Myclobutanil 10% WP @ 0.5 g /L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	9.49 (19.77)	12.33 (20.51)	2068.33 (45.48)	0.79	
T ₇	Control	28.79 (30.52)	25.19 (30.10)	1873.33 (43.28)		
	CV (%)	8.538	7.77	17.168		
	CD @ 5%	2.532	2.137	96.783		
	Sem (+/-)	0.896	0.744	38.584		

Table 5: Effect of Different Fungicide Treatments on *Alternariaster* Leaf Spot, Powdery Mildew, Seed Yield, and Benefit-Cost Ratio in Sunflower (During *Kharif* 2023)

Treatments		Powdery mildew (%)	<i>Alternariaster</i> leaf spot (%)	Seed yield (kg/ha)	B:C ratio	Increment B:C ratio (IBCR)
T ₁	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Fluopyram 17.7%+ Tebuconazole 17.7% (400 SC) @ 1 ml/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	2.95 (9.86)	7.66 (16.02)	2157.33 (46.10)	0.90	243.33
T ₂	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Azoxystrobin 18.2% + Difenconazole 11.4% SC @1ml /L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	6.66 (14.92)	9.47 (17.81)	2119.67 (45.79)	0.86	210.86
T ₃	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Boscalid 25.2% + Pyraclostrobin 12.8% WG @ 1g/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	17.44 (24.65)	16.63 (24.03)	2049.33 (45.13)	0.63	176.19
T ₄	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar spray with Dimethomorph 12% + Pyraclostrobin 6.7% @1.5 g/L (first spray at the onset of incidence or 45 days after sowing and second spray 15	20.14 (26.65)	20.95 (27.23)	2027.33 (44.85)	0.65	136.92

	days after 1st spray)					
T ₅	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar spray with Picoxystrobin 7.05%+ Propiconazole 11.71% SC @ 2 g/L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	12.42 (20.58)	15.70 (23.28)	2076.67 (45.47)	0.71	194.15
T ₆	Seed treatment with Fluxapyroxad FS @ 1.5 g/kg seed followed by foliar sprays with Myclobutanil 10% WP @ 0.5 g /L (first spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	4.69 (12.41)	22.07 (28.01)	2105.00 (45.48)	0.84	198.42
T ₇	Control	27.99 (31.93)	30.93 (33.78)	1938.33 (43.28)		
	CV (%)	6.835	10.452	16.802		
	CD @ 5%	1.910	3.019	152.488		
	Sem (+/-)	0.678	1.046	53.074		

Note: values in parenthesis are arcsin transformed

Table 6: Pooled analysis

Trt. No.	Treatment	<i>Alternariaster</i> leaf spot (%)	Powdery mildew (%)	Seed yield (Kg/ha)	B:C ratio
T ₁	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with fluopyram 17.7%+ tebuconazole17.7% (400 SC) @ 1 ml/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	6.43 (14.69)	4.95 (12.86)	2145.00 (46.32)	0.74
T ₂	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with azoxystrobin 18.2% + difenconazole 11.4% SC @1ml /L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	10.01 (18.44)	12.73 (20.90)	2109.67 (45.94)	0.73
T ₃	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with boscalid 25.2% + pyraclostrobin 12.8% WG @ 1g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	15.84 (23.45)	16.02 (23.59)	2032.00 (45.08)	0.70

T ₄	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with dimethomorph 12% + pyraclostrobin 6.7% @1.5 g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	19.78 (26.41)	17.49 (24.72)	2078.55 (45.60)	0.68
T ₅	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar spray with picoxystrobin 7.05%+ propiconazole 11.71% SC @ 2 g/L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	15.78 (23.41)	12.76 (20.93)	2038.78 (45.16)	0.71
T ₆	Seed treatment with fluxapyroxad FS @ 1.5g/kg seed followed by foliar sprays with myclobutanil 10% WP @ 0.5 g /L (First spray at the onset of incidence or 45 days after sowing and second spray 15 days after 1st spray)	18.27 (25.31)	7.12 (15.48)	2012.78 (44.87)	0.68
T ₇	Control	26.70 (31.11)	26.30 (30.85)	1833.22 (42.82)	-
	CV(%)	17.24	20.17	14.01	
	SEm+/-	1.604	2.022	47.246	
	CD@5%	4.098	6.031	141.233	

UNDER PEER REVIEW

Table 7: Weather data (AVR)

Weather parameters(Average)	Year	Months			
		July	August	September	October
Rainfall	2023	115.2	25.8	194	67.6
Temp. (Max °C)		28.3	30.5	29.3	30.1
Temp. (Min °C)		19.6	20.2	19.8	19.5
RH (Max)		81	84	86	87
RH (Min)		61	58	64	57
Rainfall	2022	149.8	211.4	142	361
Temp. (Max °C)		27.3	27.4	27.7	27.0
Temp. (Min °C)		18.7	18.6	18.5	16.9
RH (Max)		90	91	90	90
RH (Min)		61	62	59	61
Rainfall	2021	171.6	115.4	166.8	231.6
Temp. (Max °C)		28	28.4	28	28.1
Temp. (Min °C)		19.1	18.8	19	18.7
RH (Max)		87	89	91	90
RH (Min)		59	58	58	59