


Original Research Article

Evaluation of different [varieties of](#) miniature roses (*Rosachinensis minima*) under Prayagraj agro-climatic conditions



Abstract

An experiment titled “Evaluation of different miniature roses (*Rosa chinensis minima*) under Prayagraj agro-climatic conditions” was carried out in the Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology And Sciences, Prayagraj, from November, 2023 to May, 2024. The experiment was conducted in ~~a~~ Randomized Block Design with twelve varieties replicated thrice. Varieties were Red Hot, Magic Carousel, Rosy, Calpoly, Beauty Secret, Hocky Pocky, Ruby Ruby, Rainbow's End, Jet Trail, Cricket, Black Jade, and Jeane Kennealy. Variety Ruby Ruby emerged promising for parameters like plant height (25.9cm), number of branches (10.9), duration of flowering (164 days), ~~and~~ number of ~~flower~~ flowers per plant (285.9) whereas, plant spread (31.4 cm²) was better in variety Jeane Kennealy. Variety Red Hot performed better in day stake to first flowering (32 days). Bud diameter (1.1cm) was better in ~~the~~ variety Magic Carousel.

Keywords: Miniature rose, varieties, growth, flowering

1. Introduction

Roses are particularly one of the most attractive and ~~in demand~~ flowers among the world's various floriculture crops. ~~Rose~~ Rose's word springs from a Latin word “Erose” denoting ~~that~~ the “God of love” (Rajesh and Ramesh, 1999). Rose, genus *Rosa*, ~~belong~~ belongs to ~~the~~ family ~~Rosaceae~~ Rosaceae and is one of ~~the~~ nature's most beautiful ~~creation~~ creations. It is universally acclaimed as the Queen of Flowers and is one of the world's most popular flowers. The history of miniature roses is relatively unknown.

Miniature roses first came as an accidental result of rose hybridization. Miniature roses are dwarf bushes with little leaves and flowers. According to ~~the~~ American Rose Society, miniature ~~rose~~ roses are of 3 ~~type~~ types of micro-mini rose is the smallest class which only ~~reach~~ reaches 12–14 inches in height and the bloom is

the size is very small when fully opened and can be grown in a flower pot. Large miniature ~~rose~~ roses are a little bigger than the ~~micro~~ mini roses, can get 2-3 feet high and ~~its~~ their blooms are 1–2 inches across. Another is mini flor which is a relatively new class of rose due to its size. Plants reach 3–4 feet in

Comment [am1]: Add 2-3 sentences to indicate the importance of the research and also the miniature rose

Comment [am2]: Add “a”

Formatted: Font: 12 pt, Font color: Red

Comment [am3]: Put the methodology of the research including what variables to be measured

Formatted: Font: 12 pt, Font color: Red

Formatted: Font: 12 pt, Font color: Red

Formatted: Font: 12 pt, Font color: Red

Comment [am4]: Use recent publications for Introduction

Formatted: Font: 12 pt, Font color: Red

Formatted: Font: 12 pt, Font color: Red

height and blooms can be 3 inches across. It got its name because it is bigger than a miniature and not as big as a Floribunda. Miniature roses have gained much popularity mainly because of ~~the~~ [their](#) diverse and interesting flower forms and ~~colour~~ [color](#) (except true blue or black) and growth habits. Miniature roses tend to be quite hardy and can be grown in beds and ~~in~~ containers. They do best in full sun, at least six to eight hours per day. Though they can tolerate a bit of shade, ~~often times~~ [often times](#) their foliage and flowers will become sparse in shady conditions.

Miniature roses have multipurpose uses planted in pots, garden ~~display~~ [displays](#), landscape plantings, miniature bushes, little cut bouquets, etc. They are stunning and cheerful in the garden. ~~They can~~ [They can](#) also be used as lovely low hedges and borders. Many miniature rose varieties make adorable little cut bouquets for the home and special occasions. Their small size makes miniature roses ideal for edging, mass plantings, and rock or herb gardens.

Today miniature roses are reproduced by the ~~million~~ [millions](#) in countries all over the world. In some countries, most miniature roses are sold as budded or grafted stock. Miniatures are sold at the exact point when they are in full bloom. The beauty of these miniatures is that they are produced in 11 weeks and thus are much more economical to market than those currently available. The increasing popularity of miniature roses motivated the hybridizers into breeding programs and successfully developed many new miniatures.

2. Materials and Methods

The field experiment was carried out in the Department of Horticulture, Sam Higginbottom University of ~~agriculture~~ [Agriculture](#), Technology And Sciences, Naini, Prayagraj, ~~during~~ [from](#) November, 2023 to May, 2024. Twelve varieties were used in the experiment, laid out in a Randomized Block Design, which was replicated thrice along with 40 cm x 40 cm spacing. Varieties Red Hot, Magic Carousel, Rosy, Calpoly, Beauty Secret, Hocky Pocky, Ruby Ruby, Rainbow's End, Jet Trail, Cricket, Black Jade, and Jeane Kennealy were used. The source of the grafted plants was Puspanjali Nursery, Jakpur, Kharagpur, West Bengal. Up to 120 days of observations on vegetative and floral parameters were made at ~~30 days~~ [30-day](#) intervals.

3. Results and Discussion

Vegetative parameters

The mean data for different vegetative parameters, including plant height, plant spread, and number of branches of miniature roses-rose varieties are represented in Table 1.

Significantly taller plants (25.9 cm) were observed in the variety Ruby-Ruby which was at par with variety Beauty Secret (25.1 cm), Magic Carousel (22.3 cm), Calpoly (22.1 cm), Jeane Kennealy (22 cm) and Rosy (21.8 cm) whereas shorter plants (13.9 cm) were observed in variety Jet Trail. Significantly more plant spread (31.4cm^2) was observed in variety Jeane Kennealy which was at par with variety Rosy (31.3cm^2), Beauty Secret (28.9cm^2), Magic Carousel (27.2cm^2), Rainbow's End (25.1cm^2), Ruby Ruby (24.4cm^2) and Black Jade (24.2cm^2) whereas lesser plant spread (15.9cm^2) was observed in variety Jet Trail. Significantly more number of branches per plant (10.9) was observed in the variety Ruby Ruby which was at par with variety Rainbow's End (9.9) and Hocky Pocky (8.6) whereas fewer number of branches per plant (6.3) was observed in the variety Jet Trail.

The variation among the varieties for different vegetative parameters like plant height, plant spread, and number of branches per plant may be due to differences in the genetic makeup and climatic conditions of the area. Plant height variation may be due to differences in rapid meristematic activity during the early growth stages which results in more cell division and elongation in plants. Similar variation-variations in plant height were recorded by Sharova *et al.*, (1977) in gladiolus, Singh *et al.*, (2013) in floribunda rose, Ramzan *et al.*, (2014) in *Rosa hybrida*, and Soujanya *et al.*, (2014) in rose. For flower crops, plant spread is a crucial growth characteristic. It helps to utilize the sunlight to the maximum extent. This was in accordance with following the findings of Mohanty *et al.*, (2011) in rose, Chandrashekar *et al.*, (1973) in hybrid tea rose, Dias and Patil *et al.*, (2003) in elite rose, and Sharma and Sharma *et al.*, (2003) in rose. The number of branches could be because of inherent genetic factor that encourages the synthesis of substances such as auxin, cytokinin, and gibberellin at a faster rate. Similar differences in the number of branches were also noted by Wasnik *et al.*, (2016) in scented rose and Atram *et al.*, (2015) in hybrid tea rose.

Table 1. Vegetative Parameters variables of different miniature rose varieties

Variety	Plantheight (cm)	Plantspread(cm²)	Numberof branches
RedHot	14.7	18.2	6.5
MagicCarousel	22.3	27.2	7.8
Rosy	21.8	31.3	8.3
Calpoly	22.1	23.4	8.2
Beauty Secret	25.1	28.9	8.3
Hocky Pocky	21.4	22.3	8.6
Ruby Ruby	25.9	24.4	10.9
Rainbow'sEnd	19.8	25.1	9.9
JetTrail	13.9	15.9	6.3
Cricket	18.5	19.3	8.3
BlackJade	18.1	24.2	8.3
JeaneKennealy	22	31.4	8.1
F-TEST	S	S	S
SE(d)±	3.22	3.54	1.14
CD0.05	6.72	7.39	2.39
CV	19.27	17.85	16.92

~~Floral Generative variables~~ Parameters

The data for different floral parameters, including days to first flowering, bud diameter (cm), duration of flowering (days), and number of flowers per plant of miniature roses varieties are represented in Table 2.

Significantly ~~A significantly~~ lesser number of days for first flowering (32 days) was observed in variety ~~Red~~ Hot which was at par with variety Jeane Kennealy (33 days), Calpoly (38 days), Hockey Pocky (38 days), Black Jade (39 days) and Rainbow's End (39.3 days) whereas ~~more a greater~~ number of days for first flowering (44.3 days) was observed in variety Beauty Secret. Significantly bigger flower bud ~~diameter~~ diameters (1.1 cm) were observed in ~~the~~ variety Magic Carousel which was at par with variety Rosy (0.9 cm), Cricket (0.9 cm), and Rainbow's End (0.9 cm) whereas smaller flower ~~buds—diameter~~ bud diameters (0.5 cm) were observed in variety Jet Trail. Significantly longer flowering duration (164 days) was observed in ~~the~~ variety Ruby Ruby which was at par with ~~the~~ variety Beauty Secret (148 days) and Jeane Kennealy (147 days) whereas ~~a~~ shorter flowering duration (67 days) was observed in ~~the~~ variety Jet Trail. Significantly more ~~number of~~ flowers per plant (285.9) was observed in ~~the~~ variety Ruby Ruby which was followed by ~~the~~ variety Beauty Secret (241.2) whereas ~~the~~ lesser number of flowers per plant (52.6) was observed in ~~the~~ variety Jet Trail.

~~May—This~~ ~~maybe~~ due to the inherent ~~characters~~ characteristics, ~~the~~ genetic composition of the different varieties, and the ~~agro—climatic~~ agro-climatic conditions of the region. The varieties adapting well to the given agro-climatic conditions might have performed better while varieties struggling to adapt might have performed poorly. Variation in flowering parameters such as days taken to first flowering, bud diameter, flowering duration, and number of flowers per plant. Variation in days taken to first flowering was reported by Chandrashekar *et al.*, (1973) in hybrid tea ~~rose~~ roses. Variation in bud diameter was reported by Singh *et al.*, (1995) in rose. Variations in flowering duration were reported by Gaikwad and Patil, (2001) in ~~chrysanthemum—Chrysanthemum~~ and Pal *et al.*, (2003) in Gerbera. Variations in ~~the~~ number of flowers per plant ~~was—were~~ observed by Ramzan *et al.*, (2014) in *Rosa hybrida*. This might be due to ~~an~~ increase in morphological characteristics such as plant height, number of leaves, and area of leaves, which may have ~~initiate—initiated~~ the production of more photosynthates resulting in ~~a~~ greater accumulation of dry matter which in turn produces more number of flowers per plant.

Table 2. Floral Parameters Generative variables of different miniature rose varieties

Variety	Day to 1st flowering	Bud diameter (cm)	Duration of flowering (days)	Number of flowers per plant
RedHot	32	0.6	131	63.3
MagicCarousel	42	1.1	117	117.3
Rosy	43	0.9	114	97.2
Calpoly	38	0.8	112	142.7
Beauty Secret	44	0.6	148	241.2
Hocky Pocky	38	0.7	126	95.3
Ruby Ruby	41	0.7	164	285.9
Rainbow's End	39	0.9	108	144.1
JetTrail	42	0.5	67	52.6
Cricket	42	0.9	113	105.9
BlackJade	39	0.8	107	133.5
JeaneKennealy	33	0.7	147	177.8
F-TEST	S	S	S	S
SE(d)±	3.48	0.09	13.92	10.24
CD0.05	7.26	0.19	29.05	21.38
CV	10.78	13.71	14.08	9.09

4. Conclusions

It is concluded from the present investigation that 12 miniature rose varieties under study showed significant variation in all the parameters studied. Variety Ruby Ruby showed significantly better performance in vegetative parameters including plant height, plant spread, number of branches per plant, and flowering parameters including longer flowering duration, and more number of flower flowers per plant.

Variety Jeane Kenneal showed significantly better performance in vegetative parameter plants spread and flowering parameters including first flowering and longer flowering duration. Also Variety Beauty Secret performed better in vegetative parameters including plant height, plant spread, and flowering parameters like longer flowering duration and more number of flowers per plant. Variety Red Hot performed better in daystaken to first flowering. Bigger flower buds were observed in a variety of Magic Carousel.

References

Atram, V.R., Panchabhai, D.M. and Patil, S. (2015). Evaluation of hybrid tea rose varieties for flower quality and yield contributing traits. *Journal Agriculture Research*, **52**(4): 56- 60.

Chandrashekaraiah, T.S. (1973). Studies on evaluation of the hybrid tea roses for cut flowers. *M.Sc. (Agriculture) thesis* submitted to UAS, Bangalore.

Dias, S.M.F. and Patil, A.A. (2003). Performance of elite rose varieties at different population levels under the transitional tract of northern Karnataka. *Karnataka Journal of Agricultural Sciences*, **16**(2): 271-275.

Gaikwad, A.M. and Patil, S.S.D. (2001). Evaluation of chrysanthemum varieties under open and polyhouse conditions. *Journal of Ornamental Horticulture*, **4**(2):95-97.

Mohanty, C.R., Mohanty, A., Das, A.B. and Kae, S.D. (2011). Comparative performance of some rose varieties under an open and protected environment. *The Asian Journal of Horticulture*, **6**: 288-293.

Pal, K.K., Kumar, S., Srivastava, R. and Chandra, R. (2003). Evaluation of gerbera (*Gerbera Jamesonii*) cultivars under tara condition. *Journal of Ornamental Horticulture*, **6**(3):252-255.

Rajesh, M. and Ramesh, K.R. (1999). Effect of pruning height, shading and polythene covering

Comment [am5]: Put more recent primary publications of miniature roses. A journal of research articles commonly mentions the minimum number of recent references to be cited. For instance, publications in the last 10 years.

ongrowthandflowerproductionofRosecv.Raktagandha.*JournalofOrnamentalHorticulture*, **3**: 94-99.

Ramzan, A., Hanif, M. and Tariq, S. (2014). Performance of *Rosa hybrida* cultivars under agroclimatic conditions of Islamabad. *Pakistan. Journal of Horticulture*, **755**:499- 504.

Sharma, A. and Sharma, S.K. (2003). Scoring technique for rose evaluation. *Journal of Ornamental Horticulture*, **6(1)**: 50-54.

Sharova, N.L., Rybak, Y.G. and Marins, W.E. (1977). Development of gladiolus under the influence of micronutrients. *Russian Academy of Sciences*, **6(55)**: 50-54.

Singh,D.(1995).Improvementofrosessthroughcontrolledandopen-pollinatedprogenies.

Singh,P.,Dubey,R.K.,Singh,R.andKumar,R.(2013).Evaluationoffloribundarose(*Rosa hybrida*L.)cultivarsforlandscapeuseunderPunjabcondition. *JHort.Science*,**8(2)**:271- 275.

Soujanya,P.,Kulkarni,S.B.,Kumar,R.,Munikrishnappa,P.M.,Shivapriya,M.andM, H. (2018). Evaluation of rose (*Rosa hybrida* L.) varieties for growth, yield and quality undereasterndryzoneofKarnataka.*JournalofPharmacognosyandPhytochemistry*,**7(5)**: 165-168.

Wasnik,P.,Raut,V.U.,Bhande,M.H.andShambhavkar,V.B.,(2015).Evaluationofscented rose varieties for yield and quality of flowers. *Plant Archives*, **15(2)**: 895-897.