

Effect of Biocapsule and Nanozeolite in improving Growth, Development and Yield of Low chilling Apple (*Malus x domestica* Borkh.) cv. HRMN-99 in Prayagraj-agro-climatic Condition.

ABSTRACT

The present investigation was conducted at the research farm of Department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj on 4 years old plant during Dec 2023- May 2024 with the objective to find suitable treatment of Biocapsule and Nanozeolite with combination of NPK in relation to plant growth, development and yield of apple. The statistical design adopted for the experiment was randomized block design (RBD) having 03 replications and 09 treatment combinations. viz, T₀ (Control), T₁ Biocapsule 600ppm, T₂ Nanozeolite 300ppm, T₃ Biocapsule 600ppm + Nanozeolite 300ppm, T₄ Biocapsule 300ppm + Nanozeolite 300ppm, T₅ NPK+ Biocapsule 600ppm, T₆ NPK+ Nanozeolite 300ppm, T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm, T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm. The Treatments were applied by soil drenching at initiation of experiment, pre-flowering stage and the last one at the fruit setting stage to assess the effect on growth, development and yield of apple. The result of present investigation revealed that, the treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) was found significantly superior compared to other treatment combinations, which recorded highest mean value of plant height (3.41 m), primary branches (4.87), secondary branches (37.88), stem diameter (25.10 cm), leaf area (38.75 cm²), fruit set (36.12 %), number of fruits per plant (92.3) and average fruit weight (47.83 g). The highest yield per plant (9.84 kg) and yield per hectare (9.23 t/ha) were also obtained from treatment NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) and lowest yield was obtained from T₀ (control). Biocapsule and Nanozeolite with combination of NPK played an important role in fixing the nutrients in soil and protecting the apple plant against pathogen effect. That's why a positive response has to be seen in particular treatment during the research trail.

Keywords: biocapsule, nanozeolite, low-chilling apple, subtropical region, HRMN-99

INTRODUCTION

Apple (*Malus x domestica* Borkh.) is one of the most important temperate fruit crops of Northwestern Himalayan region in India. It belongs to family Rosaceae. It is liked throughout the world by all the people due to its pleasant taste and nutritional value and of its excellent taste and nutritious content; it is popular all over the world. Apple is one of the most popular fruits grown in temperate climates around the world. Its attractive look, crispy flesh, agreeable flavor, and sweet taste entice customers and bring in premium prices (Ali *et al.*, 2004).^[1] Apples are a fantastic source of fiber, healthy vitamins, and minerals. Apples are part in all food diets and its therapeutic value is well known for different illness (determines the absorption of gastric secretions, the elimination of toxins and has diuretic effect). Organic acids are an important component of fruit flavour and malic acid is the predominant organic acid in apple fruits (Campeanu *et al.*, 2009)^[2]. Malic acid is the major component of apple

that is found to maintain liver in a healthy condition and it helps in digestion process (Suni *et al.*, 2000) ^[3].

In the past it was generally regarded as a crop of the temperate zones but is increasingly cultivated under sub-tropical and even tropical conditions (Luck will 1984) ^[4]. The potential for low-chilling apple production in India's subtropical and tropical climates will improve if apple can be grown effectively in India's plains, where the climate is primarily subtropical. The choice of adaptable cultivars, as well as developments in technology and science, have made this possible ^[5].

HRMN-99 it is a low chilling apple variety noticed the resistance of an apple plant to climatic changes in 1999 by Hariman Sharma. This new variety is resistant to scab disease. Due to higher sugar acid ratio, higher total soluble solids, attractive colour and higher self-life this variety overcomes all most all the demerits of low chilling apple variety ^[6].

Recently, IISR-ICAR (Indian Council of Agricultural Research) scientists have developed the technology to pack bio-fertilizers in tiny capsules. This eliminates the need for farmers to carry the sacks of biofertilizers. It consists of a carrier medium rich in live microorganisms. When applied to seed, soil, or living plants, it increases soil nutrients or makes them biologically available. The capsule contains the microorganism in an immobilized/inactive condition and the cells can be activated by dissolving the capsule in water. This suspension can be diluted and the seeds or seedlings or rhizomes are soaked in the suspension for 30 minutes before sowing/transplanting into the main field. ^[7]

Nanozeolites are silicate minerals that have pores and channels in their crystal structure. It has a high affinity for cations like Na^+ , K^+ , and Ca^{2+} due to its unique increased Cation exchange capacity (CEC). In the soil system, zeolites are responsible for the selective retention of NH_4^+ and K^+ ions.

METHODS AND MATERIALS

The present investigation was conducted at Research farm of Department of Horticulture, SHUATS, Prayagraj, on 4- years-old plant during Dec 2023 to May 2024. The experiment was conducted in randomized block design with three replications and nine treatments T_0 (Control), T_1 Biocapsule 600ppm, T_2 Nanozeolite 300ppm, T_3 Biocapsule 600ppm + Nanozeolite 300ppm, T_4 Biocapsule 300ppm + Nanozeolite 300ppm, T_5 NPK+ Biocapsule 600ppm, T_6 NPK+ Nanozeolite 300ppm, T_7 NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm, T_8 NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm the treatments were applied by soil drenching and data were analyzed statistically.

Data was analyzed using OPSTAT.

Preparation of Biocapsule Solution.

Biocapsules were purchased from the ICAR licensed company CODAGU Agritech, Karnataka. Each biocapsule having 1gm wt. or 1000 ppm concentration and the ability of dissolving in 100 lt of water. So, according to concentration and the ability of dissolving in 100 lt of luke warm water before 12hrs of application and then after that 1 lt of solution was dissolved in 60 lt (for 600ppm) and 30 lt (for 300ppm) of water. After that apple plants were drenched with 1 lt of that solution. The biocapsule drenching were applied at different interval during the research.

Preparation of Nanozeolite Solution.

Nanozeolite were purchased from the Nano research element Haryana. According to their concentration of ppm they are first dissolve in 5ml ethyl alcohol solution after that it dissolve

in the 1 lt of water and then drenched in the apple plant. The nanozeolite drenching were applied at different interval during the research.

RESULT AND DISCUSSION

1. Growth Parameters.

The data pertaining to effect of Biocapsule and Nanozeolite on growth parameters of Apple are presented in Table 1. In case of plant height, the maximum plant height (3.41m) was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum plant height (2.69m) was found in treatment T₀(control). Increment in plant height due to the unique properties of zeolites, which affect plant cell division as well as growth. These findings are in accordance with **Bandana and Chandel, Vats et al., (2022), Jangid et al., (2023)**.

The maximum plant spread (227.25) cm was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum plant spread (184.58) cm was found in treatment T₀(control). Increment in plant spread due to the unique properties of zeolites, which affect plant cell division as well as growth. Similar findings were reported by **H. Singh and A. Mirza (2020), Deviet al.,(2022), Jangid et al., (2023)**.

The maximum No. of primary branches (4.87) was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum No. of primary branches (3.54) was found in treatment T₀(control). This might be due to rapid increase in cell division and cell elongation. Similar findings were reported by **Padhan et al., (2019), Vats et al.,(2022), Jangid et al., (2023)**.

The maximum No. of secondary branches (37.88) was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum No. of secondary branches (20.58) was found in treatment T₀(control). This might be due to rapid increase in cell division and cell elongation. The present findings are in close agreement with the finding of **Padhan et al., (2019), Vats et al.,(2022), Jangid et al., (2023)**.

In terms of stem diameter, the maximum stem diameter (25.10) cm was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum stem diameter (18.83) cm was found in treatment T₀(control). Increment in stem diameter due to different treatments of biocapsule and nanozeolite. This might be due to the unique properties of zeolites enable the prolonged and controlled introduction of necessary nutrients such as potassium, ammonium, and phosphates into the soil, which affect plant cell division as well as growth. Similar results were achieved by **Padhan et al., (2019), Vats et al.,(2022), Jangid et al., (2023)**.

2. Flowering and Fruiting Parameters.

The data pertaining to effect of Biocapsule and Nanozeolite on growth parameters of Apple are presented in Table 2.

In terms of number of flowers, the maximum No. of flowers (255.49) was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum No. of flowers (187.44) was found in treatment T₀(control). after 4 years of transplanting. This might be due to the different treatments of biocapsule and nanozeolite with combination of NPK enhanced the reproductive growth. The present findings are in close agreement with the finding of **Sharma et al., (2005), Padhan et al., (2019), Vats et al., (2022)**.

In terms of number of fruits per plant, the maximum No. of fruits (255.49) was found in treatment T₇NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) Whereas the minimum No. of fruits (187.44) was found in treatment T₀ (control) after 4 years of transplanting. This might be due to the different treatments of biocapsule and nanozeolite with combination of NPK enhanced the reproductive growth. These findings are in accordance with **Padhan et al., (2019), Vats et al., (2022), Jangid et al., (2023)**.

Coming to the fruit weight, the maximum fruit weight (255.49) g was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) (247.98) g Whereas the minimum fruit weight (187.44) g was found in treatment T₀(control) after 4 years of transplanting. This might be due to the different treatments of biocapsule and nanozeolite with combination of NPK enhanced the reproductive growth. Similar findings were reported by **Verma et al., (2009), Singh et al., (2011), Vats et al., (2022)**.

The maximum fruit yield (kg/plant) (9.83) kg was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) (9.53) kg Whereas the minimum fruit yield (4.16) kg was found in treatment T₀(control) after 4 years of transplanting. Similar findings were reported by **Padhan et al., (2019), Vats et al., (2022), Jangid et al., (2023)**.

The maximum fruit yield (t/ha) (9.23) tonnes was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) (9.05) tonns Whereas the minimum fruit yield (3.19) tonns was found in treatment T₀(control) after 4 years of transplanting. Similar findings were reported by **Padhan et al., (2019), Vats et al., (2022), Jangid et al., (2023)**.

3. Fruit Quality Parameters.

The data pertaining to effect of Biocapsule and Nanozeolite on growth parameters of Apple are presented in Table 3. The maximum T.S.S (11.63) brix was found in treatment T₇ NPK+

Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by treatment T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) (11.16) brix Whereas the minimum T.S.S (8.08) brix was found in treatment T₀(control). These results are in support with **Singh et al., (2011), Vats et al., (2022), Jangid et al., (2023)**.

In case of acidity, the highest acidity (0.74) % was found in treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) (0.72) %Whereas the lowest acidity (0.49) % was found in treatment T₀(control). Similar findings were reported by **H.A. Hifny et al., (2017), Vats et al., (2022), Jangid et al., (2023)**.

Coming to T.S.S: Acid ratio, the highest ratio (18.77) was found in treatment T₁Biocapsule 600ppm (Soil drenching) followed by treatment T₄Biocapsule 300ppm +Nanozeolite 300ppm (Soil drenching) (17.00)are found statistically at par to Biocapsule 600ppm (Soil drenching)Whereas the minimum T.S.S:Acid ratio (15.55) was found in treatment T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching). These findings are in accordance with **Singhet al., (2011), Vats et al., (2022), Jangid et al., (2023)**.

UNDER PEER REVIEW

Table 1: Growth Parameters.

Symbol	Treatment	Growth Parameters				
		Plant height (m)	Plant Spread (cm)	Primary Branches	Secondary Branches	Stem Diameter (cm)
T0	Control	2.69	184.58	3.54	20.58	18.83
T1	Biocapsule 600ppm (Soil drenching)	2.84	196.8	3.76	24.94	19.36
T2	Nanozeolite 300ppm (Soil drenching)	2.75	191.36	3.6	23.46	19.78
T3	Biocapsule 600ppm + Nanozeolite 300ppm (Soil drenching)	3.16	219.69	4.72	33.88	24.35
T4	Biocapsule 300ppm + Nanozeolite 300ppm (Soil drenching)	2.87	198.03	3.85	27.82	21.52
T5	NPK+ Biocapsule 600ppm (Soil drenching)	3.05	208.65	4.59	31.12	23.37
T6	NPK+ Nanozeolite 300ppm (Soil drenching)	2.98	201.26	4.23	28.22	22.87
T7	NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching)	3.41	227.25	4.87	37.88	25.1
T8	NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching)	3.31	225.38	4.8	35.97	24.84
	F Test	S	S	S	S	S
	SE(d)±	0.086	0.099	0.109	0.3	0.109
	CD0.05	0.184	0.212	0.232	0.642	0.233
	CV	3.507	0.059	3.155	1.253	0.602

Table 2: Flowering and Fruiting Parameters.

Symbol	Treatment	Flowering & Fruiting Parameters				
		Number of Flowers/Plant	Number of Fruits/Plant	Fruit Weight (g)	Fruit Yield (kg/plant)	Fruit Yield (t/ha)
T0	Control	187.44	44.75	28.92	4.16	3.19
T1	Biocapsule 600ppm (Soil drenching)	204.62	68.71	31.38	6.42	5.41
T2	Nanozeolite 300ppm (Soil drenching)	199.76	60.82	30.3	5.93	5.11
T3	Biocapsule 600ppm + Nanozeolite 300ppm (Soil drenching)	236.49	84.37	44.9	8.69	8.87
T4	Biocapsule 300ppm + Nanozeolite 300ppm (Soil drenching)	218.79	73.47	32.7	6.82	6.77
T5	NPK+ Biocapsule 600ppm (Soil drenching)	229.51	79.98	40.29	7.45	8.15
T6	NPK+ Nanozeolite 300ppm (Soil drenching)	221.63	77.86	37.18	7.11	7.64
T7	NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching)	255.49	92.3	47.83	9.84	9.23
T8	NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching)	247.98	87.85	45.2	9.53	9.05
	F Test	S	S	S	S	S
	SE(d)±	0.214	0.189	0.468	0.082	0.43
	CD0.05	0.458	0.404	1.001	0.176	0.92
	CV	0.118	0.311	1.524	1.376	7.476

Table 3: Fruit Quality Parameters.

Symbol	Treatment	Fruit Quality Parameters		
		T.S.S °Brix	Acidity %	T.S.S: Acid
T0	Control	8.08	0.49	16.38
T1	Biocapsule 600ppm (Soil drenching)	9.63	0.51	18.77
T2	Nanozeolite 300ppm (Soil drenching)	9.35	0.56	16.69
T3	Biocapsule 600ppm + Nanozeolite 300ppm (Soil drenching)	10.85	0.68	15.87
T4	Biocapsule 300ppm + Nanozeolite 300ppm (Soil drenching)	9.86	0.58	17.00
T5	NPK+ Biocapsule 600ppm (Soil drenching)	10.68	0.65	16.35
T6	NPK+ Nanozeolite 300ppm (Soil drenching)	10.01	0.61	16.4
T7	NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching)	11.63	0.74	15.65
T8	NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching)	11.16	0.72	15.55
	F Test	S	S	S
	SE(d)±	0.043	0.014	0.068
	CD0.05	0.093	0.03	0.146
	CV	0.523	2.829	0.507

CONCLUSION

On the basis of present investigation, it is concluded that the various treatments of Biocapsule and Nanozeolite has been applied to enhance the vegetative and reproductive growth of apple cv. HRMN-99, where treatment T₇ NPK+ Biocapsule 600ppm+ Nanozeolite 300ppm (Soil drenching) is found superior among others, followed by T₈ NPK+ Biocapsule 300ppm+ Nanozeolite 300ppm (Soil drenching) and the lowest was T₀ (control). In every aspect of growth, development and yield parameters.

REFERENCES

- Ali, Raza H, Khan, Muhammad Azam, Hussain Maaz. Effect of different periods of ambient Storage on chemical composition of apple fruit. *International Journal of Agriculture and Biology*. 2004;06(3):568– 571.
- Campeanu G, Neeta G, Darjanschi G. Chemical composition of the fruits of several apple cultivars growth as biological crop. *NotulaeBotanicae Horticulture Agrobotanici Cluj Napoea*, 2009; 37(2):161-164. 3. Considine M. F.
- Suni M, Nyman M, Errikson NA, Bjork I, Bjork I. Carbohydrate composition and content of organic acids in fresh and stored apples. *Journal of Science of Food and Agriculture*. 2000; 80:1538-1544.
- Luckwill LC. Apple growing around the World, Problems and Prospects. In *Apples and pears: report of the Royal Horticultural Society Conference, 1983*, (ed Elspeth Napier) The Royal Horticultural Society, London, 1984, 1-10.
- Padhan A, Mishra S, Bahadur V. (2019). Effect of growing media on growth, development and establishment of low chilling variety of apple HRMN-99 under Prayagrajagro climatic conditions. *Journal of Pharmacognosy and Phytochemistry*. 8(3):1227-1230. National innovation foundation. 9th national biennial grassroots innovation and outstanding traditional knowledge award; 2017.
- ICAR-Indian Institute of Spices Research, Kozhikode.
Avaialble:<https://icar.org.in/content/novelmethod-storing-and-pgprmicrobes-throughbiocapsules>.
- Bandana, Chandel JS. Effect of growth regulators, PGPR and nitrogen on growth of apple nursery plants under protected conditions. *Journal of Hill Agriculture*. 2017;8(2):176-180.
- Saurav Vats, Saket Mishra, Vijay Bahadur and Ravi Kumar Singh. (2022). The Role of Nanozeolite and Biocapsule in Improving Growth, Yield and Fruit Quality of Low

Chilling Apple (*Malus × domestica* Borkh.) Cv. HRMN-99 in Prayagraj Agro-Climatic Condition. *International Journal of Environment and Climate Change* 12(11): 866-875; Article no. IJECC.89430 ISSN: 2581-8627.

Muskan Jangid and Saket Mishra. (2023). Effect of plant growth regulators on flowering and fruit setting of low chilling apple (*Malus domestica*) cv. HRMN-99 under Prayagraj agro climatic conditions. *The Pharma Innovation Journal*;12(5): 2335-2339.

Harinder Singh, Anis Mirza. Effect of plant growth regulators on growth, yield and quality of papaya (*Carica papaya* L.) Cv. Taiwan red lady, *International Journal of Chemical Studies*. 2020;8(6):2168-2172.

Yadav SK, Jadhav SM. Deep convolutional neural network based medical image classification for disease diagnosis. *Journal of Big data*. 2019 Dec;6(1):1-8.

Sharma SD, Sharma N, Sharma CL, Sood R, Singh RP. Studies on correlations between endomycorrhizal and *Azotobacter* population with growth, yield and soil nutrient status of apple orchards in Himanchal Pradesh. *Acta Horticulture*. 2005; 696:283-287.

Verma ML, Singh C, Bhardwaj SP. Effects of biofertilizers on soil moisture, nutrient status and fruit productivity under organic cultivation of apple in Himachal Pradesh. *Indian Journal of Soil Conservation*. 2009;37(3):201-205.

Singh SR, Zargar MY, Najjar GR, Peer FA, Ishaq MI. Integrated Use of Organic and Inorganic Fertilizers with Bioinoculants on Yield, Soil Fertility and Quality of Apple. *Journal of the Indian Society of Soil Science*. 2011;59(4):362- 367.

Hifny HA, Khalifa SM, Hamdy AE, Abd El-Wahed AN. Effect of GA₃ and NAA on growth, yield and fruit quality of Washington navel orange, *Egypt Journal Horticulture*. 2017;44(1):33-43.

Devi, Y. Pabitri, and Samir E. Topno. 2022. "Effects of Nano Chitosan and Bio Capsule on Growth, Yield and Quality of Chilli (*Capsicum Annuum* L.) in Poly-House Condition". *International Journal of Plant & Soil Science* 34 (23):662-70. <https://doi.org/10.9734/ijpss/2022/v34i232473>.