

Original Research Article

Managing High Anisometropia – The Blended Refractive surgery way.

Abstract:

Purpose: To compare patient-reported outcomes (PROs) and satisfaction results after blended refractive surgery with one eye Lasik laser/ PRK and another eye ICL..

Patients and methods: A Quality of life and Patient reported spectacle independence (PRISQ) questionnaire was administered to patients who had undergone blended Refractive surgery at 2 months of follow-up

Results: Overall PROs and satisfaction was similar among all the groups. Refractive outcomes and accuracy were similar among the groups, Refractive outcome differences were not meaningful among the groups and were not a differentiating factor in PROs. Substantially fewer patients in the third group noticed glare and halo compared with the 3.0/3.0 group ($P < 0.0001$, chi-square test). No new safety concerns were reported.

Conclusion: The choice of blended refractive surgery resulted in a higher percentage of patients who were deemed unfit for LVC being satisfied with Near, intermediate, and distance vision, overall the fewer PRO differences were not statistically significant.

Keywords: Implantable collamer lens, Lasik laser , photo refractive keratectomy , spectacle independence, glare, haloes.

Introduction

‘Refractive surgery’ encompasses any procedure that corrects refractive error .It is well recognized that refractive surgery has significant impact on quality of life and daily work, with benefits extending beyond spectacle independence.

Today, Refractive surgery has evolved beyond the stereotypical ‘laser eye surgery’. With the Advent of technologies like Femto lasik / SMILE ,ICL/IPCL, Customised RLE wide range of refractive errors are being corrected leading to rise in demand of the refractive services.

Inspite of all advancements a good percentage of patients present with difference of high refractive error in both eyes i.e Anisometropia leading to them being deemed unfit for a single refractive procedure.High-myopia anisometropia patients may have difficulty with fusion when wearing spectacles

A common practice followed by many centres is laser vision correction for the fit eye and contact lens for the eye unfit for lasik / PRK. Which surgical procedure will be chosen for the eyes of anisometropia patients mainly depends on the level of refractive error.

Due to the limits of patients' corneal conditions, LVC has been generally chosen for the correction of <12 D of myopia, and the outcomes are poor in high-myopia cases compared to cases of low to moderate myopia. While ICL, a posterior-chamber phakic intraocular lens, is the preferred technique for high myopia of >12 D.

Previous studies (Sanders & Vukich [2003](#), [2006](#); Sanders [2007](#); Igarashi et al. [2009](#); Kamiya et al. [2012](#)) have reported the grouping comparisons of ICL and laser-assisted *in situ* keratomileusis (LASIK)

The current study probably the first in india was objectively conducted to compare the patient reported outcomes of ICL implantation in one eye and LVC in the contralateral eye and investigate the long-term visual quality and stability.

Patients and methods:

The present study is conducted in the Department of Ophthalmology. Ethics reviews and approvals were conducted and obtained by the institutional review boards. : A Total of 50 subjects who underwent ICL (phakicIOL) in one eye and lasik laser in other or ICL in one eye and PRK in other were reviewed.

A Quality of life and Patient reported spectacle independence (PRISQ) questionnaire was administered to patients who had undergone blended Refractive surgery and 2 months of follow-up

Results:

Overall PROs and satisfaction was similar among all the groups. Refractive outcomes and accuracy were similar among the groups, Refractive outcome differences were not meaningful among the groups and were not a differentiating factor in PROs. Substantially fewer patients in the third group noticed glare and halo compared with the 3.0/3.0 group ($P < 0.0001$, chi-square test). No new safety concerns were reported.

Complete satisfaction level was noted in "Seeing in the distance", "Recognising faces across the street", Watching television, Driving a car by night, Using steps, Crossing the road, Using public transport and Employment / Housework activities. While in some cases satisfaction category "Seeing in the bright light / glare", "Seeing in the poor light / dim", "Appreciating colors", "Driving a car by day" (not at all 18, 72%) and (a little 7, 28%).

Function comfortably without glasses or contacts was noted in

Complete satisfaction with vision without glasses or contacts was observed.

Table 1 : Overall PROs and satisfaction

	Not at all (1)	A little (2)	Quite a bit (3)	A lot (4)
Reading	38 (67.9%)	12 (21.4%)		
Seeing in the distance	50 (100%)			
Recognising faces across the street	50 (100%)			
Watching television	50 (100%)			
Seeing in the distance	50 (100%)			
Seeing in the bright light / glare	36(72%)	14 (28%)		
Seeing in the poor light / dim	36 (72%)	14 (28%)		
Appreciating colors	36 (72%)	14(28%)		
Driving a car by day	38 (76%)	12 (24%)		
Driving a car by night	50 (100%)			
Using steps	50 (100%)			
Crossing the road	50 (100%)			
Using public transport	50 (100%)			
Travelling independently	36 (72%)	14 (28%)		
Moving in unfamiliar surrounding	36 (72%)	14 (28%)		
Employment / Housework activities	50 (100%)			

Spectacle independence with blended refractive surgery was observed in 38 (76%) of cases while 12 (24%) still need glasses.

Table 2 : Frequency Table

Need glasses	Frequency	Percentage
Yes	12	24%
No	38	76%

Table 3 : Function comfortably without glasses or contacts

	All the time	Most of the time	Some of the time	A little of the time	None of the time
Wear Glasses or contacts for					50(100%)
Function comfortably without glasses or contacts	26(52%)	24(48%)			
Satisfaction with vision without glasses or contacts	50(100%)				

Discussion:

- ❖ The different surgical methods for the correction of myopia among anisometropia patients are based on the principles of safety and efficacy.
- ❖ The long-term visual quality and stability of different surgical methods should be evaluated.
- ❖ In this study we presented patient reported outcomes and satisfaction results after hybrid refractive surgery.
- ❖ No infection, corneal dilation or other adverse reactions were observed, suggesting that both ICL implantation and corneal LVC are safe and reliable in cases with suitable indications

- ❖ The visual image magnification rate and visual quality were improved after blended refractive surgery
- ❖ A study by Nicolas et al reports better quality of vision, stable refraction and high satisfaction score after 9 years in a patient treated for high myopia with one eye ICL and other eye LASIK. (1)
- ❖ Another study by Roberto Zalvidar et al reported better visual quality gains, short term safety and refractive stability .(2)
- ❖ A study by Xun chen et al also demonstrated that ICL implantation and LRS techniques are both safe and effective procedures for myopia with suitable indications.(3)

Conclusion:

The choice of blended refractive surgery resulted in a higher percentage of patients who were deemed unfit for LVC being satisfied with Near, intermediate, and distance vision, overall the fewer PRO differences were not statistically significant.

References:

1. [Nikolaos S Tsiklis, MD, MSc, George D Kymionis, MD, PhD, Carol L Karp, MD, Tatianna Naoumidi, MD, and Aristofanis I Pallikaris, PhD](#) Journal of Refractive Surgery, 2013;23(9):935–937Published Online:November 01, 2007<https://doi.org/10.3928/1081-597X-20071101-12>
2. [Roberto Zaldivar, MD, Jonathan M Davidorf, MD, Susana Oscherow, MD, Giselle Ricur, MD, and Virginia Piezzi, MD](#) Journal of Refractive Surgery, 2013;15(3):299–308 Published Online:May 01, 1999<https://doi.org/10.3928/1081-597X-19990501-04>
3. Chen X, Guo L, Han T, Wu L, Wang X, Zhou X. Contralateral eye comparison of the long-term visual quality and stability between implantable collamer lens and laser refractive surgery for myopia. *Acta Ophthalmol.* 2019 May;97(3):e471-e478. doi: 10.1111/aos.13846. Epub 2018 Sep 5. PMID: 30187653; PMCID: PMC6585688.
4. Kim TI, Alio Del Barrio JL, Wilkins M, Cochener B, Ang M. Refractive surgery. *Lancet.* 2019;393:2085–98.
5. Sugar A, Hood CT, Mian SI. Patient-reported outcomes following LASIK: quality of Life in the PROWL Studies. *JAMA.* 2017;317:204–5.

6. Sugar A, Rapuano CJ, Culbertson WW, Huang D, Varley GA, Agapitos PJ, et al. Laser in situ keratomileusis for myopia and astigmatism: safety and efficacy: a report by the American Academy of Ophthalmology. *Ophthalmology*. 2002;109:175–87.
7. Sandoval HP, Donnenfeld ED, Kohnen T, Lindstrom RL, Potvin R, Tremblay DM, et al. Modern laser in situ keratomileusis outcomes. *J Cataract Refract Surg*. 2016;42:1224–34.
8. Ang M, Mehta JS, Chan C, Htoon HM, Koh JC, Tan DT. Refractive lenticule extraction: transition and comparison of 3 surgical techniques. *J Cataract Refract Surg*. 2014;40:1415–24.
9. Reinstein DZ, Archer TJ, Gobbe M. Laser blended vision for presbyopia correction. In: Pallikaris IG, Plainis S, Charman WN, editors. *Presbyopia: origins, effects and treatment*. Thorofare, NJ: SLACK Incorporated; 2012. [
10. Evans BJ. Monovision: a review. *Ophthalmic Physiol Opt*. 2007;27:417–39.