

## Original Research Article

### **A Study on Fishing Gear and Methods commonly used at an Oxbow lake of South-West Bangladesh**

#### **Abstract:**

A total of 18 fishing gears and crafts were recorded at Bukbhora ox-bow Lake of Jashore region. Fishing gears were categorized as nets (8), traps (3), hooks and lines (3), wounding gears (2) and crafts (2). Among nets, highest mesh size recorded 5-8 cm in fash jal whereas the lowest was thela jal 0.25 cm. The highest fishing effort/day was found for Jhaki Jal (cast net) 13-15 and the lowest was recorded in case of current and punti Jal (01). Highest CPUE (kg/effort) was found in ber jal (15-18) and the lowest was found in Thela jal (1-1.5). Similarly seine net contributed to the highest catch (70.19%) and push net was the lowest (1.00 %). Maximum fishers had pieces of kochal (Seine net) and komar (drag net) 2-5 pieces and those were jointed together (each of 18 fishers groups separately work) during the harvesting period. Fishing (harvesting of carp species) started in November-December and completed on 30th June in each season. Based on the usage, the seine net was found as the most usable gear by the fishermen constituting 78.15% and fewer amount by traps. The highest price of fishing gear was observed in case of ber jal (about 1,00,000 BDT) and the lowest price was for punti jal (250-2500 BDT). Among 16 fishing gears 5 of these were found illegal in this study due to its mesh size and destructive behavior of fish biodiversity. Except some, majority fish species trapped in the recorded fishing traps were **SIS**, whereas fishing hooks and lines were used to catch carnivorous fish species mainly. Wounding fishing gears were used for wounding large size fishes during heavy flood or when water level becomes low. This study showed that 40.91% fishers had their own fishing craft (dingi nouka) where 59.09% had no craft. It is recommended to strong implementation of all conservation

**Comment [A1]:** At the beginning of the paragraph it is best to briefly state the problem of this research

**Comment [A2]:** It's best to write down the abbreviation of SIS in full

guidelines and acts to prevent illegal fishing, seizing of destructive fishing gears and also protecting potential breeding habitats.

**Key words:** CPUE, Fishing gears, Mesh size, SIS, Conservation

**Introduction:**

Fish and fisheries have been an integral part of life of the people of Bangladesh from time immemorial; many aspects of the culture, economy and tradition are based around fisheries activities. Bangladesh has the third biggest aquatic fish bio-diversity blessed with enormous and exuberant fisheries resources in Asia, after China and India, with about 800 species in fresh, brackish and marine waters. Bangladesh is endowed with an extensive network of inland waterways, including estuaries, small and large rivers, *beels*, *haors*, *baors*, canals, lakes, and seasonal channels (Ali et al., 2017). The *baor* fishery is one of the most popular fisheries resources due to its steadily rising production, which is expected to increase from 4698 MT to 11685 MT (2006–2022) (DoF, 2023). This type of fishery requires very little or no feed at all as feed is the most expensive input in aquaculture (Begum et al., 2017). The primary sources of Bangladesh's constant freshwater supply are the ancient, winding bends in the river (Haque et al., 2015). The Baor, also known as the Oxbow Lakes, are semi-closed bodies of water in the southwest region of Bangladesh that are produced by dead arms of rivers in the Ganges delta (Farid, 2013). There are about 600 *baors* in the southwest part of Bangladesh comprising 5,671 ha and 29 *baors* are being managed under some sorts of culture practices initiated by the DoF and very few of the rest are under the management of fisher's cooperatives or NGOs (DoF, 2022; 2023).

The skill of fishing involves capturing not just fish but also other aquatic creatures (Sreekrishna et al., 2001). In addition to being a source of food for humans, fishing supports a substantial

portion of rural society's livelihood and nutritional security (Hameed et al., 2000). In these inland water bodies, prevalent fishing activities include hand picking, dewatering, overfishing, undersized fishing, haphazard fishing, fishing during the nesting season, and so on (Das et al., 2023). Moreover, the biodiversity of the freshwater fish species is declined due to the intense human intervention resulting in niche loss and declination and as a result, many fish species have become highly endangered (Rahman et al. 2012).

The choice of fishing gear, method of operation, number of fishermen, fishing intensity, and level of exploitation all has a significant impact on the future of the fishing industry (Mia et al 2017). Modernized crafts have advancement to capture and transport fishes as well (Rahman et al 2017) which influences the fish biodiversity greatly (Rahman et al 2016). Thus, the evaluation and information about fishing gears and crafts are necessary to emplace developmental measures about smooth **harvesting** among fisherman's community (Baleta et al 2017) from any area.

Numerous researches rose regionally to report the biodiversity of fishers and fishing gears (Azam et al 2014; Ali et al 2015; Rahman et al 2015; Rahman et al 2017) to understand the fishing status deeply.

Numerous groups of fishing crafts and gears are operated in *baor* for commercial fishing exploitation (Rahman et al 2016) having different shapes and sizes to capture various sizes of fish species (Azam et al 2014). The shape and size of the gear depends on the locality, types of water body, different operation area, depth of water and availability of target species to be caught (Flowra et al. 2013). The major categories of fishing gears that are regularly used in Bangladesh can be counted as the following: fishing nets, fishing traps, hooks and lines, wounding gears and fish aggregation device (Chakraborty et al 1995) where various types of materials are used to make these fishing gears (Hameed and Boophendranath, 2000).

**Comment [A3]:** It's best to find the right equivalent word that is generally applicable to make it more appropriate

According to Ghosh and Biswas (2017), the catch per unit effort (CPUE) is a recognized measure of the abundance and consumption of fisheries and can be used to calculate the number of fishing units in a sustainable fishery. It also measures stock density, physical and financial productivity, and the efficiency of a fishing operation. It is expected that CPUE will reflect the fraction of the fish population used as the relative abundance index (Karim et al 2019).

Since, *Baor* is one of the natural stocks of native fish species especially SIS in south western part of Bangladesh, the biodiversity of this *baors* could be negatively affected by using harmful fishing gear and fishing system. Many researchers have studied fishing gear in different water body *i.e* river (Saberin et al., 2018, Tikadar et al., 2021); *haor* (Mia et al., 2017, Yeasmin et al., 2021 )

However, the impact of fishing gear used in *baors* on biodiversity is largely understudied. The literature outlining the description, mode of operation, and classification of *baor* fishing gear is particularly illuminating. Such information is critical for developing effective management practices for the *Baor's* commercial fisheries. Unreliable research has been done on the diversity and productivity of fish in the *baor*. In this conception, the present research was taken to reveal some aspects including different types of fishing gears operated in *baors* with their mode of action, CPUE, catch composition of various gears, the significant reasons for the eradication of fish fauna and find out some recommendations to enhance the fish biodiversity of the *baors* in Bangladesh.

Comment [A4]: and conservation

## **MATERIALS AND METHODS**

### **Study area and duration**

Bukbhora Baor (23.1759° N, 89.1170° E) is one of the prominent oxbow lakes located in Jashore Sadar Upazila. The lake is linked to the Kopothakho River by a canal spreading around 7 kilometres (4.3 mi), known as Katakhal. The present study was conducted for a period of twelve months from July 2023 to June 2024. Data was collected personally through face to face interviews using well-structured questionnaire and cross-check.

**Comment [A5]:** Write down the frequency of data collection, whether once a month, twice a month or every week?

**Table 1: Sampling stations of the study**

Sampling station	Name
Site 1	Halsa
Site 2	Gohata
Site 3	Chanduita bazar
Site 4	Mathbari
Site 5	Ichhapur

**Catch per unit effort (CPUE) and gear efficiency:**

The catch per unit effort (CPUE) of the angling gears were taken dependent on the weight of fish discovered amid an angling day ( $\text{kg gear}^{-1}\text{haul}^{-1}$ ) for the various species consolidated and the gear efficiency ( $\text{kg gear}^{-1}\text{person}^{-1}\text{haul}^{-1}$ ) additionally assessed based on the weight of fish got and a number of individuals drew in with each fishing gear per hour.




Figure 01. Map showing the study area

### Collection of primary data

Data were gathered by direct catch assessment survey (CAS) and fishing effort survey (FES) conducted at the study sites. The fishing gears were overviewed by direct physical perceptions and dependent on participatory rural appraisal (PRA), questionnaire interview (QI), focus group discussion (FGD). Ten (10) FGD were made with 6–12 members where participants were fisherman of young, middle and old ages. The QIs and FGDs were led with a semi-organized and pre-trialed questionnaire. A total of thirty fishermen in the chosen *baor* area were surveyed using a basic random sampling procedure and questionnaire interviews in order to gather data. The interview was conducted with fisherman either at their homes or at their fishing locations. Data were collected from fishermen, aratdar, fish landing center, labor and direct observation of fishing gears at the time of operation and from the local fishing gear market. Data about fishing gear with their mesh size, operation mode, building materials, fish species caught, were the

**Comment [A6]:** need to provide an explanation of why young, middle and old respondents should be selected

principal and common type of net used in the season etc. and data about operation of fishing craft with their size, shape etc. were collected by Questionnaire interviews (QI).

### **Study on fishing gears**

Fishing gears operated at selected *baor* by fishermen were studied by direct physical observation. A measuring scale was used as a tool to measure mesh size of existing fishing gears. Direct observation was conducted to record data on fishing gear like shape and mesh size (cm) of fishing gears. Information about the price of fishing gear, species catch by individual gear, catch per unit effort (CPUE)/day and operation period of individual fishing gear were collected through interviewing with fishermen.

### **Secondary data**

After regular field study, the primary data were compared with the available secondary data to make a precise and composite picture of the present study area. Books, Journals, Upazila Fisheries Officer (UFO) etc. were used to modify, cross-check and accurate the information. The collected gears were assorted under various major classes followed by Ahmed (1971). Fish samples captured by different gears were identified and then taxonomically characteristics followed by Talwar and Jhingran (1991) and Rahman (2005).

### **Data processing and analysis**


All the data were accumulated to analyze the findings. Tabular technique was applied by using simple statistical tools such as percentages and averages. For processing and analyzing the data “Microsoft Excel-2013” was used.

## RESULTS

The fishery of the ox-bow lake is multispecies and multigear in nature. Fishermen use numerous types of fishing gears for fishing based on seasons and water depth of the *baor*. Variation in net types, lengths as well as mesh sizes of the net varying based on the fishermen choice, economic conditions of the persons involved in fishing practices and also the abundance of fish. Categorization of different fishing gear with their shape, respective mesh size, major species caught, price, catch per unit effort (CPUE) and operation period in the *baor* are showed in Table 1. About eight (08) various types of fishing gear were classified under six major classes and these were gill net (current jal, puti jal, fash jal), seine net (ber jal), Drag net (komar Jal), lift net (vashal jal), push net (thela jal), cast net (jhaki jal), three types of fish traps (Charo, bitte, ghuni), three types of hook and lines gears (chip borshi, dhawn borshi, nol borshi), two types of wounding gear (koach, teta) and two types of fishing crafts (kosha, dinghi) were found for catching fishes in the following *baor* (Table 2,3).

### Fishing gear efficiency and CPUE of gears in the Baor

In the present study the fishing gear efficiency using CPUE and fishing effort of the various fishing gears operated in the Bukbhora *baor* was recorded and data were presented in Table 1. Along with all types of gears most of the fishers use nets where about 78% fishermen use seine net and only 1% use push net. Among other gears types wounding gear, hooks and lines and traps occupy 0.5%, 0.35%, 0.12%, respectively (Figure 2).



**Figure 02:** Fishing gear used by fishermen in the *baor*


**Table 2:** Different fishing net with their mesh size, shape, price, and major species caught, CPUE and operating period in *Baor* at Jashore, Bangladesh

Type of fishing	Gear		Description/Size		Mesh size (cm)	Shape and Materials Used	Fishing effort/day	CPUE (kg gear <sup>-1</sup> haul <sup>-1</sup> )	Main Fishing period	Catch Composition	Construction cost (BDT/net)
	English name	Local name	Length (m)	Breadth (m)							
Kochal	Scine net	Ber jal	80-110	15-20	2.30-5.00	Rectangular, Nylon twine	02	15-18	Oct-Dec Feb-mar May-Jun	Rui, Catla, Silver carp, Mola, Tit puti	About 1,00,000
Komar	Drag net	Moi/Komar jal	300-400	9-10	3-4	Rectangular, Nylon twine	4-5	02-2.5		Bele, Taki, Puti, Tengra, baim	1,500-4,000
Push	Push net	Thela jal	-	-	0.25-0.50	Triangular, Cod twine with bamboo frame	7-8	01-1.5	Over the year	Puti, Taki, Bele, Chapil, Icha chingri	1,500-2,500
Cast	Cast net	Khepla/Jhaki jal	8-10	8-10 (round diameter)	0.5-1.4	Conical, Nylon twine	13-15	<1.50	July-september	Punti, chapila, coilisa, Silver carp, Rui, Catla, koi	2,000-4,500
Lift	Lift net	Khora/Veshal jal	12-15	10-12	0.5 Centre ; 1.5 Front	Square, Nylon twine	05	01-1.5	June-Aug & Oct-Dec	Bata, Punti, mola, tengra, Koi	9,000-10,000
Gill netting	Gill net	Current jal	50-150	1.0-1.5	1.0-3.00	Rectangular, Nylon twine	01	08-10	Round the year	Puti, Taki, shing, magur tengra, Bele, Koi	3,500-6,000
		Fash jal	10-12	2-5	5-8	Rectangular, Mono-filament nylon	02	10-12		Puti, Taki, Shing, Bele, Icha chingri	3,500-6,500
		Puti Jal	20-30	0.5-1.0	2.0-3.0	Rectangular, cotton twine or monofilament	01	04-05		Punti, Bele, Gulsha, Koi, Taki	250-2500

\*Jal= Fishing net; BDT= Bangladeshi Taka (currency); CPUE= Catch Per Unit Effort.

**Gear wise catch composition:**

It was observed that huge number of fishes was caught by seine net (Ber jal) and the rare species were damaged through these bulk catches and it is considered as one of the most detrimental gears compared to others. Among these seine net contributed to the highest catch (70.19%) and push net was the lowest (1.00 %).



**Figure 03.** Contribution of gears to catch in *baor*

**Table 03.** Total catch and contribution of gears to catch of *baor* (Nov-Jun)

Gear Type	Total Catch/Day (kg)	Probable fishing time (Days)	Total Catch (8 Months) (Kg)	Total catch (%)
Seine Net	1797	90	161735	70.19
Drag Net	505		45417	19.71
Gill Net	163		14678	6.37
Cast net	70		6290	2.73
Push net	26		2304	1.00
<b>Total</b>			<b>230424</b>	<b>100</b>

**Catch per Unit Effort (CPUE) of Baor**

The catch per unit effort (CPUE) is the average daily catch per gear type standardized per fishing unit. CPUE is influenced by several factors, primarily the type of gear used and its efficiency, how many hours it is operated for in a day, weather conditions, and location of fishing. CPUE varies between gears and years in different *baors*.




Figure 04. Catch per Unit Effort (CPUE) of different fishing nets used in *baor*

**Fishing Traps:** This type of fishing gear was made from bamboo plate with twine. Fish traps are not considered as destructive gear until it creates any barrier on the migratory route of fish. Relatively small sized wild fishes were voluntarily trapped in this type of gear. Different types of fishing traps were operated in shallow region of the *baor* during the study period.

**Charo / Vair:** Vair is a Long box like trap with a door extending from its base to its apex used in shallow portion of the *baor*. Made of split bamboo sticks tied with creeper or cane. Small opening at the apex for collecting fish. In front of Vair a barrier is created with split bamboo made Bana. The trap checked up every one or two hour interval. Usually operated 1 to 12 hrs/day in the month of July to January.


**Ghuni/Kholsun:** Rectangular box shaped. Made of split bamboo tied with jute rope or cane. Kholsun consists of two doors from its apex for fish opening. Small opening present at its apex for collect the fish. Usually it set in shallow part of the *Baor* with the help of bamboo pole or tree branches during early morning or evening.

**Bitte:** Basket shaped, made of split bamboo with two or three entrances. There is an opening on the trap for collecting fish. The trap is sunken (1 to 1.5 feet below from water surface) in shallow, where small current is present during early morning and evening. Checked up every one or two hour interval.


**Table.4 Different fishing techniques operating in Bukbhora Baor**

<b>Fishing Traps operating in Baor</b>										
Type of gear	Local name	Description			Mesh size (cm)	Materials used	Fishing period	Number of trap used/unit	Major fish caught	
		Length (cm)	Height (cm)	Breadth (cm)						
Trap	Bitte	45	25	25	0.8	Steel wire	Jun-Aug	15-30	Punti, Koi, Baim, Punti, Tengra, Katchki, Mola, Chanda, Khalisha	
	Charo/vair	40		15	1.0-1.5				Rui, Catla, Punti, Koi, Baim, Tengra, Katchki, Mola, Boal, Gozar, Shol, Taki	
	Ghuni/Kholsun	25-40	9-20	9-20	0.2-0.5			20-30	Baim, Koi, Punti, Tengra, small prawn	
<b>Hook and lines operating in Baor</b>										
Type of gear	Name of gear	Number of hook per line of lift		Bait used	Catch/ Haul (kg)	Fishing period	Major fish caught			
Hook and line	Dhawn borshi	Several hundred		Earthworm	0.25- 2	June-Sep	Shol, taki, tengra, punti, boal etc			
	Nol/Dhap borshi			Small live fish, earthworm, small frog etc.	0.25- 1.5	June-Jan	Shol, taki, tengra, boal etc			
	Chip borshi	1 hook		Small live fish, earthworm, small frog etc.	0.03- 0.05	All seasons except rainy day and winter	Koi, Shing, Punti, Tengra and some carp species			
<b>Wounding gears used in Baor</b>										
Type of gear		Name of gear	Length (m)	No. of iron sticks	Catch/ Haul (kg)	Fishing period	Major fish caught			
Spear/Harpoon (Wounding gear)		Koach	1-3	10-20	1-2	Jul-Jan	Rui, Catla, Kakila, shol, boal, ayre, gazar, baim			
		Teta	1.5-2.1	6	0.5-2	Jul-Jan	Taki, Boal, Shol, Bele			
<b>Fishing crafts used in Baor</b>										
Craft name	Size (m)			Shape of bottom	Persons need to operate	Boat type	Life span (years)	Using gears		
	Length	Height	Width							
Kosha	8-10	1-2	2-4	Flat	2-3	Non mechanized	4-5	Khepla jal, Puti jal, Current jal		
Dinghi	5-6		2-3	Rounded	5-6			Ber jal, fash jal, veshal jal		

Fish of the Cypriniformes order was most commonly caught in fishing gears. Their percentage of being caught was 76%. Perciformes order in 72% fishing gear, Siluriformes order were caught in 68% fishing gear, Synbranchiformes order in 52% fishing gear and Beloniformes (35%). Like this, Osteoglossiformes and Clupeiformes both order were caught in 12% fishing gear.



**Figure 05.** Compositions (%) of different categories of fish harvested with different fishing gear



**Figure 06.** Cultured fish harvested with different fishing gear

**Comment [A7]:** consistent in creating image attributes




Figure 07. Captured fish (SIS) harvested with different fishing gear

Comment [A8]: consistent in creating image attributes

## DISCUSSION

About eight (08) various types of fishing gear were classified under six major classes , where three types of fish traps, three types of hook and lines gears, two types of wounding gear and two types of fishing crafts) were found for catching fishes in the following *baor* (Table 2,3,4,5). Adjacent to many researcher such as Nurullah et al. (2005), Ahmed and Hambrey, (2005); Ahmed, (2008); Sayeed et al., (2014) studied on the different fishing techniques performed in open water bodies like beels and rivers. Majumdar et al., (2020) stated that 15 different types of fishing gears (8 major categories) are used for fishing by the fishermen at Chinadi Beel. Similarly, Flowra et al. (2011) and Rahman et al. (2015) who acknowledged 16 different gears under 7 categorizes that supported the present findings. Nonetheless Rahman et al. (2016) and Jaman et al. (2019) found total seven and eighteen major types of fishing gears defined in the Payra River, respectively. Furthermore, Tikadar et al., (2021) found 10 unique gears under 4

categories to operate in the Gorai River. Ferdoushi (2018) reported that total of thirty (30) different types of fishing gears including 10 fishing nets, 9 fishing traps, 9 wounding gears, 2 fish aggregating devices and hand fishing were employed to catch fishes in Ashura beel. Siddiq et al. (2013) conducted a study on Dogger beel and mentioned 5 types of fishing nets, 3 fishing traps, 2 hooks and 2 spears.

Different types of carp species (locally known as 'Raja Mach') were used to culture (fry are released and then harvest when these are the marketable size). Furthermore, various types of non-stocked fishes especially SIS were found in *Baor* which were reared and reproduced naturally (locally known as 'Rani Mach'). Captured SIS Chapila (*Gudusia chapra*) and cultured fish Catla (*Catla catla*) were found the highest harvested fish. The terms kochal (Seine net) and komor (drag net) fishing were local name in the study area, which were mostly used for fishing the carps. Initially, the kochal jal (seine net) is operated for 5 to 10 days and by this time the runaway fish took shelter into bush parks (locally called komor) that set in nearby areas. At that stage, the fishers were stopped operating the kochal jal and started fishing in the komors by encircling those with komor jal (drag net). Maximum fishers had pieces of kochal jal and komor jal (2-5 pieces) and those were jointed together (each of 18 fishers groups separately work) during the harvesting period. Komor fishing (drag netting) was principally adopted as a method to catch the remaining fishes in the *baor* after kochal fishing. Different fishing gears such as entangle net of selective mesh size, borshi, ghuni/britti etc. were used for SIS harvesting. But now-a-days, these SIS species are going to be disappeared despite of their limited ability to reproduce naturally due to environmental degradation as well as habitat degradation.

Based on the usage, the seine net was found as the most usable gear by the fishermen constituting 78.15% followed by drag net (16.05%) among total gear type. The fishermen also

used gill net (2.32%), cast net (1.51%), push net (1.00%), wounding gear (0.50%), hooks and line (0.35%) and traps (0.12%). The result stated that the seine net was highly preferable gear by the fishermen. This investigation recorded that maximum amount of species caught by seine net (ber jal) and few amount by traps. Biswas et al. (2009), Sharif et al.,(2016), Biswas et al.,(2021) also reported that kochal (seine net) fishing were widely used for fish harvesting in different *baors* of jashore region. From Ramnabad River, Ali et al. (2015) noticed the maximum utilization of seine net (60%) and gill net (20%), additionally the lesser utilization were found in lift net (0.6%), cast net (1.2%), push net (1.8%) and trap (0.6%).

Different shapes of fishing gears were enlisted such as rectangular shape (ber jal, current jal), conical shape (jhaki jal), square shape (khora jal) and triangular shape (thela jal) with mesh size ranged from 0.5 to 10 cm, in the study area. Due to destructive nature of some fishing gear, most of the gears were banned (illegal) by departmental authority to save the fish diversity. According to Food and Agriculture Organization (2019), International plan of action to prevent, deter and eliminate Illegal, Unreported and Unregulated fishing in Bangladesh, fishing net with lower mesh was strictly prohibited. Among 16 fishing gears 5 of these were found illegal in this study due to its mesh size and destructive behavior of fish biodiversity. Rahman et al. (2017) found the highest mesh size for gill net (10 cm) and the lowest size push net (0.5 cm) in the Agunmukha River, which concurs well with the present findings. Among those, banned nets, current jal, ber jal, fash jal, puti jal were found the most destructive behavior towards fry fish. Fry fish of almost all species were caught in this net. Only Komar/moi jal, jhaki jal were announced as legal gear by the authority. According to Paul et al. (2021), Jamuna river in the Sirajganj area also faces the destructive behavior of moshari jal which destroyed the juvenile fish.

**Comment [A9]:** It really needs to be explained in detail and its impact on future fish resources in the research location lake

**Comment [A10]:** It is best to state the reasons for the ban, whether because it damages the environment or the mesh size is too small. It's a good idea to include references based on government regulations

**Comment [A11]:** What is the solution if a fishing device like that is found? better say it

In the present findings, the price of fishing gears were ranged from BDT 250-1, 00,000 (Table 1). The highest price was observed in case of seine net (ber jal: about 1, 00,000 BDT) and the lowest price was for punti jal (250-2500 BDT) (Table 1).

According to Portt et al. (2006), Efficiency of gears varies with its type. The maximum CPUE was recorded for ber jal 15-18 kg/effort depending on its massive dimension and mesh size where the CPUE of fash jal, current jal, puti jal, komar/moi jal, khepla jal, thela jal, khora jal was found 10-12, 08-10, 4-5, 02-2.5, <1.50, 01-1.5, 01-1.5 kg/effort, respectively which was fewer than ber jal and greater than any other gear (traps, hooks and line, wounding gears). Maximum fishing efforts per day were correlated with the highest catch composition of any fishing gear. Table 1 represents the highest fishing effort done by those fishing gears which are easy to handle and require less fishermen such as jhaki jal, thela jal, moia jal and khora jal.

However, most of the fishermen caught fish species by using ber jal, komar jal, current jal, thela jal to bring their livelihood. Fishing (harvesting of carp species) started in November-December and completed on 30<sup>th</sup> June in each season. Fishermen caught fish throughout the year but comparative higher amount fish was caught during winter season (November to January) which was agreed with Farid et al. (2013).

Mainly the fishermen operate their trap in rainy season when the water flows of the *baor* are much higher. Different types and sizes of trap are used by the fishermen and it's completely depending upon the target species and their sizes that they would be caught, water depth, seasonality etc. The operation method of these traps are more or less similar. Due to easy operating method and cheap rate, marginal fishermen widely used these trap for commercial and household consumption purposes.

**Comment [A12]:** When writing scientific articles, you should avoid using this term

In Bangladesh, there are about 231 trawler, 34,810 traditional, 32,859 mechanized boat and 1, 88,707 nets & others are engaged in fish harvesting (DoF, 2022). Fishermen use different types of crafts differ in length, breadth, depth, shape and size from region to region of the country. The crafts or boats are built traditionally by the rural carpenters according to their individual plan with planks. It is also difficult to classify them according to the net used because fishermen used the same boat for operating different nets. Most of the fishing boats are manually operated by using paddles and pole propulsion with or without sails. Kosha nauka (plain bottom) are used in low tide area which are non-mechanized and used to catch small fishes. In high tide area the dingi used to catch fishes. This study showed that 40.91% fishers had their own fishing craft (dingi nouka) where 59.09% had no craft.

**Conclusion:**

Size of the gear, mesh size, craft and fisherman required for operation vary with the fishing method and targeted fish species. It depends on season, water level, size of fish, targeted species, fishing area and availability of fisherman. It was found that fishing was widely done in Post monsoon (November-February) and therefore water content was comparatively low. Mainly scine net, drag net, lift net, wounding gear were used during fishing in this season. In order to reverse the trend and ensure sustainability of fish production various measures for protection, conservation and management of fisheries resources have been adopted time to time by the Government. Biodiversity management strategies should be taken to prevent illegal fishing, seizing of destructive fishing gears and also protecting potential breeding habitats. Mass awareness must be required to protect the threatened fish species from being extinct from this wetland; also fishermen and by protecting various fish fauna resources. It is recommended to

**Comment [A13]:** In conclusion, it is best to answer the research objectives related to the number of fishing gear, CPUE, sustainability and illegal fishing gear

strong implementation of all conservation guidelines and acts to stop illegal catching practices and pollution in the *baors*.

#### **CRedit authorship contribution statement**

**Anik Talukdar:** Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Writing –original draft. **Amrita Shaha:** Formal analysis, Investigation, Methodology, Writing – review & editing. **Nasima Begum:** Formal analysis, Investigation, Methodology, Writing – review & editing. **Rakibul Islam:** Writing – review & editing. **Akhery Nima:** Writing – review & editing

#### **Acknowledgments:**

We extend our sincere appreciation to the fishermen and local inhabitants of Bukbhora *baor* helped with the fieldwork.

#### **Conflicts of Interest:**

The authors have no conflicts of interest to declare.

#### **Ethics Committee Approval:**

The authors affirm that ethical approval is unnecessary for this study.

#### **References:**

- Abdullah-Bin-Farid, B.M.S., Mondal, S., Satu, K.A., Adhikary, R.K. and Saha, D. (2013). Management and socio-economic conditions of fishermen of the Baluhar *Baor*, Jhenaidah, Bangladesh. *J. of Fisheries*, 1: 30-36
- Ahmed N. (1971). Government of East Pakistan Directories of Fisheries: Fishing Gear of East Pakistan. East Pakistan Fishery Department, East Pakistan

Ahmed, K., Hambrey, J. (2005). Studies on the fish catch efficiency of different types of fishing gear in Kaptai Reservoir, Bangladesh. *Lakes Reservoirs Res. Manag.* 10, 221–234.

Ahmed, K., Hambrey, J. (2005). Studies on the fish catch efficiency of different types of fishing gear in Kaptai Reservoir, Bangladesh. *Lakes Reservoirs Res. Manag.*, 10, 221–234.

Ahmed, M.S. (2008). Assessment of fishing practices on the exploitation of the Titas floodplain in Brahmanbaria, Bangladesh. *Turk. J. Fish. Aquat. Sci.*, 8 (2), 329–334.

Ahmed, M.S. (2008). Assessment of fishing practices on the exploitation of the Titas floodplain in Brahmanbaria, Bangladesh. *Turk. J. Fish. Aquat. Sci.*, 8 (2), 329–334

Alam, M. M., Jahan, S. N., Hussain, M. A., De, M., Goutham-Bharathi, M. P., Magalhaes, A. L., Mazlan, A. G., Simon, K. D. (2013). Length-length relationship, length-weight relationship and condition factor of freshwater fish species of Bangladesh. *AAFL Bioflux*, 6(5):498-509.

Ali, M. M., Hossain, M. B., Al-Masud, M., Alam, M. A. W. (2015). Fish species availability and fishing gears used in the Ramnabad River, Southern Bangladesh. *Asian Journal of Agricultural Research*, 9(1):12-22

Ali, M. S., Islam, M. S., Begum, N., Suravi, I. N., Mia, M., & Kashem, M. A. 2017. Effect of monoculture and polyculture systems on growth and production of fishes in seasonal waterbodies of Haor villages, Sunamganj district. *Journal of Scientific Research*, 9(3): 307-316.  
<https://doi.org/10.3329/jsr.v9i3.31531>

Ali, M.M., Hossain, M.B., Masud, M.A. and Alam, M.A.W. (2015). Fish species availability and fishing gears used in the Ramnabad river, Southern Bangladesh. *Asian. J. Agric. Res.*, 9(1): 12-22

Azam, A. S., Saha, D., Asadujjaman, M., Mahbub, K. R., Minar, M. H., (2014). Fishing gears and crafts commonly used at Hatiya Island: a coastal region of Bangladesh. *Asian Journal of Agricultural Research*, 8(1):51-58.

Baleta F. N., Beltijar J. G., Bolanos J. M. (2017). Design, fabrication and operation of fishing gears used along the coastal areas of Isabela, Philippines. *International Journal of Fisheries and Aquatic Studies*, 5(2):319-323.

Begum, N., Islam, M., Haque, A., & Suravi, I. (2017). Growth and yield of monosex tilapia *Oreochromis niloticus* in floating cages fed commercial diet supplemented with probiotics in freshwater pond, Sylhet. *Bangladesh Journal of Zoology*, 45(1), 27–36.

<https://doi.org/10.3329/bjz.v45i1.34191>

Biswas A., Tripti, J.K., Farid, M.A., Rahman, M.H. and Ghosh, S. (2021). Present status of fish biodiversity and socio-economic conditions of fishermen of the Kannayadaha *baor*, Jashore, Bangladesh. *Int. J. Biosci.*, 18: 120– 12

Biswas, M.M.R., Islam, M.F, Rahman, M.M., Kawsar, M.A., Barman, S.K. (2009) Fisheries management scenarios of two *Baors* in the district of Chuadanga, Bangladesh. *J Innov. Dev. Strategy*, 3(5):11-15

Chakraborty, S.C.; Hossain, M.A. and Hoq, M.E. (1995). Traditional inland fishing methods in Bangladesh. *J. Asiat. Soc. Bangladesh Sci.*, 21 (1): 19-27.

Ferdoushi Z., M. Rana, N. Gupta, Y. Ara and A. Hossain. (2018). Fishing gears and their targeted species of ashura beel in Dinajpur of Bangladesh. *Journal of Science and Technology*. 16: 10-23

Flowra, F.A., Islam, M.A., Jahan, S.N., Samad, M.A. and Alam, M.M. (2011). Status and decline causes of fishing activities of the Baral River, Natore, Bangladesh. *J. Sci.Found.*, 9(1-2): 115-124.

Galib, S.M., Samad, M.A., Kamal, M.M., Haque, M.A. and Hasan, M.M. (2009). A study on fishing gears and methods in the Chalan Beel of north-west Bangladesh. *Journal of Environmental Science & Natural Resources*, 2(2):213-218.

Ghosh D., Biswas, J.K. (2017). Fish Fauna faces anthropogenic double trouble: erosion of fish diversity in tropical Oxbow Lake of the Ganga River Basin in Eastern India. *J. Biodivers. Endanger. Species*. 5 (2) (2017), p. 188

Hameed M.S. and Boopendranath M.R. (2000). Modern fishing gear technology. Daya Publication House, New Delhi, 186. ISBN: 9788170352235

Jaman, M.N., Hasan, M.M., Ripon, M.A.S., Debnath, S.K. and Hoque, M.S. (2019). Efficiency of fishing gears used in the Payra river at Dumki upozila in Patuakhali district. *Int. J. Oceanogr. Aquacul.*, 3(1): 000157.

Karim, E., Liu, Q., Sun, M., Barman, P. P., Hasan, S. J., & Hoq, M. E. (2019). Assessing recent gradual upsurge of marine captured Hilsa stock (*Tenualosa ilisha*) in Bangladesh. *Aquaculture and Fisheries*, 4(4), 156-165.

Majumdar, B.C., Paul, S.I., Hasan, M., Kabir, T., Islam, M. and Kabir, I.E. (2020). Fish Biodiversity Assemblages and Fishing Gears Used at Chinadi Beel in Narsingdi District of Bangladesh. *International Journal of Agriculture, Environment and Biotechnology*, 13(4): 403-413

- Mia, M., Islam, M. S., Begum, N., Suravi, I. N., & Ali, S. 2017. Fishing gears and their effect on fish diversity of Dekar *haor* in Sunamganj district. *Journal of Sylhet Agricultural University*, 4: 111-120.
- Nurullah, M., Kamal, M., Wahab, M.A., Islam, M.N., Yasmin, L., Shakuntala H. Thilsted and Mazid, M.A. (2005). Present status of harvesting, transportation and marketing of freshwater Small Indigenous Species of Fish (SIS) of Bangladesh, *Bangladesh J. Fish. Res.*,9(2): 159-168
- Paul, S. I., Majumdar, B. C., Hasan, M., Sarker, A. K., Baidya, A., & Hakim, M. (2021). Fish Biodiversity, Threat Status and Conservation Significance of the Jamuna River, Bangladesh. *Croatian Journal of Fisheries*, 79(4), 173-186.
- Portt, C. B., Coker, G. A., Ming, D. L., & Randall, R. G. (2006). A review of fish sampling methods commonly used in Canadian freshwater habitats.
- Rahman, A.K.A. (2005). Freshwater Fishes of Bangladesh. Edn 2. *Zoological Society of Bangladesh*, Dhaka, 394 p.
- Rahman, M.M., Rahman, M.B., Akhter Rithu M.N., Hoque M.S. (2016). Observation on selectivity of fishing gears and ichthyofaunal diversity in the Paira River of Southern Bangladesh. *International Journal of Fisheries and Aquatic Studies*. 4(1): 95-100.
- Rahman, M.B., Hoque, M.S., Rahman, M.M. and Nahar, A. (2017). Exploration of fishing gear and fisheries diversity of Agunmukha river at Galachipa upazila in Patuakhali district of Bangladesh. *Iran. J. Fish. Sci.*, 16(1): 108-126.
- Rahman, M.K. and J.N. Akhter. (2015). Ecology and management of inland water and fisheries resources of Bangladesh. Ideal Books, 38/2 ka Bangla Bazar, Dhaka. 417 p.

- Rahman, M.M., Hossain, M.Y., Ahamed, F., Fatematuzzhura Subba B.R., Abdallah E.M., Ohtomi J. (2012). Biodiversity in the Padma Distributary of the Ganges River, Northwestern Bangladesh: Recommendations for Conservation. *World J. Zool.* 7(4): 328-337.
- Sayeed, M.A., Hashem, S., Salam, M.A., Hossain, M.A.R., Wahab, M.A. (2014). Efficiency of fishing gears and their effects on fish biodiversity and production in the Chalan Beel of Bangladesh. *Eur. Sci. J.* 10 (30), 294–309
- Sharif, B.M.N., Hassan, R., Islam, M.M., Rahaman, M.M., Das, S.K., Miah, M.I. and Amin, M.R. (2016). A study on fishery management of Bergobindapur *baor* at Chaugachha upazila under Jessore district , Bangladesh. *Asian Australas. J. Biosci. Biotechnol.*, 1: 291–296
- Siddiq, M.A., Miah, M.I., Ahmed, Z.F. and Asadujjaman, M. (2013). Present status of fish, fishers and fisheries of Dogger beel in Hajigonj upazila, Chandpur, Bangladesh. *J. Aquat. Sci.*, 1(2): 39-45.
- Sreekrishna Y. and Shenoy L. (2001). Fishing gear and craft technology. Directorate of Information and Publications of Agriculture, Indian Council of Agriculture Research, New Delhi, 342.
- Talwar, P. K. & Jhingran, A. G. (1991). Inland fishes of India and adjacent countries., Vol.1&2, Oxford & IBH Publishing Company Pvt. Ltd, New Delhi, pp.1-1158.
- Tikadar, K. K., Kunda, M., & Mazumder, S. K. (2021). Diversity of fishery resources and catch efficiency of fishing gears in Gorai River, Bangladesh. *Heliyon*, 7(12).

Yasmin, R., Pramanik, M. M. H., Hoque, M. M., Hasan, M. M., & Karim, E. (2021). Effect of Gear Selectivity and Diversity on Catch per Unit Effort (CPUE) of Haors and Beels of Bangladesh. *Bangladesh J. Fish. Res*, 20(1-2), 143-160.

UNDER PEER REVIEW