

# A review on Integrated Nutrient Management: A roadmap for improving the soil sustainability and crop productivity

---

## ABSTRACT

The excessive use of chemical fertilizers with minimal organic inputs in modern agriculture has resulted in nutrient deficiencies and environmental problems. The inorganic chemical fertilizers cannot sustain plant nutrition in highly intensive agricultural systems. Sustainable yield and crop performance are difficult to improve without balanced application of nutrient management practices. The integrated nutrient management (INM) strategies involved the application of vermi-compost, green manure, cover crops, crop residues, vermi-compost and farmyard manure. Since organic manure cannot synchronize the concentrated requirement of plant nutrients as per crop demands, hence balanced integration of organic and inorganic fertilizers helps to improve the crop productivity and economic returns to farmers. Therefore, INM strategy has come up as a field-specific approach in the modern era. The INM strategy builds up the organic matter which improves the crop performance and efficient management of water and nutrients. However, optimizing the timing, rate, source and methods of nutrient application is helpful in achieving the higher apparent recovery efficiency and crop production synchronization with crop demands.

*Keywords:* Crop yield, fertilizers, nutrient management and soil fertility

## 1. INTRODUCTION

The most serious problem being faced by agriculture in the current scenario is excessive chemical fertilizers and groundwater depletion. To meet the increasing population demand, intensive farming using highly resistant cultivated varieties and indiscriminate use of fertilizers has been practiced [1]. Food and Agricultural Organization reported that inorganic fertilizers contribute to an increase in the world's agricultural food production. The unbalanced use of chemical fertilizers without soil testing, conventional farming practices with flooded irrigation practices and reduced or less persistent use of organic manures in appropriate quantity in the past three decades have induced the stagnation of crop yield, deteriorating the crop quality, negative nutrient balance in soil, organic carbon depletion and soil health degradation [2] and [3]. It is not possible to meet the increasing nutritional demand of population growth without balanced nutrient management strategies.

[4]. Increasing in population growth have put a lot of burden on farming community to increase the agricultural production. Most of the Indian soils are deficient in N element [5]. The judicious use of chemical N-fertilizer such as urea cannot sustain soil productivity in highly intensive agricultural systems which can lead to reduction in growth performance and yield-related parameters [6]. Relying applying chemical fertilizer, especially nitrogenous fertilizers cannot sustain soil quality which can lead to degrading soil recuperative and crop productivity [7]. The fertilizer prices increase year after year and the indiscriminate use of nitrogenous fertilizers can lead to N losses, reduces the N use efficiency and further results in an increase in total expenditure on crops [8]. The widespread application of inorganic N-fertilizers and the limited use of organic and bio-fertilizers in modern agriculture are resulting in loss of organic carbon (%) and multi-nutrient deficiencies. India consumes 16 % out of the total N-fertilizer consumed globally [9]. High fertilizer consumption in developing countries is due to knowledge gap between farmers and scientist, high population pressure and availability of highly subsidized N fertilizers. The trends of N consumption have increased substantially in India which can lead to nitrate contamination of surface and groundwater [10].

**Need for Integrated Nutrient Management:** There is significantly challenging task of developing countries to meet the agricultural food demands with increasing population growth. Now a days, farmers apply excessive use of hazardous chemical inputs such as fertilizers which leads to environmental deterioration to meet the challenges of modern era. The soil microbe system plays vital impacts on improving the soil quality and agricultural sustainability. The unbalanced use of inorganic fertilizers results adverse effects on groundwater quality which causes air contamination and ultimately causing nitrate leaching hazards. Also, with increase in population growth, it is most difficult criteria to improve the food security of nation. In India, nutrient management strategy played a significant role in increasing food grain production from 52 MT in 1951 to 252 MT during 2015. Currently, agricultural specialists suggest farmers to change their perspective and adopt an integrated nutrient management (INM) strategy to replace a portion of chemical fertilizer with more sustainable, more efficient and eco-friendly sources of nutrients. It also reduces exploitation of resources and improve soil carbon stocks [11]. Manure are eco-friendly alternative to chemical fertilizers which helps in improving the soil health without any environmental pollution [12]. It can help in sustaining the crop yield while reducing the soil and water pollution [13]. The conjoint application of organic, bio and chemical fertilizers helps to increment the agricultural production and soil sustainability, whereas sole application of chemical fertilizer degrades the soil fertility status. The INM greatly improve soil health and biological activity than sole application of chemical fertilizers. It

is now possible to optimizing the nutrients management strategies to account the variations in the nutrient supplying capacity of soil through variable rates and timings of fertilizer application [14].

The INM can improve the microbial and slow persistent decomposition of biological matter which helped in conversion of organic N to plant available form and hence can improve the soil fertility along with fruit yield [15]. Nutrient management strategy such as split application of fertilizers, applying organic manure in their integrated combined application have proven to be very effective in increasing nutrient use efficiency, crop yield and reducing nutrient losses such as leaching, denitrification, etc [16].

## **2. CONCEPT OF INTEGRATED NUTRIENT MANAGEMENT**

For optimal soil fertility and crop production, nutrients play a significant role in improving the soil structure. The right timing, rate and source of nutrients application influences the higher apparent recovery of added nutrients e.g., ammonium nitrate source of N-fertilizer is best for upland crops besides amide and ammonium sources of N-fertilizer are considered best for lowland rice crop. The split and balanced application of nitrogen and potassium fertilizer is best preferred than basal application whereas for phosphorus, basal placement is generally adopted method due to less fixation. The developed countries using nutrient management stewardship strategy such as the application of phosphorus-based fertilizer/manure, most cereal growing areas will overcome the problem of low phosphorous availability [17]. The phosphorus availability is affected by soil pH and maximum when soil pH between 5.5 and 7.5 whereas soil pH < 5.5 *i.e.* acidic soil causes dissolution of aluminium and iron minerals which precipitates with solution phosphorus but in soil pH > 7.5, basic soil cause excessive calcium present in soil solution which precipitates with phosphorus decrease the availability of phosphorus content. Integrated nutrient supply not only improves the physical, chemical and biological soil properties but also improves the availability concentration of applied and native soil nutrients. The water degradation and environmental sustainability can be improved by adopting carbon sequestration and checking the losses of nutrient to water bodies and atmosphere. The slow and controlled release fertilizers are best nutrient fertilizers which synchronizes the demand of nutrient as per crop demands both in space and time and sufficient nutrient supply from the labile soil.

## **3. COMPONENTS OF INTEGRATED NUTRIENT MANAGEMENT**

The various components of integrated nutrient management are:




Fig 1. Components of integrated nutrient management

#### 4. IMPLEMENTATION OF INTEGRATED NUTRIENT MANAGEMENT

##### Testing methodologies to assess nutrient deficiencies in plant and soil samples are:

- a) Deficiency or toxicity symptom evaluation - The general appearance of crop can provide indications of certain nutritional toxicity or deficiencies syndromes e.g., N deficient plants appear pale yellow, stunted and less vegetative growth than healthy plants
- b) Plant tissue analysis and soil analysis - where signs or syndrome of leaves are not clearly visible, tissue and soil samples can be examined in a laboratory after harvesting and compared with a reference plant sample from a healthy plant.
- c) Scientific examination of different constraints and opportunities in modern strategy for managing soil fertility and how they connect to the diagnosis of nutrients, for e.g. excessive or inadequate application of fertilisers.
- d) Evaluation of farming system's productivity and soil sustainability - The balanced application of nutrients required by crop is influenced by soil type, climatic conditions, tillage practices, nutrient management approach and *In-situ* straw management technologies. Once these physical, chemical and biological factors are understood, so that optimal INM technologies can be used.

- e) Environment friendly INM strategy research and development - The field specific need for locally available resources means that farmer participate in the soil and water analysis adopted the INM practices.

**BASIC PRINCIPLES OF INTEGRATED NUTRIENT MANAGEMENT INCLUDES:**




Fig 2. Basic Principles Of Integrated Nutrient Management Includes

The timing and levels of nutrient application in accordance with the leaf's nutrient requirements, is necessary to achieve maximum yields and improve nutrient-use efficiency [18]. The N-fertilizers with most frequently applying can potentially reduce the environmental pollution but increases yield and quality of crop. Li et al. [19] reported that nitrogen fate is an integrated consequence of nitrogen uptake, immobilization, nitrogen losses like volatilization, denitrification, leaching and runoff.

## **5. IMPACT OF INTEGRATED NUTRIENT MANAGEMENT**

### **5.1 Effects on Soil Fertility**

The ability of soil to continue functioning as critical living system within land-use boundaries is known as soil health. It is due to soil containing biological components that plays a key role in ecosystem function [20]. Plants are often unable to grow in poor soil structural condition which consequently limit their capacity to explore the soil depth for water and nutrients [21]. The main components of integrated nutrient management are farmyard manure (FYM), vermicompost, crop residues, bio-fertilizer, green manure and chemical fertilizer which help to improve soil properties in cropping system. The physico properties such as soil structure, aeration, porosity, infiltration rate and water holding capacity indirectly

improves the health status of soil. Kumar et al [22] studied that soil sustainability and crop productivity by applying chemical or synthetic fertilizers have negatively effect on bio-geochemical cycles [9]. Also, usage of inorganic fertilizer caused denitrification, leaching and run-off, especially nitrogen (N) and phosphorus (P) which results in environmental and global warmth. The different sources of nutrients such as manure, bio-fertilizers, soil, irrigation water, and atmosphere can change the crop productivity. The nutrients removal by crops from the surface soil exceeded their restoration through fertilizers; manures causing unbalanced nutrients in soil were also reported by Gangwar and Prasad [23]. Kumar et al. [24] studied that organic manures in combination with fertilizers can enhance crop performance, productivity and profitability to famers. The decrease in soil pH with the conjoint application of N fertilizer with FYM may be attributed to the release of carbon dioxide and the production of organic acid during the mineralization and decomposition of FYM [25]. Rajneesh et al. [26] revealed that an increase in the level of N fertilizer can decrease the soil pH due to the release of H<sup>+</sup> ions and hydrolysis of inorganic urea. The decrease in surface soil pH has a positive influence on the availability of different nutrients, notable Zn, Mn, Fe and P [27]. The lower EC values under organic amendment (FYM) could be attributed to increased soil water holding capacity because of improved soil aggregation [28]. Soil organic carbon (SOC) is the most imperative indicator of agricultural sustainability and soil health. The greater SOC with FYM may be attributed to FYM's slower rate of decomposition (less and persistent mineralization rate) because of polyphenol as well as lignin content. An improvement in SOC content might be due to the improvement of root anatomy and increased plant residue with the increased nutrient application through manure and chemical fertilizers [29]. The INM treatments have higher available N as compared to sole inorganic N fertilizer due to the improvement of SOC and slow mineralization of available N in surface soil [30] and [31]. The FYM with inorganic N fertilizer have higher available P content than inorganic N fertilizer which might be due to soil organic matter (SOM) coating the sesquioxides (Al-P and Fe-P), rendering them inactive and reducing the P fixing capacity of surface soil resulting in improvement of P in soil [32]. The increase in available K of surface soil with combined application of N fertilizer with FYM may be attributed to improve the water holding capacity (WHC) and slow mineralization of nutrients. The study revealed that FYM can significantly increase the available K content in INM plots as compared to inorganic N fertilizer plots [33]. The INM treatments have higher available micronutrients as compared to sole inorganic N fertilizer due to increasing soil microbial activity during mineralization and decomposition of organic matter which improved the DTPA-extractable Zn, Cu, Fe and Mn content in surface soil by preventing leaching, fixation and precipitation [34].

## 5.2 Effects on Environmental Benefits

Organic farming was shown to be successful strategy for improving soil sustainability, environmental quality and maintaining soil ecosystem services [35] and [36]. Efficient use of on-farm inputs, crop residues, and limited use of off-farm inputs sustained the soil fertility, nutrients and environmental quality in organic farming. The extensive integration of organic leftovers in organic farming lead to accumulation of high soil organic matter in organically amended soils which supports productivity of an agroecosystem [37]. Bhattacharya et al. [38] reported that organic carbon in soil increase under long-term legume-based crop rotation in organically managed soils. In addition to organic practices, i.e., incorporation of compost and manures, crop rotation, bio-fertilizers and cover crops not only improve soil organic matter but also improve physico-chemical and biological properties of soil [39] and [40]. Numerous studies have evaluated that permanent additive of organic manures, green manures, crop residues and bio-fertilizers in cropping system improved soil physical properties, i.e., bulk density, water-holding capacity (WHC), aggregate stability and hydraulic conductivity; chemical properties i.e. pH range (0.3-0.9), electric conductivity ( $0.2-0.5 \text{ dS m}^{-1}$ ), organic carbon (32%), available N (47%), available P (90%), available K (50%) and biological properties, i.e., microbial biomass carbon (132%) dehydrogenase activity (347%) and phosphatase activity (203%) relative to control treatments [41] and [42]. Significant physico-chemical properties and variation in micro-environment as a result of organic residue have shown positive effects on DTPA-extractable Fe, Cu, Zn, and Mn content in soil [43]. Dhaliwal et al. [2] observed significant apparent recovery of DTPA-extractable micronutrients with the addition of organics, crop residues, and bio-fertilizers in soils with alternate drying and wetting conditions. The integrated application of FYM, green manures and bio-fertilizers and subsequent microbial degradation of amendments significantly improved soil organic carbon (SOC), available N, P, S and DTPA extractable micronutrients. The integrated incorporation of organic manures with crop residues and bio-fertilizers results to improve the nutrient availability and crop productivity [44]. Significant improvement in crop productivity and DTPA-extractable micronutrients were observed through several researchers when 25% to 50% of the recommended dose of nitrogen was substituted through different organics, i.e., wheat straw, FYM and green manures in rice followed by wheat crop each year [32]. Walia and Dhaliwal [34] reported a positive correlation of Fe, Mn, Zn with the SOC in soil. Further, organic matter affects growth and yields either by directly supplying nutrients or indirectly by modifying soil physical properties such as stability of aggregates, porosity and available water capacity that can improve the root environment and stimulate plant growth. Organic matter not only increases the water

holding capacity of the soil but also proportionates water available for plant growth and improves physical properties of soil [39].

### **5.3 Effects on Crop Productivity**

The crop yield is influenced by many parameters, besides its most important parameter for attaining net profit for the farmers. It is combined effect of metabolic and physiological parameters. It helps us to understand the several factors that limit the adaptability of crop. The yield was increased in INM plots because the application of organic manure can supplement the N fertilizer to crop for developing plant growth that reflects on the resultant crop productivity [45]. A higher source generates a more efficient sink, which increases yield and dry matter production. The rate of photosynthesis is measured by chlorophyll content which serves as an indicator of biomass accumulation. The higher nutrient availability through microbial activity enhancement with conversions of some of the nutrients from unavailable to available forms which improve the physical, chemical and biochemical conditions for better nutrient translocation and nutrient assimilation. Babhulkar et al. [46]; Singh et al. [47] and Kumar et al. [48] studied that significant positive correlation observed between available nutrients and crop yield. The microbial and slow decomposition of FYM, crop residues and continuous slow release of nutrients throughout the growth stage of crop coupled with higher nutrient assimilation. Singh et al [49] provide conducive situation for higher yield performance of different field crops. The optimal soil physical conditions are important prerequisite for sustaining fertility management. The soil aggregates breakdown, platy soil structure, root restrict growth and consequently limit their capacity to explore the soil profile for water and nutrients [21]. Soil organic matter affects growth and yield, either directly by supplying nutrients or indirectly by modifying soil properties that results the rhizosphere environment and better plant growth. The INM strategies of sustainable crop production plays a significant role through reducing chemical fertilizers. Kamboj et al. [50] studied that integrated N management practice with FYM improve growth, quality and yield of chilli as compare to sole use of inorganic N-fertilizers. The addition of organic manure, lime and bio-fertilizers enhanced soil organic carbon, moisture retention capacity and infiltration rate [51]. Bio-fertilizers sustained fertilizer nutrient use through biological nitrogen fixation, which further solubilize fewer mobile nutrients and leading to sustainable agriculture system [52].

## **6. CONCLUSION AND FUTURE PROSPECTS**

The integrated nutrient management emphasis on the balance between organic and inorganic inputs in current context of global agricultural challenges, such as soil degradation and environmental sustainability. The INM integrates scientific concepts such as nutrient cycling, soil microbe interaction, and the environmental implications of fertilizer use. It

improves the microbial decomposition of organic matter which helped in conversion of organic N to plant available form and hence can improve the soil fertility along with fruit yield. The conjoint integration of organic, bio-fertilizers and chemical fertilizers improves the productivity and agricultural sustainability, whereas sole application of chemical fertilizer degrades the soil fertility. The nutrient management techniques such as split application of fertilizers, the use of organic manure and their combined application have proven to be very effective in increasing nutrient use efficiency, crop yield and reducing nutrient losses such as leaching, denitrification, etc. The fertilizer prices increase year after year and the indiscriminate use of nitrogenous fertilizers can lead to N losses, reduces the N use efficiency and further results in an increase in total expenditure on crops. Therefore, integrated nutrient management is an eco-friendly approach which helps to improve the soil fertility, crop productivity and environmental sustainability.

#### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### **7. REFERENCES**

1. Das K, Bhattacharyya S, Sudhishri AR, Sharma YS, Saharawat, KK, Bandyopadhyay and Bhatia A. Conservation agriculture in an irrigated cotton-wheat system of the western Indo-Gangetic Plains: Crop and water productivity and economic profitability. *Field Crops Res.* 2004;158:24-33.
2. Dhaliwal SS, Naresh RK, Mandal A, Walia MK, Gupta RK, Singh R and Dhaliwal MK. Effect of manures and fertilizers on soil physical properties, build-up of macro and micronutrients and uptake in soil under different cropping systems: A review. *J Plant Nutr.* 2019;42:2873-2900.
3. Jat HS, Datta A, Sharma PC, Kumar V, Yadav AK, Choudhary M, Choudhary V, Gathala MK, Sharma DK and Jat ML. Assessing soil properties and nutrient availability under conservation agriculture practices in a reclaimed sodic soil in cereal-based systems of North-West India. *Arch Agron Soil Sci.* 2017;64:531-45.
4. Zhang F, Gao X, Wang J, Liu F, Ma X, Cao H, Chen X and Wang X. Sustainable nitrogen management for vegetable production in China. *Front Agri Sci Eng.* 2022;9:373-85.
5. Rekha GS, Kaleena PK, Elumalai D, Srikumaran MP and Maheswari V N. Effects of vermicompost and plant growth enhancers on the exo-morphological features of (*Capsicum annuum* L.). *Int J Recycl Organic Waste Agric.* 2018;7:83-88.
6. Powlson DS, Gregory PJ, Whalley WR, Quinton JN, Hopkins DW, Whitmore AP, Hirsch PR and Goulding KW. Soil management in relation to sustainable agriculture and ecosystem services. *Food Policy.* 2011;36:72-87.

7. Majeed A, Abbasi MK, Hameed S, Imran A and Rahim N. Isolation and characterization of plant growth promoting rhizobacteria from wheat rhizosphere and their effect on plant growth promotion. *Front Microbiol.* 2015;6:78-97.
8. Chai R, Ye X, Ma C, Wang Q, Tu R, Zhang L and Gao H. Greenhouse gas emissions from synthetic nitrogen manufacture and fertilization for main upland crops in China. *Carbon Balance Manag.* 2009;14:1-10.
9. Bijay-Singh. Nitrogen use efficiency in crop production in India: trends, issues and challenges. *Agric Res J.* 2023;12:32-44.
10. Bijay-Singh and Craswell E. Fertilizers and nitrate pollution of surface and ground water: an increasingly pervasive global problem. *Springer Nature Sci Rev.* 2021;3:518-34.
11. Koshale C, Kurrey DK and Banjare LD. Effect of organic manure and inorganic fertilizer on growth, yield and physiological parameter of chilli. *Int J Chem Stud.* 2018;6:118-22.
12. Singh A and Shivay YS. Zinc application and green manuring enhances growth and yield in basmati rice (*Oryza sativa* L.). *Indian J Plant Physiol.* 2015;3:289-96.
13. Khan A, Shah SNM, Rab A, Sajid M, Ali K, Ahmed A and Faisal S. Influence of nitrogen and potassium levels on growth and yield of chillies (*Capsicum annum* L.). *Int J Farm Sci.* 2014;3:260-64.
14. Bijay-Singh and Singh VK. Advances in nutrient management in rice cultivation in achieving sustainable cultivation of rice; Burleigh Dodds Science Publishing Limited: Cambridge, UK 2017;2:25-68.
15. Sinha SK, Jha CK, Kumar V and Pandey SS. Yield and soil organic carbon pool in relation to soil fertility of sugarcane (*Saccharum* species hybrid complex) plant-ratoon system under integrated nutrient management. *Indian J Agron.* 2017;62:25-30.
16. Shafeek MR, Nagwa MK, Hassan MK, Awatef, Beheiry G and Rizk FA. Productivity of sweet pepper plant (*Capsicum annum* L.). as affected by different sources of nitrogen and sulphur fertilizers under newly reclaimed soil. *J Appl Sci Res.* 2012;8:5365-71.
17. Ortiz-Monasterio JI, Pena RJ, Pfeiffer WH and Hede AH. Phosphorus use efficiency, grain yield, and quality of triticale and durum wheat under irrigated Conditions. 2002;3:45-51.
18. Cassman KG, Dobermann A and Walters DT. Agroecosystems, nitrogen use efficiency and nitrogen management. *Ambio.* 2002;31:132-40.
19. Li Y, Simunek J, Zhang Z, Jing L and Ni L. Evaluation of nitrogen balance in a direct-seeded rice field experiment using Hydrus-1D. *Agri Water Mgt.* 2015;148:213-22.
20. Hane RB. Influence of integrated use of organics to sustain productivity of rice. M.Sc. (Ag.) Thesis, Dept. of Soil Sci. IGAU, Raipur. 2001.
21. Haynes RJ and Naidu R. Influence of lime, fertilizer, and manure applications on soil organic matter content and soil physical conditions: A review. *Nut Cycl Agroecosyst.* 1998;51:123-37.

22. Singh K, Tirunima P, Yadav SL, Yadav SS and Singh N. Effect of NPK levels with biofertilizers on growth and yield of chilli (*Capsicum annuum* L.) in alluvial soil of Gwalior, Madhya Pradesh. *Int J Plant Soil Sci.* 2022;34:1337-42. <https://doi.org/10.9734/ijpss/2022/v34i2231504>.
23. Gangwar B and Prasad K. Cropping system management for mitigation of second-generation problems in agriculture. *Indian J Agric Sci.* 2005;75:65-78.
24. Kumar AS, Sharma A and Mishra S. Application of farmyard manure and vermicompost on vegetative and generative characteristics of *Jatropha curcus*. *J Phyt.* 2009;1:206-22.
25. Singh NJ, Athokpam HS, Devi KN, Chongtham N, Singh NB, Sharma NB, Sharma PT and Dayananda S. Effect of farmyard manure and press mud on fertility status of alkaline soil under maize-wheat cropping sequence. *African J Agric Res.*2015;10:2421-31.
26. Rajneesh RP, Sankhyan NK and Kumar R. Long-term effects of fertilizers and amendments on depth-wise distribution of available NPK, micronutrients cations, productivity and NPK uptake by maize-wheat system in an acid alfisol of north western Himalayas. *Commun Soil Sci Plant Anal*2017;48:2193-2209.
27. Benbi DK, Brar K, Toor AS, Singh P and Singh H. Soil carbon pools under poplar-based agroforestry, rice-wheat and maize-wheat cropping systems in semi-arid India. *Nutr Cycl Agroecosys.* 2012;92:107-18.
28. Duhan L and Singh S. Effect of green manuring and nitrogen on yield and uptake of micronutrients by rice. *J Indian Soc Soil Sci.*2002;50:178-80.
29. Dhaliwal SS, Ram H, Walia SS, Walia MK, Kumar B and Dhaliwal MK. Long-term influence of nutrient management on carbon and nutrients in Typic-ustochrept soils. *Commun Soil Sci Plant Analysis.* 2019;50:2420-28.
30. Guo L, Wu G, Li Y, Li C, Liu W, Meng J, Liu H, Yu X and Jiang G. Effects of cattle manure compost combined with chemical fertilizer on topsoil organic matter, bulk density and earthworm activity in a wheat-maize rotation system in eastern China. *Soil Till Res.*2016;156:140-47.
31. Yan F, Zhang F, Fan X, Fan J, Wang Y, Zoa H and Li G. Determining irrigation amount and fertilization rate to simultaneously optimize grain yield, grain nitrogen accumulation and economic benefit of drip-fertigated spring maize in northwest China. *Agric Water Manage.*2021;243:106440.
32. Subehia SK, Sepehya S, Rana SS, Negi SC and Sharma SK. Long-term effect of organic and inorganic fertilizers on rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) yield, and chemical properties of an acidic soil in the western Himalayas. *Exp Agric* 2013;49:382-94.
33. Urkurkar JS, Tiwari A, Chitale S and Bajpai RK. Influence of long-term use of inorganic and organic manures on soil fertility and sustainable productivity of rice (*Oryza sativa* L.) and wheat (*Triticum aestivum* L.) in Inceptisols. *Indian J Agric Sci.*2010;80:208-12.
34. Walia MK, Walia SS and Dhaliwal SS. Long-term effect of integrated nutrient management of properties of typic ustochrept after 23 cycles of an irrigated rice (*Oryza sativa* L.) - wheat (*Triticum aestivum* L.) system. *J Sustain Agric.*2010;34:724-43.

35. Ram M, Davari MR and Sharma SN. Direct, residual and cumulative effects of organic manures and biofertilizers on yields, NPK uptake, grain quality and economics of wheat (*Triticum aestivum* L.) under organic farming of rice-wheat cropping system. *J Organic Syst.* 2014;9:16-30.
36. Thuithaisong C, Parkpian P, Shipin OV, Shrestha RP, Naklang K, Delaune RD and Jugsujinda A. Soil quality quality indicators for predicting sustainable organic rice production. *Commun Soil Sci Plant Analysis.* 2011;42:548-68.
37. Dotaniya ML, Datta SC, Biswas DR and Kumar K. Effect of organic sources on phosphorus fractions and available phosphorus in typic haplustept. *J Indian Soc Soil Sci.* 2014;62:80-83.
38. Bhattacharya RS, Chandra RD, Singh S, Kundu AK, Srivastva and Gupta HS. Long-term farmyard manure application effects on properties of a silty clay loam soil under irrigated wheat-soybean rotation. *Soil Till Res.* 2007;94:386-96.
39. Naresh RK, Kumar P, Malik S, Kumar M and Choudhary U. Effects of tillage, residue and nutrient management on top soil carbon stocks and soil labile organic carbon fractions in the Indo-gangetic plains of North West India: A review. *J Pharm Phytochem* 2018;7:1818-42.
40. Singh G, Kumar D and Sharma P. Effect of organics, biofertilizers and crop residues application on soil microbial activity in rice-wheat and rice-wheat-mungbean cropping systems in Indo-Gangetic plains. *Cogent Geosci.* 2015;1:1-14.
41. Biswas DR, Ghosh S, Ramchandran BB, Basak and Moharana PC. Dependence of thermal and moisture sensitivity of soil organic carbon decomposition on manure composition in an Inceptisols under 5 years old maize-wheat cropping system. *J Geophysical Sci.* 2018;123:1-14.
42. Ghosh A, Bhattacharyya R, Meena MC, Dwivedi BS, Singh G, Agnihotri R and Sharma C. Long-term fertilization effects on soil organic carbon sequestration in an Inceptisol. *Soil Till Res.* 2018;177:134-44.
43. Dhaliwal SS, Manchanda JS, Walia SS and Dhaliwal MK. Differential response of manures in transformation of DTPA and total zinc and iron in rice transplanted on light soil of Punjab. *Int J Sci Environ Technol* 2013;2:300-12.
44. Li-Li BF, Xiang-chao, De-gui T, Bin W, Lin B and Tie-jun Y. Evaluation and comparison of composting rabbit manure mixed with mushroom residue and rice straw. *J Agric Sci Technol.* 2013;11:1069-81.
45. TambeAJ, DhawanAS and Gourkhede PH. Effect of integrated nutrient management on yield, quality improvement and nutrient uptake of chilli. *Int Trop Agric.* 2015;33:3777-81.
46. Babhulkar PS, Wandile RM, Badole WP and Balpande SS. Residual effect of long-term application of farmyard manure and fertilizers on soil properties and yield of soybean. *J Indian Soc Soil Sci.* 2000;48:89-92.
47. Singh S, Singh RN, Prasad J and Singh BP. Effect of integrated nutrient management on yield and uptake of nutrients by rice and soil fertility in rainfed

- uplands. J Indian Soc Soil Sci 2006;54:327-30.
48. Kumar B, Gupta RK and Bhandari AL. Soil fertility changes after long term application of organic manures and crop residues under rice-wheat system. J Indian Soc Soil Sci. 2008;56:80-85.
  49. Singh H, Singh RP, Meena BP, Lal B, Dotaniya ML, Shirale AO and Kumar K. Effect of integrated nutrient management (INM) modules on late sown Indian mustard (*Brassica juncea* L.) and soil properties. J Cereals Oilseeds. 2018;9:37-44.
  50. Kamboj D, Gill AAS and Jindal SK. Effect of timing and levels of nitrogen application with farmyard manure application on growth and yield of chilli (*Capsicum annuum* L.). J Adv BioBiotechnol.2024;27:11-22. <https://doi.org/10.9734/jabb/2024/v27i91269>.
  51. Meena OP, Maurya BR and Meena VS. Influence of K- solubilizing bacteria on release of potassium from waste mica. Agri Sustain Dev. 2013;1:53-56.
  52. Subedi P, Bhattarai P, Lamichhane B and Kansal A. Effect of different levels of nitrogen and charcoal on growth and yield traits of chilli (*Capsicum annuum* L.) Heliyon. 2023;9:2405-12.