

Preliminary analysis of the feeding habits of two species of the genera *Mobula* (*Mobulajaponica* and *Mobulatarapacana*) in the coastal waters of Côte d'Ivoire, Gulf of Guinea

ABSTRACT

This study provides baseline information on the feeding habits of *Mobula japonica* and *Mobulatarapacana* using data on stomach contents. Specimens were sampled from the small-scale driftnet fishery operating along the coast of Côte d'Ivoire between March 2020 and April 2022. Diet was described using the Prey-Specific Index of Relative Importance (%PSIRI) combining occurrence, numerical and weight percentages. A total of 215 *M. tarapacana* and 210 *M. japonica* ranging in sizes of 154–371 cm DW and 80–369 cm DW, respectively were examined. The large number of empty stomachs for *M. japonica* (71.43%) and *M. tarapacana* (73.02%) may be the result of using hooks, which cause high stress resulting in stomach regurgitation. *Mobulajaponica* and *M. tarapacana* were found to primarily consume the commonly abundant crustaceans such as *Sergestes atlanticus* and euphausiid *Pseudeuphausialatiffrons*, accounting for 95% and 71.13% of the diet, respectively. However, diet of *M. tarapacana* was more diverse, including crabs, algae of the genus *Sargassum*, teleost fishes, molluscs and insects. Analysis of Costello diagram showed that both species were specialist predators even if *M. japonica* has more restricted niche. The Morisita-Horn similarity showed no significant changes in diet according fish sizes and marine seasons.

Keywords: Devil rays, diet, specialist predator, stomach contents, West Africa

1. INTRODUCTION

Rays of Mobulid family are pelagic fishes found in shallow and deep waters in tropical, subtropical and temperate coastal waters [1,2,3]. These species have a crucial ecological role, contributing to the regulation of prey populations and the balance of marine ecosystems [4]. In Côte d'Ivoire, they are a common by-catch in artisanal tuna driftnet fishery targeting tunas [5]. The international market demand causes increased fishing pressure, as ray body parts have high economic value for meat, skin and gill plates [6]. Due to their biological characteristics (low natural mortality, slow growth, late sexual maturation and low offspring [7-2], their excessive exploitation could have a serious impact on the renewal of their population. At this time, the entire *Mobula* spp. has been registered in the Appendix II of the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) whereas *M. tarapacana* and *M. japonica* are classified as Vulnerable or Endangered according to the International Union of Conservation of Nature [8-9].

Understanding the diet of fishes is essential in explaining variations in growth, aspects of reproduction, migration and feeding behaviour [10]. According to Costello [11], it provides information on the diversity of prey ingested, food preferences and trophic niche. For Stewart et al. [12], analysing feeding habits of tropical populations will determine diet preference and allow inference on habitat use and feeding behaviour, which are both very important to the conservation and management of rays. Although studies on the biology and ecology of Mobulid species started since the 17th century, information on their feeding habits still very limited [12]. Couturier et al. [1] reported that their diet is based on zooplankton and ichthyoplankton, but specific differences can occur. For instance, *M. thurstoni* is known to feed on mysid shrimps and euphausiids (Notarbartolo di Sciara, [13]) whereas *M. mobular*, *M. hypostoma*, *M. rochebrunei*, *M. tarapacana*, *M. birostris*, *M. alfredi*, *M. japonica*, and *M. munkiana* feed generally on small fishes and crustaceans like krill (*Meganyctiphanes norvegica*, *Nyctiphanes simplex*), the mysid *Mysidium* sp., and other planktonic organisms [13, 14, 1]. However, no information is known on the feeding habits of *M. tarapacana* and *M. japonica* in the central eastern Atlantic and specifically in the Gulf of Guinea. Thus, the aim of this work is to analyse the composition of their stomach contents and deduce their feeding habits.

2. MATERIALS AND METHODS

Study area

This study was carried out in Côte d'Ivoire's Exclusive Economic Zone (EEZ) bordering the Gulf of Guinea (Atlantic Ocean). This zone extends from Cape des Palmes (8°W) in the West to Cape Three Points (2°30'W) in the East [15], over a length of about 600 km (Fig. 1). Its characteristic is the presence of four marine seasons including two warm seasons (March–June and November–December) and two cold seasons (July–October and January–February) [16]. The surface sea water temperatures vary from 27.0 to 30.6°C and the salinity below 35‰.




Fig. 1. Map showing the Exclusive Economic Zone (EEZ) of Côte d'Ivoire and the study site.

Sampling and laboratory analysis

Samples were obtained monthly at the fishing harbour of Abidjan (Côte d'Ivoire) from March 2020 to April 2022. For each individual, the sex was determined and the body weight was recorded to the nearest kg. The disc width (DW) was then measured to the nearest 0.1cm using a measuring tape. After dissection, stomachs were removed and kept frozen at – 20°C for further analysis in the biology laboratory of the Oceanological Research Centre, Abidjan.

In the laboratory, the stomach contents were washed individually under running water using a 0.5 mm mesh sieve, which retained any undigested food elements in the faeces that could be used for identification (e.g. scales, otoliths, exoskeletons, teeth, etc.). Each prey was then identified to the lowest taxonomic level according to the identification keys of Fisher et al. [17], Schneider [18] and Carpenter et al. [19] and weighed in grams (g). The preys were then counted and weighted and all undetermined preys were considered as unknown preys and classified into a group.

Data analysis

The index of vacuity was used to express the number of empty stomachs encountered as a percentage of the total stomachs examined.

$$\%CV = \frac{\text{Number of empty stomachs}}{\text{Total number of stomachs}} \times 100$$

The contribution of each prey item to the diet was assessed using the prey-specific relative importance index (%PSIRI), as %FO was found to be unknowingly squared in the IRI calculation [20]. The frequency of occurrence (FO), the average percentage abundance (%Ai) and the prey-specific abundance (%PAi) were calculated with the following equations:

Prey-specific abundance (%PN_i, %PW_i):

$$\%PA_i = \frac{\sum_{j=1}^n \%A_{ij}}{n_i}$$

Average percentage abundance (%Ni, %Wi):

$$PA_i = \frac{\sum_{j=1}^n \%A_{ij}}{n}$$

Frequency of occurrence (FO):

$$\%FO = \frac{n_i}{n}$$

where %A_{ij} is the abundance (by counts or weights) of prey category i in stomach sample j, n_i is the number of stomachs containing prey i, and n is the total number of stomachs.

The %PSIRI was then calculated:

$$\%PSIRI_i = \frac{\%FO_i \times (\%PN_i + \%PW_i)}{2}$$

where %FO_i is the percent frequency of occurrence (the number of stomachs containing prey category i divided by the total number of stomachs n) and %PN_i and %PW_i are the prey-specific abundances by counts and weight, respectively.

The classification of prey items followed the method of Rosecchi and Nouaze [21]. For this purpose, preys were first sorted in decreasing order of importance according to their PSIRI and then a cumulative %PSIRI was calculated. The first single item, or group of items, for which cumulative %PSIRI was ≥ 50% was regarded as the preferred food. The %PSIRI values of other important prey items were then added to that of the preferred food until the %PSIRI reached 75%, and these were regarded as secondary prey. Food items between a %PSIRI of 75 and 100% were regarded as incidental prey. The food Overlaps between sexes and marine seasons were analysed using the Morisita [22] Index as follow:

$$C\lambda = \frac{2 \sum_{i=1}^n (P_{ix} \times P_{iy})}{(\sum_{i=1}^n P_{ix}^2 + \sum_{i=1}^n P_{iy}^2)}$$

Where Cλ the Morisita-Horn [22] index, P_{ix} and P_{iy} are proportions (based on %PSIRI) of a prey i consumed by the predator x and y respectively, following the sex and marine seasons. Values between 0 and 0.29 indicate a lower overlap, between 0.30 and 0.59 a medium overlap, and values over 0.60 show a higher overlap [23].

Food strategy

The Costello diagram [24] modified by Amundsen et al. [25] was used to describe the feeding strategies of the species studied. This diagram illustrates the specific abundance of the prey (S_i) as a function of the frequency of occurrence

(%FO) and allows analysis of the importance of the prey, the feeding strategy and the contribution to the niche. The prey-specific abundance was determined through following formula:

$$\%S_i = \left(\frac{\sum P_i}{\sum P} \right) \times 100$$

Where P_i = weight of the prey i and P = total weight of preys into stomach containing prey i .

3. RESULTS AND DISCUSSION

Results

Vacuity index

A total of 215 specimens of *M. tarapacana* and 210 specimens of *M. japonica* with sizes ranging from 154 to 371 cm DW and from 80 to 369 cm DW, respectively were used. Of the stomachs examined, 59 (27.44%) had food and 156 (72.56%) were empty in *M. tarapacana* whereas 60 (28.57%) were full and 150 (71.43%) were empty in *M. japonica*. The vacuity index varied greatly according to marine seasons. For *M. japonica*, the vacuity index was higher in the warm season (78.18%) than in the cold season (69.68%). On the other hand, the values were higher in the cold season (79.81%) than in the warm season (65.77%) for *M. tarapacana*.

Overall diet composition

The dietary spectrum of *M. tarapacana* included 12 prey items, belonging to 8 families which can be grouped into five major categories: crustaceans, algae, teleost fishes, molluscs and insects. The quantitative analysis of the diet showed that crustaceans (%PSIRI = 68.62%), mostly represented by *Sergestes atlanticus* (% PSIRI = 49.83) and euphausiid *Pseudeuphausialatiffrons* (%PSIRI = 17.87%) were the main ingested prey of *M. Tarapacana*. These prey items were followed by the marine algae of the genus *Sargassum* (%PSIRI = 15.76%), composed of *Sargassum natans* (%PSIRI = 8.14) and *Sargassum fluitans* (%PSIRI = 7.63) and several fish species, namely *Remora remora* (%PSIRI = 2.54), *Sardinella aurita* (%PSIRI = 1.81), *Trichiuruslepturus* (%PSIRI = 0.72), *Priacanthusarenatus* (%PSIRI = 0.60) and *Chloroscombrus chrysurus* (%PSIRI = 0.59). The other taxonomic groups (insects and molluscs) representing less than 5.00% of the %PSIRI were incidental food in the diet of this species (Table 1). In contrast, the food spectrum of *M. japonica* is less varied and consisted of two main categories of prey namely crustaceans and algae. Crustaceans were the most important taxonomic group in the diet (PSIRI = 94.67%), followed by prey belonging to the algae of the genus *Sargassum* (PSIRI = 5.33), in particular *S. natans* (%PSIRI= 2.00) and *S. fluitans* (%PSIRI = 3.33) which can be considered as incidental food (Table 2). Among crustaceans, *S. atlanticus* (%PSIRI = 63.33) was the most important prey, followed by *P. latiffrons* (%PSIRI = 31.34%).

Table 1. General overview of the diet of *Mobulatarapacana* captured from March 2020 to December 2022 in Ivorian waters by the artisanal gillnet fishery. %FO = percentage frequency of occurrence; %PN = prey specific number

percentage; %N= number percentages (%N); %PW= percentage of specific weight of these same prey; W = weight percentage and %PSIRI = prey-specific relative importance index; S_i= specific abundance; (ind) = indeterminate.

Categories	Prey	%FO	%PN	%NOT	%PW	%W	%PSIRI	S _i	
Crustaceans		72.88	2.58	69.76	2.31	67.47	68.62	-	Tab
Sergestidae	<i>Sergestes atlanticus</i>	52.53	0.97	50.84	0.93	48.81	49.83	93.62	le 2.
Euphausiidae	<i>Pseudeuphausialatiffrons</i>	18.64	0.95	17.80	0.96	17.95	17.87	99.51	Gen
Portunidae	<i>Portunussayi</i>	1.69	0.66	1.12	0.42	0.71	0.92	42.86	
Algae		22.03	1.28	14.07	1.59	17.46	15.76	-	eral
	<i>Sargassum natans</i>	11.86	0.61	7.29	0.76	8.98	8.14	73.46	over
	<i>Sargassum fluitans</i>	10.17	0.67	6.78	0.83	8.47	7.63	93.46	vie
Teleosts		18.64	2.64	9.14	3.31	10.92	10.03	-	w of
Priacanthidae	<i>Priacanthusarenatus</i>	3.39	0.03	0.12	0.32	1.08	0.60	35.84	the
Clupeidae	<i>Remora remora</i>	3.39	0.75	2.54	0.75	2.54	2.54	75.00	diet
Echeneidae	<i>Sardinellaaurita</i>	3.39	0.50	1.69	0.57	1.92	1.81	32.29	of
Carangidae	<i>Chloroscombrus chrysurus</i>	1.69	0.30	0.51	0.39	0.66	0.58	39.12	
Trichiuridae	<i>Trichiuruslepturus</i>	1.69	0.32	0.54	0.53	0.9	0.72	53.68	<i>Mob</i>
	<i>Fish (ind)</i>	5.08	0.73	3.73	0.75	3.81	3.77	48.81	<i>ulaj</i>
Mollusc		3.39	0.49	0.83	0.45	0.76	0.80	-	<i>apa</i>
Sepiidae	<i>Sepia officinalis</i>	1.69	0.16	0.27	0.16	0.27	0.27	16.09	
Bivalves (ind)	<i>Ind</i>	1.69	0.33	0.56	0.29	0.49	0.53	29.16	<i>nica</i>
Insects (Ind)	<i>Ind</i>	6.78	0.92	6.20	0.50	3.40	4.80	2.60	

captured from March 2020 to December 2022 in Ivorian waters by the artisanal gillnet fishery. %FO = percentage frequency of occurrence; %PN = prey specific number percentage; %N = number percentages (%N); %PW = percentage of specific weight of these same prey; W = weight percentage and %PSIRI = prey-specific relative importance index; S_i= specific abundance.

Categories	Prey	%FO	%PN	%N	%PW	%W	%PSIRI	Si
Crustaceans		94.99	94.99	2.00	94.36	1.99	94.67	-
	<i>Sergestes atlanticus</i>	63.33	63.33	1.00	63.33	1.00	63.33	100.00
	<i>Pseudeuphausialatiffrons</i>	31.66	31.66	1.00	31.03	0.99	31.34	97.70
Algae		6.66	5.01	1.58	5.63	1.69	5.33	-
	<i>Sargassum natans</i>	3.33	1.69	0.58	2.30	0.69	2.00	86.42
	<i>Sargassum fluitans</i>	3.33	3.33	1.00	3.33	1.00	3.33	100.00

Diet in relation to fish size and marine seasons

Fish were grouped into two categories: juvenile individuals (n = 59) with sizes of 154–256 cm, DW and adults (n = 157) with sizes of 257–371 cm, DW for *M. tarapacana*. In *M. japonica*, sizes of juveniles (n = 32) ranged between 80–177 cm DW while adults (n = 178) ranged in sizes of 177–369 cm DW. For *M. tarapacana*, the trophic spectrum of juveniles consisted of crustaceans, in particular *S. atlanticus* (%PSIRI = 50) and *P. latiffrons* (%PSIRI = 50), whilst the diet of adults was more diversified, including crustaceans (%PSIRI = 62.95), algae (%PSIRI = 21.12), teleost fishes (%PSIRI = 9.18), insects (%PSIRI = 5.81) and molluscs (%PSIRI = 0.93) (Table 3). Relatively to *M. japonica*, the juveniles as well as adults consumed mainly crustaceans (PSIRI = 100% for juveniles and 94.64% for adults), followed by the algae of the genus *Sargassum* (PSIRI = 5.36) in adult individuals (Table 4). The Morisita-Horn index indicated a high trophic overlap between size groups for *M. tarapacana* ($C\lambda = 0.766$) and *M. japonica* ($C\lambda = 0.939$), indicating that both species preferentially feed on the same preys.

As for marine seasons, the prey was similar in both seasons, but the proportions varied considerably (Table 3). During the cold season, the trophic spectrum of *M. tarapacana* consisted mainly of *S. atlanticus* (%PSIRI = 41.16), *P. latiffrons* (%PSIRI = 19.05), *S. natans* (%PSIRI = 8.82%), *R. remora* (%PSIRI = 7.14), fish remains (%PSIRI = 4.76) and *S. fluitans* (%PSIRI = 2.31). The same prey items were found in the warm season, with variations in the order of importance: *S. atlanticus* (%PSIRI = 54.84), *P. latiffrons* (%PSIRI = 17.88), *S. natans* (%PSIRI = 8.03%), *S. fluitans* (%PSIRI = 9.49), *S. aurita* (%PSIRI = 3.08) and fish remains (%PSIRI = 3.23). The other teleost families and Molluscs were incidental foods. Concerning *M. japonica*, Crustaceans including *S. atlanticus* and *P. latiffrons* represented 100% of the PSIRI in warm season (Table 4). During this season, *S. atlanticus* (%PSIRI = 66.67) was the preferential prey, while *P. latiffrons* (%PSIRI = 33.33%) was eaten as secondary prey. In cold seasons (n = 47), apart *S. atlanticus* and *P. latiffrons* which constituted 63.83% and 31.53% of the %PSIRI, the algae of the genus *Sargassum* were lightly encountered in stomachs (%PSIRI = 4.64). The Morisita-Horn indices calculated separately for *M. japonica* ($C\lambda = 0.997$) and *M. tarapacana* ($C\lambda = 0.95$) indicated a high level of trophic overlap between the warm and cold seasons even if no significant difference was observed in diet.

Table 3. Diet composition based on sex, marine seasons and size class groups of *Mobulatarapacana* captured from April 2020 to May 2022 in waters Ivoirian.

Item prey	Prey-specific relative importance index (%PSIRI)					
	Males	Females	Cold	Warm	Juveniles	Adults
Crustaceans	65.42	73.92	60.21	74.15	100.00	62.95
<i>Sergestes. atlanticus</i>	47.29	52.88	41.16	54.84	50.00	50.71
<i>Pseudeuphausia. Latiffrons</i>	16.57	21.04	19.05	17.88	50.00	10.20
<i>Portunussayi</i>	1.56	0.00	0.00	1.43	0.00	2.04

Algae	20.57	9.59	11.13	17.52	0.00	21.12
<i>Sargassum natans</i>	8.64	5.42	8.82	8.03	0.00	12.96
<i>Sargassum fluitans</i>	11.93	4.17	2.31	9.49	0.00	8.16
Teleosts	6.87	13.23	13.6	7.96	0.00	9.18
<i>Priacanthus arenatus</i>	0.64	0.46	1.69	0.00	0.00	0.73
<i>Remora remora</i>	0.00	6.25	7.14	0.00	0.00	2.04
<i>Sardinella aureata</i>	2.55	0.00	0.00	3.08	0.00	0.58
<i>Chloroscombrus chrysurus</i>	0.00	1.46	0.00	0.89	0.00	0.71
<i>Trichiurus lepturus</i>	0.82	0.00	0.00	0.76	0.00	0.59
<i>Fish (ind)</i>	2.86	5.06	4.76	3.23	0.00	4.54
Molluscs	0.00	1.87	1.47	0.38	0.00	0.93
<i>Sepia officinalis</i>	0.00	0.58	0.00	0.00	0.00	0.29
Bivalves	0.00	1.29	1.47	0.38	0.00	0.63
Insect (ind)	7.14	1.39	13.59	0.00	0.00	5.81

Table 4. Diet composition according to marine seasons and size class groups of *Mobula japonica* captured from April 2019 to May 2021 in Ivorian waters.

Item prey	Prey-specific relative importance index (%PSIRI)				Feeding strategy
	Cold	Warm	Juveniles	Adults	
Crustaceans	95.36	100.00	100.00	94.64	The modified
<i>Sergestes atlanticus</i>	63.83	66.67	50	64.29	
<i>Pseudeuphausia latifrons</i>	31.53	33.33	50	30.36	
Algae	4.64	0.00	0.00	5.36	
<i>Sargassum natans</i>	2.51	0.00	0.00	1.79	
<i>Sargassum fluitans</i>	2.13	0.00	0.00	3.57	

Costello Diagram constructed (Fig. 2) established crustaceans, in particular, *S. atlanticus* and *P. latifrons* as the main food item of both species in the coastal waters of Côte d'Ivoire. In other words, these two species display specialist feeding behaviour that focuses on *S. atlanticus* and *P. latifrons*. These species appeared in the upper right area of the diagram, whereas most of the other prey species are located in the left corner or close to the vertical axis, indicating low values for prey importance. These low frequency values suggest that almost all the other species are rare or unimportant prey and are therefore consumed by few individuals.




Figure 2. Costello Diagram modified by Amundsen et al. [24] described feeding strategy and niche width of *Mobulatarapacana* and *Mobulajaponica* from EEZ Côte d'Ivoire. The points represent different prey types in the stomach contents. *Sergestes atlanticus* (*S. atl*), *Pseudeuphausialatifrons* (*P. lat*), *Portunussayi* (*P. say*), *Sargassum natans* (*S. nat*), *Sargassum fluitans* (*S. flu*), *Priacanthusarenatus* (*P. are*), *Remora remora* (*R. rem*), *Sardinelaureta* (*S. aur*), *Chloroscombrus chrysurus* (*C. chr*), *Trichiuruslepturus* (*T. lep*), *Sepia officinalis* (*S. off*), Insects indeterminate (*I. ind*) Bivalves indeterminate (*B. ind*).

DISCUSSION

The value of the vacuity index obtained in this study was very high (over 70%) for both species, indicating that these species can be included in the category of relatively low-feeding species. The high number of empty stomachs is thought to be due to several factors, one of which is the availability of food in nature. The prey eaten by both species were shrimps, which have the characteristic of being digested quickly. In fact, all stages of digestion were observed in the stomachs, although most of them were in an advanced or well-digested state. In addition, Shibuya et al. [26] stated that the high empty stomach could be caused by the fishing time that is not following the activities of fish that are looking for food. This high incidence of empty stomachs may also reflect a long delay between capture and examination. In fact, the duration of the tide was 4–5 days and the fishing activities took place at night, with sets were performed at 17h00, captures occurred habitually between 00h00 and 05h00 and landings occur between 07h00 and 16h00. These results would suggest that most of specimens had fed at least 10 hours before being captured. On the other hand, the hooks used in association to driftnet by fishermen to attract rays cause high stress at the time of capture often resulting in the regurgitation of stomach contents or the whole stomach of several individuals during the data collection. This vomiting reflex is a protective mechanism for the bulk ejection of noxious material which is common in vertebrates [27].

The diet composition of *M. tarapacana* is more diverse, consisting of crustaceans (68.62%), algae of the genus *Sargassum*, teleost fishes, molluscs and insects. On the other hand, the dietary spectrum of *M. japonica* is essentially made up of crustaceans (94.70%) and algae of the genus *Sargassum*. As we can see, both species had a diet based mainly on crustaceans. Stevens[28] noted that in the East Atlantic areas of the Indian Ocean, *M. tarapacana* appear to be specialized in catching small fish, whereas *M. japonica* feeds mainly on zooplankton and micronectonic crustaceans such as euphausiid *P. latifrons* and sergestid shrimps *Acetes intermedius* and *Lucifer* spp., copepods and other rare prey items [14-29]. This suggests that the prey dominance in the diet varies according to space and their abundance in the environment[30]. Based on the quantitative analysis of the diet through the %PSIRI, the diet of both species was made up mainly of crustaceans, in particular the Shrimp *S. atlanticus* and *P. latifrons*. Their importance in the diet of these Mobulid species could be due to their abundance in the environment, as [31]reported high concentrations of shrimp along the coast of Côte d'Ivoire. The presence of teleost fishes, crabs, molluscs, cephalopods and insects attests the ability of *M. tarapacana* to adjust its diet according to the availability of some preys, relying on seasonal variations or environmental conditions to optimise its diet. Some previous studies carried out on *M. japonica* reported a low diversity of ingested prey composed exclusively of pelagic crustaceans such as euphausiids and other zooplankton species [1,13,14,29]. Two algae of the genus *Sargassum* (*S. natans* and *S. fluitans*) which seasonally invade marine waters [32] were identified in stomachs. The hypothesis that could explain the presence of algae in stomachs is linked to the Mobulidae's hunting and predation strategy, which seems to be closely linked to the way they take food, by propelling themselves to capture small schooling fish (Stevens [29]) or by extracting prey buried in the carpet formed by this algae [33-34]. According to Coston-Clements et al.[35], *Sargassum* is a group of brown algae that provides food, refuge, and breeding ground for many marine animals, such as turtles, crabs, shrimp, fish, and seabirds. In this condition, they can be considered as incidental items. The presence of the shrimps (*S. atlanticus* and *P. latifrons*) and the crab *Portunussayi* indicated that this species migrates between pelagic and benthic zones.

The feeding habits of both species did not change significantly according seasons and the Morisita-Horn index 0.997 (*M. japonica*) and 0.950 (*M. tarapacana*) indicated a similarity of diet. However, the proportions and relative importance of each prey type vary. Thus, the diet was richer in the cold season, due to variations in the abundance of groups of benthic and pelagic fauna constituting the majority of prey. This study also showed ontogenetic variations in the feeding habits of both species. In fact, the trophic spectrum of juveniles consisted of crustaceans (essentially *S. atlanticus* and *P. latifrons*), whilst that of adults was more diversified, including in addition to crustaceans, algae, teleost fishes, insects and molluscs. As we can see, as fish sizes increase, they gradually add other prey to their diet. This is consistent with Jacobsen and Bennett [30], who found that the diet of rays varies depending on body size, morphology, and sexual maturity status.

Varghese et al. [36], reported that prey diversity in the adult diet could also be associated with the increase of energy requirements that increases with fish sizes. The feeding strategy involves supplementing the diet with more mobile, calorie-rich prey; reducing the time spent foraging and devoting the energy gained to reproductive activity. The Costello diagram indicates that these *Mobula* species are selective predators or specialise in eating shrimp species throughout the year.

4. CONCLUSION

This study highlights the feeding habits of *M. tarapacana* and *M. japanica*, two species of *Mobula* rays regularly captured in the coastal waters of Côte d'Ivoire. Both species feed mainly on benthic shrimps of the Sergestidae and Euphausiidae families. However, the diet of *M. tarapacana* is more diversified, including fish, Cephalopods, Crabs, Insects and Bivalves. Two algae of the genus *Sargassum* (*S. natans* and *S. fluitans*) were identified in stomachs, indicating that, these species hunt their prey by extracting prey buried in the mat formed by these algae. These species can be considered as carnivorous predators targeting shrimp species. These current results provide a useful contribution to the local conservation and management of *Mobulidae* in Côte d'Ivoire.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that no generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

REFERENCES

1. Couturier LI, Marshall AD, Jaine FR, Kashiwagi T, Pierce SJ, Townsend KA, et al. Biology, ecology and conservation of the *Mobulidae*. *J Fish Biol.* 2012;80: 1075–1119.
2. Croll D A, Dewar H, Dulvy NK, Fernando D, Francis MP, Galván-Magaña F et al. Vulnerabilities and fisheries impacts: The uncertain future of manta and devil rays. *AquatConserv.* 2016; 26:562–575.
3. Francis MP, Jones EG. Movement, depth distribution and survival of spinetail devil rays (*Mobula japanica*) tagged and released from purse-seine catches in New Zealand. *AquatConserv.* 2017;27(1):219–236.
4. White WT, Bazinet AL, Yang L, Corrigan S, Swofford DL, Naylor GJ et al. Phylogeny of the manta and devil rays (*Chondrichthyes: Mobulidae*), with an updated taxonomic arrangement for the family. *Zool J Linn Soc.* 2017; 182(1): 50–75.
5. Kouamé KY. Caractérisation de la pêche artisanale aux filets maillants et bioécologie du requin peubleue *Prionace glauca* (Linnaeus, 1758) de la Zone Economique Exclusive (ZEE) de la Côte d'Ivoire. Mémoire de Thèses de doctoratenHydrobiologie, Université Félix Houphouët-Boigny (Abidjan, Côte d'Ivoire). 2020; 200 p. French.

6. Wijayanti F, Abrari MP, Fitriana N. Keanekaragamanspesies dan status konservasi ikan pari di tempatpelelangan ikan Muara Angke Jakarta Utara [Species diversity and conservation status of stingrays in Muara Angke Fish Auction, North Jakarta]. *JurnalBiodjati*. 2018;3(1) :23–55.
7. Garcia VB, Lucifora LO, Myers RA. The importance of habitat and life history to extinction risk in sharks, skates, rays and chimaeras. *Proc R Soc Lond B Biol Sci*. 2008; 275: 83–89.
8. White W.T., Clark T.B., Smith W.D. & Bizzarro J.J. *Mobulajapanica*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2011.2. Retrieved from www.iucnredlist.org. 2006.
9. Marshall A, Barreto R, Carlson J, Fernando D, Fordham S, Francis MP et al. *Mobulamobular* (amended version of 2020 assessment). The IUCN Red List of Threatened Species 2022: e.T110847130A214381504. <https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T110847130A214381504.en>
10. Du Buit MH. Diet of hake (*Merluccius merluccius*) in the Celtic Sea. *Fish Res*.1996; 28(4):381–394.
11. Costello M.J. Predator feeding strategy and prey importance: a new graphical analysis.*J Fish Biol*. 1990; 36: 261-
12. Stewart JD, Hoyos-Padilla EM, Kumli KR, Rubin RD. Deep-water feeding and behavioural plasticity in *Manta birostris* revealed by archival tags and submersible observations. *Zoology (Jena)*. 2016;119(5):406–413.
13. Notarbartolo di Sciara G. A revisionary study of the genus *Mobula* Rafinesque, 1810 (Chondrichthyes: Mobulidae) with the description of a new species. *Zool. J Linn Soc*. 1987; 91(1): 1–91.
14. Sampson L, Galván-Magaña F, De Silva-Dávila R, Aguiñiga-García S, O'Sullivan JB. Diet and trophic position of the devil rays *Mobulathurstoni* and *Mobulajapanica* as inferred from stable isotope analysis. *J Mar Biol Assoc U K*. 2010; 1:1–1.
15. Diaha C, N'da K, Soro Y. Diet of *Scomberomorus tritor* (Cuvier, 1831) in the Gulf of Guinea. *Int J Biol Chem Sci*. 2010; 4(3): 669-680.
16. Morlière A, Rebert JP. Hydrological study of the Ivorian continental shelf. Scientific documents from the Oceanographic Research Center, Abidjan.1972; 3(2):1–30.
17. Fischer W, Bianchi G.FAO species identification sheets for fishery purposes. Eastern Central Atlantic, fishing areas 34 and 47. 1981; 324 p.
18. SchneiderW. FAO species identification sheets for fishery purposes: Field guide to the commercial marine resources of the Gulf of Guinea. Prepared and published with the support of the FAO regional office for Africa. FAO, Rome. 1992; 268 p.

19. Carpenter KE, De Angelis N. The Living Marine Resources of the Eastern Central Atlantic. Volume 1: Introduction, Crustaceans, Chitons and Cephalopods. Rome: Food and Agriculture Organization of the United Nations. 2014;683.
20. Brown S, Bizzarro J, Cailliet G, Ebert D. Breaking with tradition: redefining measures for diet description with a case study of the Aleutian skate *Bathyrhajaaleutica* (Gilbert 1896). *Environ Biol Fishes*. 2012;95(1):3–20.
21. Rosecchi E, Nouaze Y. Comparaison de cinq indices utilisés dans l'analyse des contenus stomacaux. *Rev. Trav. Inst Pêches Marit*. 1987; 49:11–123.
22. Morisita M. Measuring of the dispersion and analysis of distribution patterns. *Memories of the Faculty of Science, Kyushu University. Series E Biology*. 1959; 2:215–235.
23. Smith PE, Zaret MT. Bias in estimating niche over-lap. *Ecology*. 1982; 63 (5): 1248–1253.
24. Costello MJ. Predator feeding strategy and prey importance: a new graphical analysis. *J Fish Biol*. 1990; 36:261–263.
25. Amundsen PA, Gabler HM, Staldvik FJ. A new approach to graphical analysis of eating strategy from stomach contents data—modification of the Costello (1990) method. *J Fish Biol*. 1996; 48:607–614
26. Shibuya A, Araújo ML, Zuanon JA. Analysis of stomach contents of freshwater stingrays (Elasmobranchii, Potamotrygonidae) from the middle Negro River, Amazonas, Brazil. *Panam J Aquat Sci*. 2009; 4(4):466–475.
27. Sims DW, Andrews PL, Young JZ. Fish behaviour: Stomach rinsing in rays. *Nature*. 2000; 404(6778):566–566.
28. Stevens G. Field guide to the identification of Mobulids ray (Mobulidae): Indo-West Pacific. The manta trust. 2011; 19 p.
29. Shirlamaine IG, Ephrime BM, Ken-Ichi H, Satoru T, Shuhei N. Feeding Habits of *Mobula japonica* (Chondrichthyes, Mobulidae) in Butuan Bay, Mindanao Island, Philippines. *Sci Diliman*. 2018; 30 p.
30. Jacobsen IP, Bennett MB. A Comparative Analysis of Feeding and Trophic Level Ecology in Stingrays (Rajiformes; Myliobatoidei) and Electric Rays (Rajiformes: Torpedinoidei). *PLoS ONE*. 2013; 8(8): 1–10.
31. Troadec UP, Garcia S, Petit P. L'exploitation des crevettes en Côte-d'Ivoire, de solides raisons d'espérer. *Pêche Marit*. 1969, 1095: 434–436.
32. Sankaré Y, Komoe K, Aka KS, Fofie, NB, Bamaba A. Répartition et abondance des sargasses *Sargassum natans* et *Sargassum fluitans* (Sargassaceae, Fucales) dans les eaux marines ivoiriennes (Afrique de l'Ouest) *Int J Biol Chem Sci*. 2016; 10(4): 1853–1864. French.
33. West LE. Habitat location and selection by the Sargassum Crab *Portunussayi*: the role of sensory cues. Thesis State University System of Florida. 2012.

34. Brown and Barry B. Sargassum Swimming Crab, Portunussayi, Curacao. 2011.
35. Coston-Clements L, Settle LR, Hoss DE, Cross FA. Utilization of the Sargassum habitat by marine invertebrates and vertebrates - a review. NOAA Technical Memorandum. 1991;30 p.
36. Varghese SP, SomvanshiVS, Dalvi R.S. Diet composition, feeding niche partitioning and trophic organization of large pelagic predatory fishes in the eastern Arabian Sea. Hydrobiologia. 2014;736(1):99–114.

UNDER PEER REVIEW