

***Clitoria ternatea* (Aparajita) : A Versatile Medicinal Plant with Diverse Benefits**

**ABSTRACT**

*Clitoria ternatea* Linn., (aparajita/shankapushpi; family Fabaceae), is a traditional medicinal plant with diverse applications- both medicinal and commercial. It is a perennial twinning herb with prominent flowers. It is native to tropical Asia. The aim of review is to provide in-depth information on the phytochemistry along with commercial and medicinal uses of *C. ternatea*. Exhaustive review of available literature was carried out to collect and critically evaluate scientific information available on the plant. According to available literature, the plant possesses diverse health benefits due to which it is used in traditional as well as modern systems of medicine. Traditionally it is used as antidote for snake bite, scorpion bite, memory enhancer and anxiolytic agent and in modern system of medicine it shows various pharmacological activities that includes antidepressant, antioxidant, anti-inflammatory, anti-pyretic, analgesic etc. The activities are attributed to the presence of varied constituents predominantly the unique acylated anthocyanins – ternatins, cyclotides, phenolic compounds etc. Commercially it is valued as a natural dye and food colorant because of the presence of acylated anthocyanins in it. This plant has marked antioxidant and antiaging properties and hence it is gaining popularity for making cosmetics and cosmeceuticals. It is also used for protecting crops in agrotechnology. This review highlights the numerous medicinal properties, commercial uses, as well as its phytochemical investigation prove multifaceted potential of this plant. Further directions for developing the plant and its products as effective and stable medicine, nutraceutical and cosmeceuticals are also suggested.

**Keywords:** Acylated anthocyanins, *Clitoria ternatea*, Cosmetics, Food colorant, Medicinal uses, Shankapushpi, Traditional medicine.

## 1. Introduction

Globally medicinal plants are employed to impart health and to cure several illnesses. Plant products are used in both modern and traditional systems of medicine. The Ayurvedic medicinal system is notable for the impact of its "Medhya Rasayana" collection of herbal remedies on the nervous system. "Medhya drugs" are described as having the capacity to improve cognition in Ayurvedic literature.

*Acorus calamus*, *Areca catechu*, *Celastrus panniculatus*, *Centella asiatica*, *Clitoria ternatea*, *Tinospora cordifolia*, and *Withania somnifera* a few examples said to influence the neurological system. There is greater emphasis in the current time to generate evidence to support traditional claims hence many plants are being investigated in detail to assess their medicinal potential and to identify the active ingredients.

One important "Medhya Rasayana"- *Clitoria ternatea* (commonly called aparajita/shankapushpi; family Fabaceae). This plant has been examined from various perspectives showing its phytochemistry, pharmacological as well as commercial uses. It is a perennial climber. This ornamental plant is valued due to its attractive colour of the flower (Gomez and Kalamani., 2003). It is a self-pollinated plant with diploid chromosome number (2n=16) (J. B. Morris., 2009). Because of the presence of high protein content, it is also used as fodder for the cattle. It is a drought tolerant plant (Patel and Mishra., 2023).

Distributed in many tropical countries like in Caribbean, India, Madagascar, Philippines, South and Central America, and other tropical Asian countries, Two types of varieties of this flower, one is white flower, and blue flower.

The aim of this review article is to highlight the commercial and medicinal applications of *C. ternatea*.

## 2. Taxonomic classification


*C. ternatea* is known by different name in different countries some of the synonyms are presented in table 1.

**Table 1: Synonyms of *C. ternatea***

Language	Synonyms
Brazilian	Cunha
Chinese	Lan hu die
Hindi	Koyala
Indonesian	Bunga biru, tembang telang
Malaysia	Telang
Philippines	Pokindang
Portuguese	Fulacriquacunha~ fula criqua
Sanskrit	Shankapushpi, Aparajita, Saukarnika, Supuspi, Vishnukranta
Spanish	Clitoria azul
Sudan	Butterfly pea, Kordofan pea
Thai	Dangchan
Turkish	Mavi kelebek sarması
<b>References:</b> Mukherjee <i>et al.</i> , 2008; Jayaraj <i>et al.</i> , 2021.	

### 3. Plant cultivation

*C. ternatea* is a perennial ornamental plant which can be cultivated in gardens and grows wild as well.



The plant needs some requirements for growth which are presented in table 2.

**Table 2: Requirements for growth of the plant.**

Requirements for growth	
<b>Altitude</b>	0-1600 metres
<b>Lifecycle of flower</b>	Flowering takes 4 to 5 weeks
<b>pH</b>	5.5 to 8.9
<b>Season</b>	Perennial
<b>Soil Type</b>	Variety of soils including calcareous soils
<b>Spacing</b>	15 - 30 cm (slight row gap is preferred)
<b>Sowing depth (damp soils)</b>	2.5 - 6.5 cm
<b>Sunlight/Shade</b>	Prefer full sunlight, but occasionally partially shaded areas
<b>Temperature Range</b>	19-28°C
<b>References:</b> Gomez and Kalamani., 2003; Chahal <i>et al.</i> , 2010; Multisona <i>et al.</i> , 2023; Patel and Mishra., 2023	

In American, recently this plant cultivated in wet tropical climate, according to ecology this plant usually require direct sunlight, but occasionally partially shaded areas are preferable. For the germination, seeds are soaked in water overnight after these seeds will germinate within 1 or 2 weeks and flowering takes 4 to 5 weeks. The stem method is also used for the cultivation of *C. ternatea* (Multisona *et al.*, 2023).

This plant is drought sensitive, recent study in Malaysia was conducted for comparing the germination rate in drought stress and normal conditions and the germination rate was reduced in drought stress encountered plant as compared to normal condition plants. The germination rate in drought stress and normal conditions was 42.5% and 56.25% respectively (Jamil *et al.*, 2018).

### 3.1. Micropropagation

Micropropagation of plant material is required to fulfill the demand of the natural product in the market. In-vitro culture techniques are the alternative technique to minimize the consequence of extinction of the plant (Malabadi and Nataraja, 2001). Studies of *in-vitro* propagation of the plant are summarized in table 3.

**Table 3: Conditions for *in-vitro* propagation**

Explant	Types of media	Further supplementation	Observations	Reference
Shoot	Semisolid Murashige and Skoog (MS) growth medium	Sucrose (3% w/v), and different concentrations of growth hormones [6- benzylaminopurine or kinetin] + 1-naphthaleneacetic acid or Indole-3-acetic acid.	No growth of the explants observed when explants were cultured on the media without auxin or Cytokinin. Supplementation with different concentrations of growth hormones promoted multiplication of shoots.	Rout., 2005
Shoot	MS media	Sucrose (3% w/v), Agar (0.8% w/v) and [6-benzyl aminopurine, 6- furfuryl aminopurine and thidiazuron]	MS medium supplemented with 2.22 $\mu$ M BAP showed the maximum regeneration of shoots.	Mishra <i>et al.</i> , 2019

#### 4. Plant Description

In different regions and systems of medicines, diverse parts of the plant are employed (figure 1; Tamanna., 2023).




Figure 1: Different parts of *C. ternatea*

##### 4.1. Flowers



- It is the attractive part of the plant. These are available in single or paired form with various colors like mauve, white, dark and light blue.
- The pedicels and bracteoles can grow up to 4-9 mm and 12 m long respectively and its corolla consists of one standard petal, 2 keels and 2 wing petals. The standard petal is the largest one among all the petals (Multisona *et al*, 2023; Ashok and Geetha., 2012).
- Ashok and Geetha., 2012 suggest that typically the flower is bilateral, but the petal mutants are also present in nature that results in radial appearance of the flower. The colour variability of the flower is because of the presence of flavonoids which was reported by Kazuma *et al.*, 2003.

##### 4.2. Leaves

- The leaves show pinnate venation with 7 leaflets (Oguis *et al.*, 2019).
- The terminal and base leaflets are bigger and smaller respectively.
- The dorsiventral structure is observed when the transverse section of the leaves is carried out.
- Calcium oxalate crystals are present along the veins prismatic. The palisade ratio and vein-islet number is 6.0 and 7.5 respectively (Jagbir Chahal *et al.*, 2010).
- Leaf blade have linear trichomes on both sides. The shape of the lamina is ovate, and its surface is smooth with hairy texture (Rampalli & Bindu., 2022).

Organoleptic characters of *C. ternatea* flowers and leaves are presented in table 4 (Tamanna., 2023).

Table 4: Organoleptic characters of *C. ternatea* flowers and Leaves

Plant	Plant Part	Picture	Organoleptic Parameters	Observation
<i>C. ternatea</i>	Flowers	 <p>Figure 2: <i>C. ternatea</i> Flowers</p>	Color	Indigo
			Odour	Odour less
			Shape	Funnel-shaped
			Size	Length:1-1.5 in; Width: 2-3 cm
			Taste	Woody
			Season	Perennial
			Inflorescence	Solitary
	Sex	Bisexual		
	Leaves	 <p>Figure 3: <i>C. ternatea</i> Leaves</p>	Color	Green
			Odour	Characteristic
			Shape	Pinnate
			Size	Length: 4-6 in; Width: 5-6 cm
			Taste	Bitter and acrid
			Type	Compound
Apex			Emarginate	
Base	Symmetrical			
Margin	Entire			

**4.3. Pods:** The pods are flat, linear, olive, brown and black in colour, about 5-7 cm long containing 6-10 seeds in each pod (Suarna and Wijaya., 2021).

**4.4. Seeds:** The seeds can be germinated by immersing them in water over night. After that, germination happens in 1-2 weeks, and blooming happens in 4 weeks. The plant needs full sunlight or partially shaded area to grow properly (Multisona *et al.*, 2023).

**4.5. Stems:** Stems are 3-5 meters long, hairy, bald, or occasionally upright. It is long, slender, and flexible in nature. It is light green to brownish in colour (Jain and Shukla., 2011).

**4.6. Roots:** The root system of this plant is a strong taproot system (Multisona *et al.*, 2023). Nodules of the root shows symbiotic association with nitrogen fixing bacteria, so it has capability to fix atmospheric nitrogen in them (Suarna and Wijaya., 2021). It has a brown colour with a bitter taste and characteristic odour (Taur and Patil., 2010).

## 5. Traditional Uses:

The plant is valued greatly for its varied therapeutic benefits in different traditional systems and folk medicine. Medicinal benefits of different parts of *C. ternatea* are presented in table 5.

**Table 5: *C. ternatea* Traditional Uses**

Plant Part	Uses	References
<b>Flower</b>	<ul style="list-style-type: none"> <li>The paste of flower is used for the treatment of eye infection and headache.</li> <li>Flowers are also used as an antidote for snakebites.</li> </ul>	Alok <i>et al.</i> , 2015
<b>Leaves</b>	<ul style="list-style-type: none"> <li>When a headache or swelling of a nearby gland occurs, juice from the leaves is combined with salt and placed around the ears to reduce discomfort.</li> <li>Juice of leaves is used as an antidote against snakebite.</li> <li>To treat swelling joints and used as poultices.</li> </ul>	Alok <i>et al.</i> , 2015
<b>Seeds</b>	<ul style="list-style-type: none"> <li>Used to treat colic, dropsy, swelling joints and enlargement of abdominal viscera.</li> <li>It also possesses laxative, mild emetic and vermifugal properties.</li> <li>Used as green manure and as an antidote for poisons.</li> </ul>	Ashraf, <i>et al.</i> , 2024
<b>Stem</b>	<ul style="list-style-type: none"> <li>Used as antidote for snake bite and scorpion.</li> <li>It acts as brain tonic because of the presence of some phytochemical, and it is also useful in urinary problems, throat, and eye related problems.</li> </ul>	Sarma <i>et al.</i> , 2023
<b>Roots</b>	<ul style="list-style-type: none"> <li>Ascetics, epilepsy, enlargement of the abdominal viscera, skin diseases, and sore throat.</li> <li>Used as diuretic, laxative, mind tonic and purgative.</li> <li>Serves to treat different ailments such as constipation, dyspepsia, eye conditions, eye conditions, enlarged abdominal organs, fever.</li> <li>Rheumatism and ear problems are also treated using roots in the form of powder or decoction.</li> <li>White variety flowered root juice is blown up the nose as a treatment for hemicrania.</li> </ul>	Mukherjee <i>et al.</i> , 2008; Alok <i>et al.</i> , 2015; Ashraf, <i>et al.</i> , 2024

### 5.1. Other uses:

- *C. ternatea* is locally called as Sangu Pushpam in Kancheepuram district of Tamil-Nadu, root powder with water is consumed to get relief from indigestion, eye diseases and headache.
- In Chhattisgarh, the fresh root bark (powder) acts as diuretic when administered with warm milk for 2 weeks and the seed act as purgative when administered with warm water once a day for the period of 3 days.
- In Irulas of the Kodiakkarai, *C. ternatea* flower paste is used to treat eye infection and headache.
- According to Dharampuram Taluk, Tamil-Nadu the seed powder of *C. ternatea* used for the treatment of constipation when mixed with pepper.
- According to Uttara Kannada district of Karnataka:
  - ✓ The root juice is used to reduce fever when it is applied on the body.
  - ✓ The root ash is involved in facial care.
  - ✓ The burned seeds fume inhalation is useful to get relief from hiccups.
  - ✓ The infusion of leaves are used for the treatment of eruptions (Jagbir Chahal *et al.*, 2010).
  - ✓ In Malaysia, the color of the flower and leaves are used as colorant (Ashraf, *et al.*, 2024).
  - ✓ In Philippines, different parts of the plant are consumed as vegetables (Ashraf, *et al.*, 2024).
  - ✓ In Saudi Arabia, flowers of the plant are used as caffeine-free herbal tea (Gomez and Kalamani., 2003).

### 6. Nutritional content:

There are various amounts of nutrients present in plants. Some of the nutrients of *C. ternatea* present in different parts of the plant are mentioned in table 6.

**Table 6: Nutritional composition of *C. ternatea*.**

<b>Plant part</b>	<b>Nutrients value</b>	<b>References</b>
<b>Flowers</b>	Iron = 0.14 mg/g Magnesium = 2.23 mg/g Potassium = 1.25 mg/g Protein = 0.32 % Sodium = 0.14 mg/g Zinc = 0.59 mg/g	Jeyaraj <i>et al.</i> , 2021
<b>Leaves</b>	Fiber = 8.45 % Fat = 5.5 % Protein = 14.99 % Reducing sugar = 0.036 % Starch = 0.038 %	Swati <i>et al.</i> , 2014
<b>Seeds</b>	Oil = 10 % Protein = 25 to 38 % Sugar = 5 %	Turnos., 2021
<b>Stem</b>	Sugar = 112 ± 0.30 % Starch = 53 ± 0.47 % Protein = 39 ± 0.13 % Phenol = 37 ± 0.56 % Lipids = 18 ± 0.35 % (mg/gdw = milligram per gram dry weight)	Shekhawat & Vijayvergia., 2010
<b>Roots</b>	Sugar = 102 ± 0.59 % Starch = 42 ± 0.35 % Protein = 21 ± 0.49 % Phenol = 43 ± 0.13 % Lipids = 41 ± 0.14 % (mg/gdw = milligram per gram dry weight)	

## 7. Phytoconstituents:

The plant is rich in primary and secondary metabolites. Different parts contain varied phytoconstituents presented in table 7.

**Table 7: Phytoconstituents reported from different parts of *C. ternatea*.**

Plant Part	Phytochemical constituents	References
Flower	Alkaloids, Carbohydrates, glycosides, phytosterols, saponins and tannins.	Sarma <i>et al.</i> , 2023
Leaves	Alkaloids, flavonoids, glycosides, reducing sugar and Steroids	
Seeds	Alkaloids, amino acids, glycosides, mucilage, proteins, resins tannins	
Stem	Carbohydrates, saponins, fixed oil, flavonoids, phenols	Jain and Shukla., 2011
Roots	Flavonoids, resins, starch, tannins, taraxerone and taraxerol.	Alok <i>et al.</i> , 2015

### 7.1. Anthocyanins:

The biologically and commercially important constituents includes anthocyanins, cyclotides, flavonoids, steroids etc. Cyclotides are the stable peptides which are present in the whole plant. Butelase 1 is a cyclotide processing enzyme with having capability of ligating various range sizes (26 to >200 residues) of peptides which further used as cyclotide containing pesticide (Oguis *et al.*, 2019). Anthocyanins, which are flavonoid chemicals give plants various colour, from mild pink to deep blue and purple (Cavalcanti *et al.*, 2011). Because of the intense blue colour of the petals, it has been determined that the primary phytochemicals in the *C. ternatea* flower is anthocyanins.

The evaluation of anthocyanin content: Amount of anthocyanin present in *C. ternatea* can be evaluated by formula:

$$\gamma = (A.M.F) / (m. \epsilon .l)$$

where  $\gamma$  = Anthocyanin mass concentration.

A = Absorbance.

M = Anthocyanin's molar mass, 500.8 gmol<sup>-1</sup>

F = Dilution ratio of *C. ternatea* flower.


V = Extraction solution volume.

m = Mass of *C. ternatea*,  $\epsilon$  is attenuation coefficient, 27300<sup>-1</sup> mol<sup>-1</sup> cm<sup>-1</sup> and the width of cuvette, 1 cm (Syafa'atullah *et al.*, 2020).


The anthocyanins extracted from this plant, are ternatins A1-A3, B1-B4, C1-C4, and D1-D3. These are blue acylated anthocyanins that are derived from delphinidin. Ternatins structure is identified as malonylated delphinidin 3,3',5'-triglucosides, having 3',5'-side chains with have ability alternate D-glucose and p-coumaric acid units because they are highly acylated with p-coumaric acid (Terahara *et al.*, 1998; Mukherjee *et al.*, 2008). All ternatins produced 6-malonyl-glucose while hydrogen peroxide oxidation which indicates that malonic acid is attached to 3-glucoside of each ternatin. Anthocyanins are extracted by polar solvent because they have the affinity to extract them. Lower pH was found to the optimal for the extraction of anthocyanins (Ludin *et al.*, 2018; Mauludifia *et al.*, 2019). Depending on the pH, anthocyanin's colour changes from acid to base. The blue colour anthocyanins appear in the pH range of 5–7 by the quinonoid base form which can be easily degraded. The stableness of ternatins can be improved by high degree of acylation and glycosylation (Fu *et al.*, 2021).

The stability of anthocyanins will be calculated by using formula:


$$\text{Absorbance decay percent (\%)} = [(A_0 - A_t)/A_0] \times 100\%$$




(a)




(b)




(c)




(d)



(e)




(f)



(g)

**Figure 4:** Anthocyanins present in *C. ternatea*: (a) Ternatin A1, (b) Ternatin A2, (c) Ternatin B1, (d) Ternatin B2, (e) Ternatin D1, (f) Ternatin D2, (g) Delphinidin-3-O-glucoside


### 7.2. Flavonol Glycosides:



**Figure 5.** Major flavonol glycosides are present in *C. ternatea*: (a) Kaempferol; (b) Quercetin; (c) Myricetin; and (d) Delphinidin-3-malonyl glucoside.

### 7.3. Phytosterols:

Phytosterols are present in the plant (Multisona *et al.*, 2023) are presented in figure 6.



**Figure 6:** Phytosterols in *C. ternatea* (a)  $\beta$ -sitosterol; (b) stigmasterol; (c) taxaxerol; (d) campesterol; (e) sitostanol

### 8. Extraction methods:

Extraction is the essential step for the isolation of valuable phytochemical from the plant material. There are various methods available for the extraction of phytochemicals and these methods are selected based on the nature of the plant material (Handa., 2008). Extraction methods are classified into two types: conventional methods (traditional methods) and non-conventional methods (modern methods) (Jeyaraj *et al.*, 2021). There are various factors which affect the yield of extraction like pH, solvent (polar or non-polar), temperature etc. (Lakshan *et al.*, 2019).

Various methods have been employed to extract phytoconstituents from this plant (Table 8).

**Table 8: Extraction of different compounds from *C. ternatea* by different extraction methods and solvents.**

Extraction method		Solvent	Compound/phytochemical	Yield	Reference
Maceration		Water	Dye strain	45.52 %	Baskaran <i>et al.</i> , 2019
Cold extraction	water	Water	Phenols	185.3 mg/g	Weerasinghe <i>et al.</i> , 2022
Hot extraction	water	Water	Phenols	239.5 mg/g	
			Flavonoids	128.3 mg/g	
Ultrasound assisted extraction		Distilled water	Anthocyanins	-	Chong <i>et al.</i> , 2015
Maceration		Hydro alcohol	Flavonoids	246.6 %	
Microwave assisted extraction (decoction)		Distilled water	Dye strain	48.61 %	Marsin <i>et al.</i> , 2020

Among all the methods extraction assisted with ultrasound (modern method) have proven to be more effective method for anthocyanins extraction.

#### **9. Pharmacological activities:**

Numerous pharmacological studies (*in-vitro* and *in-vivo* investigations) have been carried using various extracts of the plant. **Table 9** sums up the findings of investigations in the last 10 years.

**Table 9: Various pharmacological activities of *C. ternatea*.**

**Antioxidant activity:**

<b>Plant (Part used)</b>	<b>Extract</b>	<b>Dose</b>	<b>Model</b>	<b>Result/ Observations</b>	<b>Reference</b>
White and Blue (roots)	Petroleum ether, Chloroform and Methanol extract	50-600 µg/ml	DPPH assay	Methanol extract of white variety roots show most significant antioxidant activity.	Al-Snafi, 2016; Divya <i>et al.</i> , 2018
White and Blue (leaves)	Petroleum ether, Chloroform and Methanol extract	50-600 µg/ml	Hydroxyl Radical Scavenging Activity	Methanol extract (white flower) leaves show most marked activity.	Al-Snafi, 2016; Divya <i>et al.</i> , 2018
White and Blue (leaves)	Petroleum ether, Chloroform and Methanol extract	50-600 µg/ml.	DPPH assay	Methanol extract of white variety show most significant activity.	Jiji & Muralidharan, 2020
White and Blue (Flowers)	Flower petal extract	400 µg/ml	Oxygen Radical Absorbance Capacity method and DPPH assay	The results of the study showed that petal extract prevent morphological alteration of erythrocytes.	Jiji & Muralidharan, 2020

Flowers	Anthocyanin extract	pH 5-7	DPPH assay	Strong antioxidant effect in the pH range 4–7 and a sharp decrease when the pH exceeded 7.	Fu <i>et al.</i> , 2021
Flowers	Aqueous extract	100 mg/kg	DPPH assay and Bisphenol A	Protective effects in female murine reproductive system with regards to improve the percentage of pregnancy and litter size and potent antioxidant activity.	Goh <i>et al.</i> , 2021
Flowers	Ethanol extract lotion	0.05%, 0.1% and 0.5%	DPPH assay	0.1% lotion showed potent antioxidant activity with IC <sub>50</sub> 37.92 ppm.	Jayanti <i>et al.</i> , 2021
Flowers	Ethanol extract	Different concentration	ABTS, DPPH assay	Ethanol extract exhibits potent antioxidant activity in ABTS and DPPH assay with IC <sub>50</sub> 10.23 ± 0.186; 2.77 ± 0.020, respectively.	Tuan Putra <i>et al.</i> , 2021

Flowers	Aqueous extract	Different concentration	ABTS, DPPH, FRAP and Hydrogen Peroxide (H <sub>2</sub> O <sub>2</sub> ) Scavenging Assay	Strong antioxidant activity in ABTS, DPPH and Hydrogen Peroxide (H <sub>2</sub> O <sub>2</sub> ) Scavenging Assay with (IC <sub>50</sub> : 2.51%, 17.07%, 5.56% and 26.62%) respectively.	Widowati <i>et al.</i> , 2022
Flowers	Fresh, dried flower and <i>C. ternatea</i> flowers + citrus species	Different concentrations	DPPH, Hydrogen Peroxide (H <sub>2</sub> O <sub>2</sub> ) Scavenging Assay	Dried flower showed potent antioxidant activity.	Widowati <i>et al.</i> , 2022

#### Antidiabetic activity:

Plant (Part used)	Extract	Dose	Model	Result/ Observations	Reference
Leaves	Methanol, Water, Petroleum ether and Chloroform extracts	200 -400 mg/kg	Streptozotocin Induced Diabetic Rats.	Methanol extract (200 mg/kg) decreased blood glucose level.	Al-Snafi, 2016
Roots	Alcohol extract	100 mg/kg	Juvenile diabetic rat experimental models.	Alcohol extract prevented the issues which were related to brain hippocampal CA3 area and pancreatic tissue.	Gollen <i>et al.</i> , 2018
Leaves	Ethanol extract	400 mg/kg	Streptozotocin induced model	Decrease in the blood glucose, creatinine, glycosylated hemoglobin, insulin, liver marker enzymes, and urea levels.	Kavitha <i>et al.</i> , 2018

Flowers	Hydro-ethanol extract	400 and 800 mg/kg	Diabetes mellitus and dyslipidemia in rats	Pancreatic CAT, Superoxide Dismutase (SOD) and protein levels were increased. Pancreatic Malondialdehyde (MDA), IL-18 levels, glycogen gene expression of pancreas and IL-6 protein expression of pancreas in DM and dyslipidemia rats were reduced.	Widowati <i>et al.</i> , 2023
---------	-----------------------	-------------------	--	--	-------------------------------

UNDER PEER REVIEW

**Nootropic potential:**

<b>Plant (Part used)</b>	<b>Extract</b>	<b>Dose</b>	<b>Model</b>	<b>Result/ Observations</b>	<b>Reference</b>
Leaves	Ethanol extract	200 and 400 mg/kg	Y-maze; Morris water maze; Radial arm maze	Reduction in cholinesterase activity, reduction in nitric oxide as well as lipid peroxide level and enhances the levels of catalase, superoxide dismutase and glutathione levels.	Talpate <i>et al.</i> , 2014
Roots	-	200 mg/kg and 300 mg/kg	Cerebral hypoperfusion induced memory deficits in a rat model	Inhibition of acetylcholinesterase activity in the frontal cortex and hippocampus of the rat brain.	Damodaran <i>et al.</i> , 2020
	Alcohol extract	50 and 100 mg/kg	Scopolamine induced amnesia model.	AChE activity was significantly decreased leading to increase in the level of ACh content .	Jiji & Muralidharan, 2020
	Aqueous and Hydroalcoholic extracts	100, 300, and 500 mg/kg	100, 300, and 500 mg/kg	Hydroalcoholic extract provides protection against STZ-induced cognitive impairment by reducing oxidative stress, cholinesterase activity, and ROCK II expression.	Jiji & Muralidharan, 2020
Aerial and Root parts	Ethanol extract	300 and 500 mg/kg	Electroshock-induced amnesia in rats.	Significant memory retention was produced, and the root extract was more effective.	Multisona <i>et al.</i> , 2023

**Anti-inflammatory and Antipyretic effects:**

<b>Plant (Part used)</b>	<b>Extract</b>	<b>Dose</b>	<b>Model</b>	<b>Result/ Observations</b>	<b>Reference</b>
Leaves	Water, ethanol, and petroleum ether	100-400 mg/kg	Carrageenan Induced paw oedema	Ethanol extract decreased inflammation.	Bhatia <i>et al.</i> , 2014
Blue flowered (roots)	Methanol extract	200, 300 and 400 mg/kg	Pyretic potential on normal body temperature and Yeast-induced pyrexia in albino rats	Substantial reduction in normal body temperature and yeast-provoked elevated temperature.	Al-Snafi, 2016
Flowers	Ethanol extract	400 mg/kg	Carrageenan-induced paw oedema in rats	Attenuation of inflammation.	Singh <i>et al.</i> , 2018
Roots	Methanol extract	200-400 mg/kg	Carrageenan-Induced paw oedema in rats	Marked anti-oedematous activity.	Swathi <i>et al.</i> , 2021
Roots	Methanol extract	200, 300, and 400 mg/kg	Yeast-induced pyrexia	Marked decline in normal body temperature.	Shirodkar <i>et al.</i> , 2023

**Antibacterial and Antihistaminic activity:**

Plant (Part used)	Extract	Dose	Model	Result/ Observations	Reference
Roots	Ethanol extract	100, 125 and 150 mg/kg	Antihistaminic activity using Clonidine and haloperidol induced catalepsy in mice.	Significant inhibition of clonidine induced catalepsy.	Al-Snafi, 2016
Flowers	Ethyl acetate and Dichloromethane fraction	Various concentrations	Antibacterial activity against <i>Staphylococcus aureus</i> and <i>Escherichia coli</i> .	Potent antibacterial activity exhibited by both with a Minimum Inhibitory Concentration (MIC) value of 0.156 mg/ml.	Indrianingsih <i>et al.</i> , 2021

**Anticonvulsant activity and Analgesic activity:**

Plant (Part used)	Extract	Dose	Model	Result/ Observations	Reference
Leaves	Water, ethanol and petroleum ether	100-400 mg/kg	Tail flick method	Ethanol extract exhibits significant analgesic activity.	Bhatia <i>et al.</i> , 2014
Aerial parts	Methanol extract	100 mg/kg	Pentylene-tetrazol and Maximal electroshock induced seizures.	Postponed the onset of convulsions and reduction of the duration of tonic hind limb extensions.	Gollen <i>et al.</i> , 2018
		230 and 460 mg/kg		Did not show effective results against PTZ and MES induced seizures in rats.	

**Antidepressant, Anti-anxiety activity:**

Plant (Part used)	Extract	Dose	Model	Result/ Observations	Reference
Roots	Methanol extract	100 and 400 mg/kg	Elevated plus and light/dark exploration test.	Time spent in the open arms of EPM increased and there was increase in time spent in the light box at higher doses of CT in light/dark exploration test.	Jiji & Muralidharan, 2020
Roots	Methanol extract	100 and 400 mg/kg	Tail suspension test in mice	Noteworthy decrease in the duration of immobility.	Yanuarto <i>et al.</i> , 2023

**Hepatoprotective activity:**

<b>Plant (Part used)</b>	<b>Extract</b>	<b>Dose</b>	<b>Model</b>	<b>Result/ Observations</b>	<b>Reference</b>
White and Blue Flower (Leaves)	Alcohol extract	200 mg/kg	Paracetamol induced liver toxicity.	Protection against paracetamol-induced liver toxicity and acts by reducing the increased levels of aminotransferase, and bilirubin as well as histopathological damages caused by the liver toxicity.	Gollen <i>et al.</i> , 2018
		200 mg/kg	Carbon tetrachloride induced liver toxicity.	White-Flowered leaves extract showed more better hepato-protective activity as compared to the blue-flowered leaves extract.	

## 10. Safety and Toxicity Studies:

Toxicity studies are evaluated by using methods to determine the safe dose of product to ensure the safety of human being. The toxicity of the natural compounds will depend on the availability of the chemical constituents present in them. Medicinal plants commonly have beneficial effect but some of the naturally occurring compounds show toxic effect (Kurniawati *et al.*, 2023). *C. ternatea* extracts have been examined and are reported to be safe.

- Ethanol extract of roots has LD<sub>50</sub> >1300 mg/kg in mice with no deaths observed up to 3000 mg/kg in mice in acute oral toxicity studies. There was no mortality or disorder observed up to 72 hours after the administration of single dose of 1000 mg/kg in rats.
- Ethanol extract of aerial parts of *C. ternatea* was safe up to 2 g/kg (p.o.) in rats. Animals involved in the study exhibit reduced mobility but no sign of convulsions (Verma *et al.*, 2013).

## 11. Clinical trials:


Clinical trials are carried out to make drug development more efficient and informative for the population. The clinical data will help to find out the suitable range of dose for human being. Some of the clinical trials of *C. ternatea* were reported which are presented in table 10.

Table 10: Clinical trials of *C. ternatea*.

Plant (Part used)	Extract/Formulation	Dose/Route	Participant	Result/Observations	Reference
Flowers	Aqueous	1-2 g in 400 ml water (orally)	Healthy Adult Males	Antioxidant capacity of plasma increased. postprandial sucrose and insulin levels were decreased. Postprandial antioxidant status was enhanced.	Chusak <i>et al.</i> , 2018b
Leaves	Lepa (Ayurvedic formulation)	-	Either gender	Formulation was effective for skin diseases (Dadru).	Narnavar <i>et al.</i> , 2023


## 12. Applications:

Along with medicinal benefits the plant has multifaceted applications:



### 12.1. Agriculture:

In agriculture it is used as:



#### 12.1.1. Cultivation of *C. ternatea* as a fodder crop:

It is palatable hay for cattle and its yield reaches approximately 17-29 tons/hectare. The quantity of dry matter content and digestibility of the plant depends on the timing of harvesting. It increases the energy density of the feed as well as possess high amount of nitrogen because of the presence of low acid detergent fibre content. It also served as source of carotenoids (Oguis *et al.*, 2019).

#### 12.1.2. Biopesticide Activity:

##### 12.1.2.1. Antibacterial and Antifungal:

Finotin protein also possesses antibacterial activity against *Xanthomonas axonopodis*. Ethanol extract of the plant exhibits antibacterial activity against different *Bacillus* species, *Enterococcus faecalis*, *Staphylococcus* species, *Streptococcus* species, and (Shahid *et al.*, 2009).

Ultrasound assisted extraction of the plant shows greater amount of antibacterial property due to high anthocyanin content (Anthika *et al.*, 2015).

Methanol leave extract of the plant shows antifungal activity against mold fungus *Aspergillus niger* ((Kamilla *et al.*, 2009).

##### 12.1.2.2. Anthelmintic:

Numerous investigations revealed varying levels of resistance against the parasitic root-knot nematode (*Meloidogyne incognita*) were shown by 27 homozygous lines of *C. ternatea*. Its methanolic extract exhibit 93% inhibition towards *M. incognita* (Oguis *et al.*, 2019).

##### 12.1.2.3. Insecticide:

Kelemu *et al.*, 2004; Poth *et al.*, 2011a reported that *C. ternatea* act as insecticide due to the presence of proteins and peptides. It contains a protein named finotin which cause 100% larval mortality in concentration 1% w/w and 5% w/w to bruchids *Acanthoscelides obtectus* and *Zabrotes subfasciatus*, respectively.

Larval growth retardation is also reported in dose dependent manner lepidopteran species (*Helicoverpa armigera*), when cyclotide (Cter M) was added in the diet (Poth *et al.*, 2011a).

The extracts of *C. ternatea* served as environmental-friendly natural insecticide (Mensah *et al.*, 2015).

### 12.2. Cosmetics:

The flowers are rich in antioxidant constituents which are valuable to combat oxidative stress and delay skin ageing. According to Dziok *et al.*, 2021 due to the antiaging property of *C. ternatea* it is highly demanded in the market as pharmaceutical and cosmetic product because it helps in improving the youth appearance and keep skin hydrated. Different formulations of *C. ternatea* are presented in table 11.

**Table 11: Different cosmetic formulations of *C. ternatea*.**

Plant part	Formulation	Composition	Benefits	Reference
Whole plant	Face wash gel	<i>C. ternatea</i> alcoholic extract Water Propylene glycol Sodium lauryl sulphate Triethanolamine	Provides clear glossy and healthy skin.	Simanchal Panda.,2018
Seeds	Sunscreen	<i>Cucurbita reticulata</i> extract <i>Cucurbita moschata</i> extract <i>C. ternatea</i> extract <i>Aloe vera</i> Sun products	<i>C. ternatea</i> showed higher sun protective activity than sunscreen product.	Mustaffa <i>et al.</i> , 2018
Flower	Sun block lotion	<i>C. ternatea</i> extract <i>Pandanumusa</i> <i>Paradisiaca</i> extract Virgin coconut oil Glycerin Stearic acid Triethanolamine Methylparaben	Provide protection against the UV rays.	Ritonga <i>et al.</i> , 2020
Flower	Lotion (0.05%, 0.1% and 0.5%)	<i>C. ternatea</i> extract Stearic acid Triethanolamine Liquid Paraffin Cetyl alcohol Glycerin Methyl paraben Propylparaben Aquadest	0.1% concentration of telang flower extract has a very strong antioxidant activity because it has an IC <sub>50</sub> value 37.92 ppm.	Jayanti <i>et al.</i> , 2021
Flower	Eye shadow	<i>C. ternatea</i> extract <i>Caesalpinia sappan</i> extract Cetyl Alcohol Stearic Acid Petrolatum Polawax Polysorbate 80 Sorbitan Monooleate 80 Mineral Oil Butylate Hydroxy Toluene Triethanolamine Propylene Glycol Methyl Paraben Propyl Paraben Water Talcum	This combination of eye shadow showed mild irritation but when combined with some other natural dye provide more stable formulation.	Yuniaty <i>et al.</i> , 2023

### 12.3. Food Product:

Some of the studies reported that if the anthocyanins present in *C. ternatea* flowers were used as food colorant they possess potent antioxidant activity. Leong *et al.*, 2017 reported that anthocyanins can play two roles: one is coloring agent, and the other one is bio preservative.

Application of *C. ternatea* in different food products are presented in figure 7.




Figure 7: *C. ternatea* in food products application

Figure 8 highlights the milestones in the systematic study of this plant.




Figure 8: Milestones of *C. ternatea* Studies (1954 to 2017)

### 13. Challenges and Future direction:

To get the regular supply of the plant better Good Agricultural Practices (GAP) to be documented. A limitations of *C. ternatea* plant is that the colour pigments of the plant are unstable in nature and readily degrade when expose to different environmental conditions like light, pH, and temperature (Kaushik *et al.*, 2022). To prepare stable formulations of the plant pigments, new drug delivery systems may be investigated. Microencapsulation or nanoparticle formulations may be a suitable way to safeguard the natural plant pigments. In order to develop the plant/ plant product further evidence should be strengthened with detailed pharmacological as well as human trials. Comprehensive investigations will ultimately lead to the development of a medicine/ nutraceutical or cosmeceutical from this delicate but powerful plant.

### 14. Conclusion:

The present review highlights the versatile role of this beautiful plant. *C. ternatea* is not only an ornamental plant but also offers diverse benefits in the medicine fields, cosmetics, food industry and agrotechnology. *C. ternatea* is an abundant source of different phytoconstituents which offer health benefits to humans. It acts as an additive when supplemented with functional food, pharmaceutical drug, and results in an increase in treatment efficiency. The plant demonstrates a variety of actions, and it exhibits a low toxicity profile. From the above-mentioned data, we conclude that *C. ternatea* is versatile, safe, and effective.

## References:

- Alok, S., et al. (2015). "An update on Ayurvedic herb vishnukanta (*Clitoria ternatea* Linn.): A review." *International Journal of Life Sciences and Review* 1(1), 1-9.
- Al-Snafi, A. E. (2016). Pharmacological importance of *Clitoria ternatea*—A review. *IOSR Journal of Pharmacy*, 6(3), 68-83.
- Ashraf, K., Adlin, N. F., Basri, A. N., Ahmad, W., & Sultan, S. (2024). "The Traditional Uses, Phytochemistry, and Pharmacological Effects of *Clitoria ternatea*: A Review." *Indian Journal of Pharmaceutical Education and Research*, 58(1), 1-14.
- Baskaran, A., Mudalib, S. K. A., & Izirwan, I. (2019). "Optimization of aqueous extraction of blue dye from butterfly pea flower". *Journal of Physics: Conference Series* (Vol. 1358, No. 1, p. 012001). IOP Publishing.
- Bishoyi, A. K., & Geetha, K. A. (2012). "Polymorphism in flower colour and petal type in Aparajita (*Clitoria ternatea*)". *Open Access Journal of Medicinal and Aromatic Plants*, 3(2), 12.
- Bujak, T., Zagórska-Dziok, M., Ziemlewska, A., Nizioł-Łukaszewska, Z., Wasilewski, T., & Hordyjewicz-Baran, Z. (2021). "Antioxidant and cytoprotective properties of plant extract from dry flowers as functional dyes for cosmetic products." *Molecules*, 26(9), 2809.
- Cavalcanti, R. N., Santos, D. T., & Meireles, M. A. A. (2011). "Non-thermal stabilization mechanisms of anthocyanins in model and food systems—An overview." *Food Research International*, 44(2), 499-509.
- Chong, F. C., & Gwee, X. F. (2015). "Ultrasonic extraction of anthocyanin from *Clitoria ternatea* flowers using response surface methodology". *Natural Product Research*, 29(15), 1485-1487.
- Chusak, C., Thilavech, T., Henry, C. J., & Adisakwattana, S. (2018). "Acute effect of *Clitoria ternatea* flower beverage on glycemic response and antioxidant capacity in healthy subjects: a randomized crossover trial." *BMC Complementary and Alternative Medicine*, 18(1), 1-11.
- Damodaran, T., Cheah, P. S., Murugaiyah, V., & Hassan, Z. (2020). "The nootropic and anticholinesterase activities of *Clitoria ternatea* Linn. root extract: Potential treatment for cognitive decline." *Neurochemistry International*, 139, 104785.
- Deshmukh, S., & Jadhav, V. (2014). "Bromatological and mineral assessment of *Clitoria ternatea* Linn. leaves." *Energy (KJ)*, 459(9.20), 489.424-482.483.
- Divya, A., Anbumalarmathi, J., & Sharmili, S. (2018). "Phytochemical analysis, antimicrobial and antioxidant activity of *Clitoria ternatea* blue and white flowered leaves." *Advances in Research*, 14(5), 1-13.
- Fu, X., Wu, Q., Wang, J., Chen, Y., Zhu, G., & Zhu, Z. (2021). "Spectral characteristic, storage stability and antioxidant properties of anthocyanin extracts from flowers of butterfly pea (*Clitoria ternatea* L.)". *Molecules*, 26(22), 7000.
- Goh, S. E., Kwong, P. J., Ng, C. L., Ng, W. J., & Ee, K. Y. (2021). "Antioxidant-rich *Clitoria ternatea* L. flower and its benefits in improving murine reproductive performance". *Food Science and Technology*, 42, e25921.
- Gollen, B., Mehla, J., & Gupta, P. (2018). "*Clitoria ternatea* Linn: a herb with potential

pharmacological activities: future prospects as therapeutic herbal medicine". *Journal of Pharmacological Reports*, 3(1), 1-8.

Gomez, S. M., & Kalamani, A. (2003). "Butterfly pea (*Clitoria ternatea*): A nutritive multipurpose forage legume for the tropics—an overview". *Pakistan Journal of Nutrition*, 2(6), 374-379.

Gupta, G. K., Chahal, J., & Bhatia, M. (2010). "*Clitoria ternatea* (L.): Old and new aspects." *Journal of Pharmacy Research*, 3(11), 2610-2614.

Handa, S. S. (2008). "An overview of extraction techniques for medicinal and aromatic plants. Extraction technologies for medicinal and aromatic plants", 1(1), 21-40.

Indrianingsih, A. W., Wulanjati, M. P., Windarsih, A., Bhattacharjya, D. K., Suzuki, T., & Katayama, T. (2021). "In vitro studies of antioxidant, antidiabetic, and antibacterial activities of *Theobroma cacao*, *Annona muricata* and *Clitoria ternatea*." *Biocatalysis and Agricultural Biotechnology*, 33, 101995.

Jain, R. A., & Shukla, S. H. (2011). "Pharmacognostic Evaluation and Phytochemical Studies on Stem of *Clitoria ternatea* linn." *Pharmacognosy Journal*, 3(24), 62-66.

Jamil, N., Zairi, M. N. M., Nasim, N. A. I. M., & Pa'ee, F. (2018). "Influences of environmental conditions to phytoconstituents in *Clitoria ternatea* (butterfly pea flower)—A review". *Journal of Science and Technology*, 10(2). Jayanti, M., Ulfa, A. M., & Yasir, A. S. (2021). "The Formulation and Physical Evaluation Tests of Ethanol in Telang Flower (*Clitoria ternatea* L.) Extract Losio Form as Antioxidant." *Biomedical Journal of Indonesia*, 7(3), 488-495.

Jeyaraj, E. J., Lim, Y. Y., & Choo, W. S. (2021). "Extraction methods of butterfly pea (*Clitoria ternatea*) flower and biological activities of its phytochemicals." *Journal of Food Science and Technology*, 58(6), 2054-2067.

Jiji, K., & Muralidharan, P. (2020). "Neuropharmacological Potential of *Clitoria ternatea* Linn.- A Review." *Research Journal of Pharmacy and Technology*, 13(11), 5497-5502.

Kamilla, L., Mansor, S. M., Ramanathan, S., & Sasidharan, S. (2009). "Effects of *Clitoria ternatea* leaf extract on growth and morphogenesis of *Aspergillus niger*." *Microscopy and Microanalysis*, 15(4), 366-372.

Kavitha, R. (2018). "Biochemical studies on the effect of ethanolic extracts of *Trichosanthes dioica* and *Clitoria ternatea* in streptozotocin induced diabetic male wistar rats". *International Journal of Pharmaceutical Sciences and Research*, 9, 4682-4689.

Kazuma, K., Noda, N., & Suzuki, M. (2003). "Flavonoid composition related to petal color in different lines of *Clitoria ternatea*." *Phytochemistry*, 64(6), 1133-1139.

Kelemu, S., Cardona, C., & Segura, G. (2004). "Antimicrobial and insecticidal protein isolated from seeds of *Clitoria ternatea*, a tropical forage legume". *Plant Physiology and Biochemistry*, 42(11), 867-873.

Kurniawati, E. Y., Pramono, N., Hidayat, S. T., & Mahati, E. (2023). "In Silico Pharmacokinetic and Toxicity Analysis on *Clitoria Ternatea* Flower". *Jurnal Farmasi Indonesia*, 20(2), 124-135.

Lakshan, S. A. T., Jayanath, N. Y., Abeysekera, W. P. K. M., & Abeysekera, W. K. S. M. (2019). "A commercial potential blue pea (*Clitoria ternatea* L.) flower extract incorporated

beverage having functional properties". *Evidence-Based Complementary and Alternative Medicine*, 2019.

Leong, C. R., Azizi, M. A. K., Taher, M. A., Wahidin, S., Lee, K. C., Tan, W. N., & Tong, W. Y. (2017). "Anthocyanins from *Clitoria ternatea* attenuate food borne *Penicillium expansum* and its potential application as food biopreservative." *Natural Product Sciences*, 23(2), 125-131.

Ludin, N. A., Al-Alwani, M. A., Mohamad, A. B., Kadhum, A. A. H., Hamid, N. H., Ibrahim, M. A., Teridi, M. A. M., Al-Hakeem, T. M. A., Mukhlus, A., & Sopian, K. (2018). "Utilization of natural dyes from *Zingiber officinale* leaves and *Clitoria ternatea* flowers to prepare new photosensitisers for dye-sensitised solar cells." *International Journal of Electrochemical Science*. 13(8), 7451-7465.

Malabadi, R. B., & Nataraja, K. (2001). "Shoot regeneration in leaf explants of *Clitoria ternatea* L. cultured *in vitro*". CABI Digital Library.

Manisha, B., Jagbir, C., & Sumeet, G. (2014). "Analgesic and anti-inflammatory activities of *Clitoria ternatea* Linn. leaves extract on rat model." *International Journal of Pharmaceutical Sciences and Research (IJPSR)*, 5(2), 600- 606.

Marsin, A. M., Jusoh, Y. M. M., Abang, D. N., Zaidel, Z. H., Yusof, A. H. M., & Muhamad, I. I. (2020). "Microwave-assisted encapsulation of blue pea flower (*Clitoria ternatea*) colourant: Maltodextrin concentration, power, and time". *Chemical Engineering Transactions*, 78, 199-204.

Mauludifia, F., Astrinia, S., Meiranti, K., & Djaeni, M. (2019). "Production of natural colorant powder from *Clitoria ternatea* L. using tray dryer which is dehumidified by zeolite". *Journal of Physics: Conference Series*.

Mensah, R., Leach, D., Young, A., Watts, N., & Glennie, P. (2015). "Development of *Clitoria ternatea* as a biopesticide for cotton pest management: assessment of product effect on *Helicoverpa* spp. and their natural enemies". *Entomologia Experimentalis et Applicata*, 154(2), 131-145.

Mishra, A. K., Singh, J., & Tiwari, K. N. (2019). "*In vitro* Regeneration of *Clitoria ternatea* (L.) from Nodal Explant." *International Journal on Emerging Technologies*, 10(1), 35-41.

Morris, J. B. (2009). "Characterization of butterfly pea (*Clitoria ternatea* L.) accessions for morphology, phenology, reproduction and potential nutraceutical, pharmaceutical trait utilization." *Genetic Resources and Crop Evolution*, 56, 421-427.

Mukherjee, P. K., Kumar, V., Kumar, N. S., & Heinrich, M. (2008). "The Ayurvedic medicine *Clitoria ternatea*— From traditional use to scientific assessment". *Journal of Ethnopharmacology*, 120(3), 291-301.

Multisona, R. R., Shirodkar, S., Arnold, M., & Gramza-Michalowska, A. (2023). "*Clitoria ternatea* flower and its bioactive compounds: Potential use as microencapsulated ingredient for functional foods". *Applied Sciences*, 13(4), 2134.

Narnavar, N., Nangare, N., Deshpande, M., & Mahajan, M. (2023). "Clinical Study of Aparajita (*Clitoria ternatea* Linn.) in Dadru (Tinea)." *Korean Journal of Physiology and Pharmacology*, 27(3), 154-159.

Oguis, G. K., et al. (2019). "Butterfly pea (*Clitoria ternatea*), a cyclotide-bearing plant with applications in agriculture and medicine." *Frontiers in Plant Science* 10: 645.

- Panda, S. (2018). "Formulation and evaluation *Clitoria ternatea* linn. alcoholic extract facial wash gel". *Journal of Emerging Technologies and Innovative Research*, 10(5).
- Patel, A., & Mishra, S. "Clitoria ternatea L.: a medicinal climber of India." *Bhagwati Prashad Sharma Banshidhar Behera Sweta Mishra*, 47.
- Poth, A. G., Colgrave, M. L., Lyons, R. E., Daly, N. L., & Craik, D. J. (2011). "Discovery of an unusual biosynthetic origin for circular proteins in legumes". *Proceedings of the National Academy of Sciences*, 108(25), 10127-10132.
- Rampalli, S. (2022). "Pharmacognostical investigation of *Clitoria ternatea* L. leaves." *Plant Science Today*, 9(4), 814-819.
- Ritonga, N. B., Rini, R., & Anggraini, T. (2020). "Formulation and Evaluation of Sun Block Lotion Made from Virgin Coconut Oil (VCO) with the addition of the Extract of Telang Flower (*Clitoria ternatea*, L) and Pandan Leaves (*Pandanumusa paradisiaca*, L)". *Asian Journal of Applied Research for Community Development and Empowerment*, 4(1), 59-63.
- Rout, G. R. (2005). "Micropropagation of *Clitoria ternatea* Linn. (Fabaceae)—An important medicinal plant". *In Vitro Cellular & Developmental Biology-Plant*, 41, 516-519.
- Sarma, D. S. K., Kumar, D., Yamini, C., Santhalahari, C., Lahari, C., Kumar, G. C., & Lahitha, M. (2023). "Review on *Clitoria ternatea*. *International Journal of Pharmaceutical Sciences and Medicine*, 8(9), 43-58.
- Shahid, M., Shahzad, A., & Anis, M. (2009). "Antibacterial potential of the extracts derived from leaves and *in vitro* raised calli of medicinal plants *Pterocarpus marsupium* Roxb., *Clitoria ternatea* L., and *Sanseveiria cylindrica* Bojer ex Hook". *Advances in Traditional Medicine*, 9(2), 174-181.
- Shekhawat, N., & Vijayvergia, R. (2010). "Comparative study of primary metabolites in different plant parts of *Clitoria ternatea* (L.), *Guazuma ulmifolia* (Lam.) & *Madhuca indica* (Gmel.)". *Journal of Chemical and Pharmaceutical Research*, 2(2), 168-171.
- Shirodkar, S. M., Multisona, R. R., & Gramza-Michalowska, A. (2023). "The Potential for the Implementation of Pea Flower (*Clitoria ternatea*) Health Properties in Food Matrix." *Applied Sciences*, 13(12), 7141.
- Singh, N. K., Garabadu, D., Sharma, P., Shrivastava, S. K., & Mishra, P. (2018). "Anti-allergy and anti-tussive activity of *Clitoria ternatea* L. in experimental animals". *Journal of Ethnopharmacology*, 224, 15-26.
- Suarna, I. W., & Wijaya, I. M. S. (2021). "Butterfly pea (*Clitoria ternatea* L.: Fabaceae) and its morphological variations in Bali". *Journal of Tropical Biodiversity and Biotechnology*, 6(2), 63013.
- Swathi, K., Jayaram, S., Sugumar, D., & Rymbai, E. (2021). "Evaluation of anti-inflammatory and anti-arthritic property of ethanolic extract of *Clitoria ternatea*". *Chinese Herbal Medicines*, 13(2), 243-249.
- Syafa'atullah, A. Q., Amira, A., Hidayati, S., & Mahfud, M. (2020, June). "Anthocyanin from butterfly pea flowers (*Clitoria ternatea*) by ultrasonic-assisted extraction". In *AIP Conference Proceedings* (Vol. 2237, No. 1). AIP Publishing.
- Talpatte, K. A., Bhosale, U. A., Zambare, M. R., & Somani, R. S. (2014). "Neuroprotective and nootropic activity of *Clitoria ternatea* Linn.(Fabaceae) leaves on diabetes induced cognitive

decline in experimental animals". *Journal of Pharmacy & Bioallied sciences*, 6(1), 48.

Tamanna.,2023 M. Pharm thesis, Evaluation of *C. ternatea* Linn. and *Tagetes erecta* Linn. extracts: Potential Pharmacological Therapy for Dementia.

Taur, D. J., & Patil, R. Y. (2010). "Pharmacognostic evaluation of *Clitoria ternatea* root". *Journal of Pharmacy Research*, 3(2), 205-207.

Terahara, N., Toki, K., Saito, N., Honda, T., Matsui, T., & Osajima, Y. (1998). "Eight New Anthocyanins, Ternatins C1- C5 and D3 and Preternatins A3 and C4 from Young *Clitoria ternatea* Flowers". *Journal of Natural Products*, 61(11), 1361-1367.

Tuan Putra, T., Zainol, M., Mohdlsa, N., & Mohdmaidin, N. (2021). "Chemical characterization of ethanolic extract of Butterfly pea flower (*Clitoria ternatea*).". *Food Research*, 5(4), 127-134.

Turnos, L. J. N. (2021). "Blue Ternate (*Clitoria ternatea* L.): Nutritive analysis of flowers and seeds". *Asian Journal of Fundamental and Applied Sciences*, 2(2), 103-112.

Tyan, C. Y., Radhakrishnan, L., Mustafa, F., & Sahgal, G. (2018). "Antioxidant, antimicrobial and SPF protective activity of *Cucurbita moschata*, *Cucurbita reticulata* and *Clitoria ternatea*." *Reports De Pharmacie*, 4, 488-491.

Verma, P. R., Itankar, P. R., & Arora, S. K. (2013). "Evaluation of antidiabetic antihyperlipidemic and pancreatic regeneration, potential of aerial parts of *Clitoria ternatea*." *Revista Brasileira de Farmacognosia*, 23(5), 819-829.

Weerasinghe, T., Perera, D., De Silva, N., Poogoda, D., & Swarnathilaka, H. (2022). "Butterfly pea: An emerging plant with applications in food and medicine". *The Pharma Innovation Journal*, 11, 625-37.

Widowati, W., Darsono, L., Lucianus, J., Setiabudi, E., Obeng, S. S., Stefani, S., Wahyudianingsih, R., Tandibua, K. R., Gunawan, R., & Wijayanti, C. R. (2023). "Butterfly pea flower (*Clitoria ternatea* L.) extract displayed antidiabetic effect through antioxidant, anti-inflammatory, lower hepatic GSK-3 $\beta$ , and pancreatic glycogen on Diabetes Mellitus and dyslipidemia rat". *Journal of King Saud University-Science*, 35(4), 102579.

Widowati, W., Wargasetia, T. L., Marthania, M., Hanifa, T. S., Zakaria, T. M., Gunadi, M. S., Halim, N., & Santiadi, S. (2022). "Antioxidant Properties of Tenan Herbal Tea Formulation "Telang (*Clitoria ternatea*) and Pineapple (*Ananas comosus*)". *Jurnal Kedokteran Brawijaya*, 87-93.

Yanuarto, T., Haque, A. F., Mama, M. R., Edriani, A., Asfarina, A., & Setiawati, L. (2023). "Pembuatan masker organik dari Sari Telang (*Clitoria ternatea* L.) dan bahan-bahan alami lainnya di sma negeri 11 kota bengkulu." *Jurnal sapta mengabdikan*, 3(1), 1-6.

Yuniaty, D. A., Rahmat, D., & Rachmat, R. (2023). "Formulation of eyeshadow cream combination of extract spissum butterfly pea flower (*Clitoria ternatea* L.) with Secang wood (*Caesalpinia sappan* L.) as a natural color". *Medical Sains: Jurnal Ilmiah Kefarmasian*, 8(3), 1185-1196.