

## Effect of botanicals with carbendazim against anthracnose (*Colletotrichum lindemuthianum*) of black gram (*Vigna mungo* L.)

### ABSTRACT

The present study was carried out to evaluate the effect of selected botanicals and carbendazim for the management of anthracnose disease on black gram caused by *Colletotrichum lindemuthianum* under field conditions. Three replications of black gram were planted in a randomized block design at the research plot of the Central Research Field, Department of Plant Pathology, SHUATS, Prayagraj during *kharif* season of 2023. Carbendazim 50% WP @2g/kg was applied for seed treatment of black gram before sowing. Foliar spray was conducted using five botanicals and one fungicide viz. T<sub>1</sub>-neem oil (2ml/L), T<sub>2</sub>- eucalyptus oil (2ml/L), T<sub>3</sub>- tea tree oil (2ml/L), T<sub>4</sub>- Lavender oil (2ml/L), T<sub>5</sub>- Thyme oil (5%), T<sub>6</sub>- mancozeb 75 WP (0.2%) and T<sub>0</sub>- control with each treatment replicating three times. It was observed that all treatments significantly reduced the intensity of disease and increased growth parameters of garlic. From the study of the results, it was concluded that among all the treatments, the most effective in managing the disease was T<sub>6</sub>- Carbendazim @0.1% ST+FS (22.33%) followed by T<sub>1</sub>-Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (23.14%) as compared to T<sub>0</sub>- Control (43.48%) and in influencing the yield (t/ha.) T<sub>6</sub>- Carbendazim @0.1% ST+FS (7.86) followed by T<sub>1</sub>-Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (6.76) as compared to T<sub>0</sub>- Control (3.15) in field condition and maximum cost benefit ratio was recorded in T<sub>6</sub>-Carbendazim @0.1% ST+FS (1:2.43) followed by T<sub>1</sub>-Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (1:2.07) as compared to T<sub>0</sub>- Control (1:1.03).

**Key words:** Anthracnose, Botanicals, Carbendazim, *Colletotrichum lindemuthianum*, Per cent Disease Intensity, Yield.

### INTRODUCTION

Black gram (*Vigna mungo* L.) commonly known as urd bean, mash or black maple is an annual, semi erect to spreading herb belonging to the family Fabaceae and it is grown as a *Kharif* crop in tropical and sub-tropical countries including India, Nigeria, Thailand, Philippines, Upper Volta, Zambia, Palmira and Columbia. This versatile pulse crop has a relatively short growth duration. As a leguminous crop, black gram not only enriches soil fertility but also contributes to the nitrogen economy of subsequent crops in the rotation [11].

Comment [B1]: (Ref)

The urd bean production of India was 2.78 million tonnes from acreage of 4.63 million hectares with a productivity of 600 Kg/ha. India is the largest producer and consumer of urd bean. During 2021-22, the total quantity of import of urd bean was 6.12 lakh tonnes, whereas, export of urd bean was 55.18 thousand tonnes. In the case of area, the share of Madhya Pradesh is highest (24 %), followed by Uttar Pradesh (10 per cent) and Maharashtra (7 %) during 2020-21. Among the major producing states, productivity was highest in Andhra Pradesh (915 kg/ha), followed by Jharkhand (879 kg/ha) during 2020-21 [4].

*Colletotrichum lindemuthianum* Lams. Scrib., the causal organism of pulse anthracnose, is a serious soil borne as well as seed borne pathogen of pulse crops throughout the world and has been reported to occur on *Kharif* pulses in several countries [12]. Anthracnose disease of mung bean was observed red or brown lesions on leaves, stems, branches and pods of different cultivars of *Vigna radiata* and *Vigna mungo* in Gurdaspur [1]. In India, the black gram anthracnose was first reported from Uttar Pradesh state in 1984 [16]. The pathogen was causing **saviour** yield losses nearly 80 to 100 % under cool and humid environmental conditions. Black gram yields are highly reduced due to anthracnose disease up to 40-67 % [7].

In the present investigation, it is evident that seed treatment + foliar spray of carbendazim gave maximum cost benefit ratio [10]. Use of botanicals is the safe alternative control. To prevent anthracnose disease which is caused by different species of *Colletotrichum*, botanicals viz. eucalyptus oil, lavender oil, tea tree oil and thyme oil are used as natural fungicides and resulted in effective way [16].

## MATERIALS AND METHODS

### Isolation of fungal pathogen

Prepared PDA was melted and 80 mg of streptomycin, an antibiotic was added during preparation of PDA to each 500 ml preparation of the PDA to inhibit probable bacterial growth. The leaves of black gram showing typical symptoms were collected and the standard tissue isolation procedure was followed to isolate the pathogen. The infected 14 leaf parts were cut into small pieces of **two to three** mm dimension in a manner so that pieces may have some green portion also. Such leaf bits were surface sterilized with 1% sodium hypochlorite (NaOCl) solution for 30 seconds and washed three times with sterile distilled water to remove any traces of mercuric chloride adhered with leaf bits [18]. **2-3** leaf bits were transferred on PDA medium contained in Petri plates aseptically with the help of sterilized forceps. These

Petri plates were incubated at room temperature ( $25 \pm 2^\circ$  C). After 3-4 days mycelia growth

Comment [B2]: Use another word for easily comprehension

Comment [B3]: delete

Comment [B4]: In full

Comment [B5]: 2 to 3 mm

Comment [B6]: Two to three

Comment [B7]: Back space

was observed around leaf bits from this colony growth, a portion from the periphery that is, single hyphal tip was separated and transferred to **other**.

**Comment [B8]:** another

### Field Preparation

An experiment was conducted during *Kharif* season of 2023 at the central research farm of Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj. The selected field area was prepared by ploughed and harrowing during summer season, Experimental plots were laid out as per statistical Randomized Block design. Total area was divided into 21 plots and plot size 2 x 1 m<sup>2</sup>. To evaluate the effect of botanicals with carbendazim against *Colletotrichumlindemuthianum* with Randomized Block Design in three **replications**. The crop was raised as per recommended package of practices and protective irrigation was given as and when required.

**Comment [B9]:** , the crop

### Seed Treatment with carbendazim

The seeds of black gram were treated with carbendazim 50% WP at a rate of 2 grams per kilogram of seed. **Thoroughly mixed the carbendazim with the seeds**. This was done by shaking the seeds in a bag or container to ensure the fungicide coats the seed uniformly. The seeds were soaked in this treatment for 24 hours before sowing to ensure effective coating and absorption of the fungicide. This treatment is designed to protect the seeds from fungal pathogens during germination and early growth stages, promoting healthier seedling development and ensuring better crop establishment.

**Comment [B10]:** The carbendazim was thoroughly mixed with the seeds

### Preparation of botanicals and spraying

The botanicals *viz.*, neem oil, eucalyptus oil, tea tree oil, lavender oil and thyme oils were mixed in appropriate water quantity to make solution at concentration as per treatments. All the solutions were prepared freshly and being sprayed at 30, 45, and 60 **DAS** of interval.

**Comment [B11]:** In full Days after sowing

Treatments were imposed after appearance of the first disease symptoms. Observations on disease intensity (%) of anthracnose of black gram were recorded at 15 days interval, yield (t/ha) and C:B ratio data were obtained after the harvest on physiological maturity.

Assessment of per cent disease intensity (%) of anthracnose of black gram was recorded at 30, 45 and 60 days after sowing. The disease intensity was calculated using the formula **[20]** which is given below

$$\text{Disease Intensity (\%)} = \frac{\text{Sum of all disease ratings}}{\text{Total number of rating} \times \text{Maximum disease grade}} \times 100$$

**Table No.1 Scale/grade used for Treatments**

Scale/Grade	Description
0	No symptoms on leaves
1	Small size lesions covering 1% or less of leaf area.
3	Small size lesions covering 1-10% of leaf area.
5	Lesions size big but not coalescing, covering 11-25% of the leaf area.
7	Lesions on leaves covering 26-50% of leaf area. Cankers on stem and pod infection.
9	Lesions on leaves covering 51% or more of leaf area. Defoliation of leaves, deep cankers on stem and pods, blighting of plant occurs.

**Comment [B12]:** Table 1:  
The table should be on one page

#### Economics:

Gross returns were calculated by multiplying total yield with market price of the produce. Cost of cultivation and cost of treatments were deducted from the gross returns, to find out returns and cost benefit of ratio by following formula [14].

$$\text{Cost Benefit Ratio} = \frac{\text{Gross return}}{\text{Total cost of cultivation}}$$

#### Statistical analysis:

The data recorded during the trial was subjected to be statistically analyzed by randomized block design (RBD) for drawing the conclusion. The significant and non-significant values were evaluated with the help of "F" (variance ratio) table. The calculated values were compared with the tabulated values at 5% level of probability for the appropriate degree of freedom[3].

## RESULTS

A field study was carried out to assess on various aspects of anthracnose of black gram (*Vigna mungo*L.) caused by *Colletotrichum lindemuthianum* with reference to evaluation of disease intensity (%), yield (q/ha) and cost benefit ratio among the treatments.

The results of the field experiment presented in **Table No.2** two clearly indicate that the disease

intensity was significantly low in all the treated plots compared to the unsprayed control plot after two sprays. Disease intensity was recorded three times before spray, fifteen days after first spray and fifteen days after second spray of botanicals along with carbendazim, respectively.

Comment [B13]: Table 2

Comment [B14]: Table 2 clearly

### Per Cent Disease Intensity of anthracnose of black gram

Among the treatments the significant reduction in the disease intensity (%) at 30, 45 and 60 DAS was recorded in the treatments. The minimum disease intensity (%) at 30, 45 and 60 DAS of black gram was recorded in T<sub>6</sub>- Carbendazim @0.1% ST+FS (14.37, 18.93, 22.33, respectively), followed by T<sub>1</sub>- Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (15.23, 21.25, 23.14, respectively), T<sub>2</sub>- Carbendazim @2g/Kg(ST) + Eucalyptus oil @0.2% (FS) (16.55, 23.31, 24.70, respectively), T<sub>4</sub>- Carbendazim @2g/Kg(ST) + Lavender oil @0.2% (FS) (18.73, 25.29, 29.22, respectively), T<sub>3</sub>- Carbendazim @2g/Kg(ST) + Tea Tree oil @0.2% (FS) (22.29, 25.93, 33.92, respectively) and T<sub>5</sub>- Carbendazim @2g/Kg(ST) + Thyme oil @0.2% (FS) (24.84, 30.07, 35.99, respectively) as compared to non-treated control, T<sub>0</sub>- Control (28.14, 36.57, 43.48).

### Yield(t/ha)

According to the current findings, study of data on yield of black gram reveals that the yield (q/ha.) significantly increased in the treatment T<sub>6</sub>- Carbendazim @0.1% ST+FS (7.86), followed by T<sub>1</sub>- Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (6.76), T<sub>2</sub>- Carbendazim @2g/Kg(ST) + Eucalyptus oil @0.2% (FS) (6.38), T<sub>4</sub>- Carbendazim @2g/Kg(ST) + Lavender oil @0.2% (FS) (6.25), T<sub>3</sub>- Carbendazim @2g/Kg(ST) + Tea Tree oil @0.2% (FS) (5.48) and T<sub>5</sub>- Carbendazim @2g/Kg(ST) + Thyme oil @0.2% (FS) (4.98) as compared to non-treated check, T<sub>0</sub>- Control (3.15).

### Cost Benefit Ratio of the treatment

Results of cost benefit ratio among the treatments were observed. The cost of cultivation (Rs. 8,258/ha) and the highest gross returns (Rs. 55,826/ha), net returns (Rs. 28,856/ha) and C:B ratio (1: 2.07) were recorded in T<sub>1</sub>- Carbendazim @2g/Kg(ST) + Neem oil @0.2% (FS) (6.76). The lowest gross returns (Rs. 26,013/ha) and net returns (Rs. 783/ha) and C:B ratio (1: 1.29) were recorded in T<sub>0</sub>- control (untreated check).

**Table No.2**disease intensity

Treatments	Percentdisease intensity				Yield (t/ha)	C:B ratio
	30DAS (Before spray)	45 DAS (After 1st spray)	60 DAS (After 2nd spray)	Mean		
Control (Untreated check)	28.14 <sup>a</sup>	36.5 <sup>a</sup>	43.8 <sup>a</sup>	36.15	3.15	1: 1.03
Carbendazim@2g/kg seed (S.T) + Neem oil @0.2% (F.S)	15.23 <sup>f</sup>	21.25 <sup>e</sup>	23.14 <sup>f</sup>	19.87	6.76	1: 2.07
Carbendazim@2g/kg seed (S.T) + Eucalyptus oil @0.2% (F.S)	16.55 <sup>e</sup>	23.31 <sup>d</sup>	24.70 <sup>e</sup>	21.52	6.38	1: 1.80
Carbendazim@2g/kg seed (S.T) + Tea Tree oil @0.2% (F.S)	22.29 <sup>c</sup>	25.93 <sup>c</sup>	33.92 <sup>c</sup>	27.38	5.48	1: 1.47
Carbendazim@2g/kg seed (S.T) + Lavender oil @0.2% (F.S)	18.73 <sup>d</sup>	25.29 <sup>c</sup>	29.22 <sup>d</sup>	24.41	6.25	1: 1.73
Carbendazim@2g/kg seed (S.T) + Thyme oil@0.2% (F.S)	24.84 <sup>b</sup>	30.07 <sup>b</sup>	35.99 <sup>b</sup>	30.3	4.98	1: 1.24
Carbendazim 50WP @2g/kg seed (S.T)+ @0.1% (F.S)	14.37 <sup>g</sup>	18.93 <sup>f</sup>	22.33 <sup>g</sup>	18.54	7.86	1: 2.43
<b>C.D (5%)</b>	0.26	0.80	0.49		0.02	
<b>C.V</b>	0.74	1.75	0.92		0.63	

S.T. – Seed Treatment

F.S. – Foliar spray

## DISCUSSION

The probable reasons for such findings may be due to the antifungal properties of neem oil and carbendazim, a systemic fungicide. Carbendazim disrupts fungal cell division by interfering with DNA biosynthesis, specifically inhibiting the formation of mitotic microtubules, which halts the development of fungal germ tubes and mycelial growth by **Chaudhari and Gohel [2]**. However, its persistence and overuse may lead to the development of resistance by **Purushotham et al. [13]**. Neem oil, on the other hand, offers a more sustainable option for managing plant diseases. Its most active component, azadirachtin, along with other compounds like nimbidol, nimbin, sodium nimbin, and quercetin, exhibits both antifeedant and toxic effects on pathogens. Azadirachtin, a terpene limonoid, inhibits microbial growth by disrupting the cell walls of pathogens by **Vijaykumar et al. [19]**. These findings align with the research of **Kulkarni and Raja [8][9]**, **Khan et al. [6]**, **Manjunath et al. [10]**, **Jagtap et al. [5]** and **Singh et al. [17]**. This combination of systemic fungicides and botanical oils represents an effective strategy for reducing disease intensity and increasing yield in crops.

**Comment [B15]:** Table 2: Disease intensity  
The table should be on the same page

**Comment [B16]:** delete

**Comment [B17]:**

## CONCLUSIONS

From the present studies it can be concluded that many have been relied on chemicals and this resulted in many undesirable problems. It is necessary to incorporate alternative control components and the use of indigenous sources for the plant disease management that are effective in field and are organically developed for growing concern for both public health and environment hazards with less cost. Thus, use of botanicals have been taken in consideration for the management of anthracnose of black gram caused by *Colletotrichum lindemuthianum*. The treatment, T<sub>1</sub>- Carbendazim @2g/Kg (ST) + Neem oil @0.2% (FS) have shown good result in managing anthracnose disease of black gram compared to other treatments.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

## REFERENCES

1. Bains, S. S., Kaur, I., Dhaliwal, H.S. and Gill, A.S. (1989). Outbreak of new anthracnose of mung and mash in Punjab. *Indian Bot Repr.* 8(2):164-165.
2. Chaudhari, K. A. and Gohel, N. M. (2016). Management of anthracnose disease of mungbean through new fungicidal formulations. *Journal of Pure and Applied Microbiology.* 10 (1): 691-696.
3. Fisher, R.A. and Yates, F. (1963). The design of experiments (5<sup>th</sup> edition). Oliver and Boyd, Edinburgh, and the design and analysis of factorial experiments. *Imperial Bureau of Soil Science, Harpenden.*
4. ICAR-Indian Institute of Pulses Research portal : <https://iipr.icar.gov.in/urdbean/>
5. Jagtap, G. P., Gavate, D. S. and Dey, U. (2013). Management of *Colletotrichum truncatum* causing anthracnose/pod blight of soybean by fungicide. *Indian Phytopath.* 66(2):177-181.
6. Khan, A. A., Khan, R.U., Singh, R., (2005). Management of *Cercospora* leaf spot and anthracnose diseases of mungbean by fungicides. *Annals of Plant Protection Sciences.* 13(2), 465-529.
7. Kalpana, M., Karthiba, L., Senguttuvan, K. and Parimalarangan, R. (2023). Diagnosis of major foliar diseases in black gram (*Vigna mungo* L.) using Convolution Neural Network. *Legume Research-An International Journal.* 46(7): 940-945.
8. Kulkarni, S. and Raja (2019). Evaluation of botanicals and bioagents against

- Colletotrichum truncatum* (SCHW.) Andrus and Moore, causing anthracnose of greengram. *Journal of Pharmacognosy and Phytochemistry*. 8(2): 2370-2373.
9. **Kulkarni, S. and Raja (2019)**. Integrated disease management of anthracnose of greengram. 8(3): 200-203.
  10. **Manjunath, B., Jayaram, N., Mallikarjuna, N., Sudarshan, G. K., Ravikumara, B. M. and Byregowda, M. (2016)**. Integrated management of anthracnose of field bean caused by *Colletotrichum lindemuthianum*. *Environment and Ecology*. 34 (3): 1013-1017.
  11. **Mishra, A., Pidurkar, S. K., Rathore, T. and Rai, G. (2024)**. Response of integrated nutrient management on growth and yield of black gram (*Vigna mungo* L. (Hepper) Egyptian *Journal of Agricultural Research*. 102(3):345-353.
  12. **Nandeesh, C.V., Akbari, L.F., Jaiswal, A., Harsha, B.R., Patil, B., Bhaliya, C.M., Kumar, N., Singh, T. and Jain, S. (2023)**. Control efficacy and yield response of different fungicides evaluated against anthracnose of green gram. *Crop Protection*. 174:0261-2194.
  13. **Purushotham, P., Rakholiya, K. B. and Vanani, K. D. (2023)**. Effectiveness of fungicides against *Colletotrichum lindemuthianum* causing anthracnose of greengram [*Vignaradiata* (L.) Wilczek]. *Journal of Advances in Biology & Biotechnology*. 27(1):94-103.
  14. **Reddy, T.Y. and Reddi, G.H. (1995)**. Principles of Agronomy (6<sup>th</sup> edition). *Kalyani Publishers*. Ludhiana, 72.
  15. **Saxena, R. M. and Gupta, J. S. (1984)**. Field survey for suspected seed transmitted disease of mungbean and urd bean in Uttar Pradesh. *Indian Phytopathology*. 34(3):340-345.
  16. **Sarkhosh, A., Schaffer, B., Vargas, A.I., Palmateer, A.J., Lopez, P. and Soleymani, A. (2017)**. *In vitro* evaluation of eight plant essential oils for controlling *Colletotrichum*, *Botryosphaeria*, *Fusarium* and *Phytophthora* fruit rots of avocado, mango and papaya. 54(3): 153–162.
  17. **Singh, H., Arora, A., Kaur, S. and Kaur, K. (2020)**. Evaluation of germplasm, fungicides and biocontrol agents against anthracnose (*Colletotrichum gloeosporioides*) in mango (*Mangifera indica*) nursery. *Indian Journal of Agricultural Sciences*. 90(6):1125-1129.
  18. **Tuite, J. (1969)**. Plant pathological methods: fungi and bacteria. *Burgess Publishing, Minnesota*. 239 p.
  19. **Vijaykumar, K. N., Kulkarni, S., Patil, P. V., Kambrekar, D. N. and Shashidhar, T. R. (2022)**. Eco-friendly management of purple blotch of garlic caused by *Alternaria porri* (Ellis) Ciferri. *Biological Forum - An International Journal*, 14(2): 1406-

1412.

**20. Wheeler, B. E. J. (1969).** An Introduction to Plant Diseases. *John Wiley and Sons Ltd., London.*

UNDER PEER REVIEW