

## Original Research Article

# Evaluation of selected plant extracts and carbendazim against stem rot of mustard (*Brassicajuncea* L.) caused by *Sclerotiniasclerotiorum* (Lib.) de Bary

---

### ABSTRACT

Sclerotinia stem rot disease caused by *Sclerotinia sclerotiorum* (Lib.) de Bary is a serious problem of mustard. The effect of five plant extracts using neem, garlic, eucalyptus, datura, and calotropis were tested against *S. sclerotiorum*. Maximum inhibition (%) of mycelial growth of *Sclerotinia sclerotiorum* was recorded in garlic bulb extract at 10%, followed by calotropis extract after 96 hours of inoculation. In a field experiment during rabi season 2023-24, seed treatment with carbendazim 2g/kg and foliar sprays of garlic bulb extract at 10% was most effective and recorded the lowest disease incidence (20.08%) of sclerotinia stem rot of mustard (*Brassica juncea* L.), maximum production (13.26 q/ha) and maximum cost-benefit ratio (1:1.87).

*Keywords:* Botanicals, carbendazim, mycelial growth, *Sclerotinia sclerotiorum*, stem rot,

### 1. INTRODUCTION

“Mustard (*Brassica juncea* L.) is the most important oil seed crop in India. It belongs to the family of Cruciferae. It is grown in certain tropical and subtropical regions as a cold-season crop. Mustard is the key source of income mainly for the marginal and small farmers cultivated mainly in the rain-fed and resource-vulnerable regions of the country” [18]. “Mustard is second in oilseed production, with soybean leading at 12.4 million metric tons, while mustard production is 11.8 million metric tons. The third most important source of edible oil next to soybean and palm oil” [8].

Sclerotinia stem rot was first reported by Shaw and Ajreker [15] from Pusa (Bihar). “Later on, the occurrence of this disease has also been observed in other parts of the country” [13]. Stem rot now become a serious problem in the mustard crop in northern India. It’s now economically important yield-reducing disease that has been widely reported in the last few years in India. This disease has now attained the next position to Alternaria blight in terms of its economy.

Plant extracts or botanical pesticides are essential part of an integrated pest control program because of low persistence, biodegradability and toxicity to mammals, making them ecologically beneficial. Therefore, study on the use of botanicals to treat plant diseases is necessary since they are widely available, environmentally friendly, reasonably priced, and safe for both human and animal health [20].

## 2. MATERIAL AND METHODS

### 2.1.1 Symptoms of stem rot of mustard

Symptoms of stem rot of mustard caused by *Sclerotinia sclerotiorum* were observed on lower leaves turning yellow and wilting, while a white, fluffy mold appeared around the base of the plant or on the stem collar [9]. As the disease advance, the stems become soft and can be peeled away easily. Water-soaked spots then form and grow into larger, sunken lesions that might completely encircle the stem, leading to plant collapse [6]. Black, irregularly shaped sclerotia, ranging from 2 to 12 mm, develop on the infected areas [3]. Wet conditions make this disease worse, and it tends to affect older plants more than young ones. Flowers and pods can also be affected, causing stunted growth and lower yields [17].

### 2.1.2 Morphological characters of *Sclerotinia sclerotiorum*

"*Sclerotinia sclerotiorum* hyphae are hyaline, septate, multinucleate, thin-walled 9-18  $\mu$ m in width and branching is never at right angles. Mycelia may appear white to tan in culture. Individual sclerotia are embedded in white mycelial net and are round, semi-spherical to irregular in shape, measuring 2-10 x 3-15 mm in size. Sexually produced apothecia are cup-shaped with concave discs, light yellowish brown, and vary in size from 2-11 mm (average 4-5mm) in diameter" [14].

### 2.1.3 Potato Dextrose Agar (PDA) medium:


The culture media used the experiment was prepared according to standard formula. For isolating and culturing of pathogen (*Sclerotinia sclerotiorum*) Potato Dextrose Agar (PDA) medium was used.

Procedure:

The potato was peeled and cut into small pieces and boiled in 500 ml of distilled water till they became soft. The extract obtained was filtered through muslin cloth and all the liquid was squeezed a beaker. Twenty grams of dextrose was added bit by bit to the remaining 500 ml of hot water. Then 20 g of agar were added to solidify. The volume of broth was made up to 1000 ml by adding more distilled water. Then in each conical flask 200 ml of this solution was dispensed and sterilized at 121°C at 15 lbs. pressure/square inch for 15 minutes in an autoclave.

### 2.1.4 Isolation of *Sclerotinia sclerotiorum*:

"Potato Dextrose Agar (PDA) was prepared and 80 mg of streptomycin, an antibiotic was added to each 500 ml preparation of the PDA to inhibit probable bacterial growth. The infected stem parts were cut into small pieces of two to three mm in dimension in a manner so that pieces may have some green portion also. Such stem bits were surface sterilized with 1 percent sodium hypochlorite (NaOCl) solution for 30 seconds and washed three times with sterile distilled water to remove any traces of sodium hypochlorite adhered with stem bits" [19]. Two to three stem bits were transferred on PDA medium contained in Petri plates aseptically with the help of sterilized forceps. These Petri plates were incubated at 25±1°C. After 3 days, mycelia growth was observed around stem bits from this colony growth, a portion from the periphery that is, a single hyphal tip was separated and transferred to the other.



**Plate 1. Fungal culture of *Sclerotinia sclerotiorum***

### 2.1.5 Preparation of plant extracts

The fresh leaves of selected plants and bulb extract were gently washed under running tap water and finally in sterile distilled water. They were separately grinded in sterile water at the rate of 1 ml/g of plant material in a pestle and mortar. Then they were filtered through a double layer of muslin cloth and finally through sterilized Whatman No.1 filter paper. This forms 100 % standard plant extract solution. Further, its dilution was performed of required concentration with sterilized water. By grinding 10 g of leaves in 100 ml of sterile water, 10 % of leaf extract can be obtained.

### 2.1.6 Poisoned food technique:

Five mm diameter of culture disc of *Sclerotinia sclerotiorum* was kept at the center of each Petri plate containing the botanicals of 10 percent concentration dissolved in PDA. Three replications were maintained. The plates were incubated at  $25 \pm 1^\circ\text{C}$  for seven days and colony diameter was recorded [21].

Percent inhibition of mycelial growth was calculated by using the formula:

$$I = \frac{C - T}{C} \times 100$$

where,

I = Per cent inhibition,

C=Growth (mm) of test fungus in untreated control plates,

T=Growth (mm) of test fungus in treated plates.

### 2.1.7 Mass multiplication of pathogen:

"The fungus inoculum was multiplied on sterilized sorghum grains. The sorghum grains were soaked in sterilized water overnight. The excess water drained out and 40 grams of grains were taken in each 250 ml conical flask and sterilized in autoclave at  $1.045 \text{ kg cm}^{-2}$  pressure

for 20 minutes. The sorghum grains in flasks were inoculated aseptically with 5 days old mycelial discs (5 mm) of the pathogen and inoculated for 15 days at 20±2 °C. The inoculum was mixed in rows during sowing, using 20 g of inoculum per meter of row" [2].

### 2.1.8 Disease incidence:

The incidence of disease was visually assessed in all the plots at 15 intervals from first appearance of the disease for each treatment. The data was analyzed statistically. Disease intensity was calculated by following formula:

$$\text{Disease incidence} = \frac{\text{Number of infected plants}}{\text{Total number of plants}} \times 100$$

### 2.1.9 Cost benefit ratio:

"Cost benefit ratio is the ratio of gross return to cost of cultivation, which can also be expressed as return per rupee invested. This index provides an estimate of the benefit a farmer derives from the expenditure incurs in adopting a particular cropping system. Any value above 2.0 is considered safe as the farmer gets Rs. 2 for every rupee invested. The cost benefit ratio was calculated using the formula" [12]:

$$\text{C:B ratio} = \frac{\text{Gross return}}{\text{Total Cost of treatment}}$$


## 3. RESULTS AND DISCUSSION

The *in vitro* screening aimed to find out the efficacy of selected botanicals against *Sclerotinia sclerotiorum* using a poisoned food technique. The experiment was analyzed by using CRD (Completely randomized design) with six treatments including control which were T<sub>1</sub>- neem leaf extract @ 10 %, T<sub>2</sub> -@ 10 %, T<sub>3</sub>- eucalyptus leaf extract @ 10 %, T<sub>4</sub> - datura leaf extract @ 10 %, T<sub>5</sub> – garlic bulb extract @ 10 %, T<sub>6</sub> –carbendazim @ 0.1% and T<sub>0</sub> – untreated check observations were recorded at 96 hours. Minimum radial growth at 96 hours was recorded in garlic bulb extract @ 10 % (37.50 mm) followed by calotropis leaf extract @ 10 % (44.37 mm), as compared to treated control carbendazim (30.87 mm) and untreated control (90.00 mm). Maximum percent inhibition of radial growth was recorded with garlic bulb extract @ 10 % (58.33) followed by calotropis leaf extract @ 10 % (50.70%), as compared to treated control carbendazim (65.70%) and untreated control (0.00%).


In the present studies, the maximum percent inhibition at 96 hours was recorded with garlic bulb extract followed by calotropis extract which was found effective over other treatments. The probable reasons may be the fungicidal properties of *Allium sativum* due to the presence of allicin and di-allyl sulphide [7]. Allicin is the most important biologically active substance of garlic, it is formed from its precursor, allin, by the action alliinase enzyme. Antifungal activities of allicin can be attributed to its interaction with the thiol group of proteins and amino acids and that, especially with the latter, allicin forms S-allyl derivatives [1]. These results were similar to the findings of Shivpuri and Gupta [16], Chattopadhyay *et al.* [5], Sharma *et al.*[14]. and Kewate *et al.*[10]. Maximum percent inhibition was recorded in garlic bulb extract against *Sclerotinia sclerotiorum*.

**Table 1. Mean colony diameter and per cent inhibition**

Treatment Number	Treatments	Concentration (%)	Meancolony diameter(mm)	Per cent inhibition (%)
T <sub>0</sub>	untreated control	-	90	0
T <sub>1</sub>	Neem leaf extract	10	61.96	31.15
T <sub>2</sub>	Calotropis leaf extract	10	44.37	50.70
T <sub>3</sub>	Eucalyptus leaf extract	10	85.65 <sup>a</sup>	5.94 <sup>a</sup>
T <sub>4</sub>	Datura leaf extract	10	83.90 <sup>a</sup>	6.77 <sup>a</sup>
T <sub>5</sub>	Garlic bulb extract	10	37.50	58.33
T <sub>6</sub>	Carbendazim	0.1	30.87	65.70
C.D. (5%)				0.825
SEM (±)				0.26



**Plate 2. Effect of selected botanicals on radial growth of *Sclerotinia sclerotiorum***



**Figure1. Effect of selected botanicals on per cent inhibition and radial growth of *Sclerotinia sclerotiorum***

The experiment was analysed by using RBD (randomized block design) with three replications having plot size  $2 \times 1 \text{ m}^2$  with six treatments including control which were T<sub>1</sub>- neem leaf extract @ 10 % (Foliar Spray) + carbendazim @ 2g/kg (Seed Treatment), T<sub>2</sub> – calotropis leaf extract @ 10 % (Foliar Spray) + carbendazim @ 2g (Seed Treatment), T<sub>3</sub> - eucalyptus leaf extract @ 10 % (Foliar Spray) + carbendazim @ 2g (Seed Treatment), T<sub>4</sub> - datura leaf extract @ 10 % (Foliar Spray) + carbendazim @ 2g (Seed Treatment), T<sub>5</sub> – garlic bulb extract @ 10 % (Foliar Spray) + carbendazim @ 2g (Seed Treatment), T<sub>6</sub> –carbendazim @ 2g (Seed Treatment) and T<sub>0</sub> – untreated control observations were recorded at 75 and 90 DAS.

Minimum per cent disease incidence (%) at 75 DAS was recorded in garlic bulb extract + carbendazim (14.28%) followed by neem leaf extract + carbendazim (16.66%) as compared to treated check carbendazim (20.27%) and untreated control (34.28%).

Minimum per cent disease incidence (%) at 90 DAS was recorded with garlic bulb extract + carbendazim (20.08%) followed by neem leaf extract + carbendazim (20.87%) as compared to treated check carbendazim (27.61%) and untreated control (42.85%).

Minimum number of sclerotia per stem at harvest was recorded with garlic bulb extract + carbendazim (2.13) followed by neem leaf extract + carbendazim (3.73), as compared to treated check carbendazim (13.26) and untreated check (16.13).


Maximum yield (q/ha) was recorded with garlic bulb extract + carbendazim (13.26q/ha) followed by neem leaf extract + carbendazim (12.81q/ha) as compared to treated check carbendazim (9.80q/ha) and untreated check (9.24q/ha).

Maximum cost benefit ratio was recorded with garlic bulb extract + carbendazim (1:1.87) followed by neem leaf extract + carbendazim (1:1.81) as compared to treated check carbendazim (1:1.47) and untreated check (1:1.44).

**Table2. Effect of carbendazim and selected botanical extracts on disease incidence (%), number of sclerotia and yield (q/ha) of mustard**

Tr. No.	Treatments	Per cent Disease Incidence (%)			Number of sclerotia	Yield (q/ha)
		60 DAS	75 DAS	90 DAS		
T <sub>0</sub>	Control (untreated check) Waterspray	28.57	34.28	42.85	16.13	9.24
T <sub>1</sub>	Carbendazim (ST) + Neem leaf extract @ 10% (FS)	11.12	16.66	20.87	3.73	12.81
T <sub>2</sub>	Carbendazim (ST) + Calotropis leaf extract (aak) @ 10% (FS)	12.14	18.09	22.85	6.40	12.10
T <sub>3</sub>	Carbendazim (ST) + Eucalyptus leaf extract @ 10% (FS)	14.28 <sup>a</sup>	19.62 <sup>a</sup>	25.23 <sup>a</sup>	11.00 <sup>a</sup>	10.75 <sup>a</sup>
T <sub>4</sub>	Carbendazim (ST) + Datura leaf extract @ 10% (FS)	13.88 <sup>a</sup>	19.44 <sup>a</sup>	25.07 <sup>a</sup>	10.53 <sup>a</sup>	11.07 <sup>a</sup>
T <sub>5</sub>	Carbendazim (ST) + Garlic bulb extract @ 10% (FS)	8.57	14.28	20.08	2.13	13.26
T <sub>6</sub>	Carbendazim (ST) (Treated check)	16.66	20.27	27.61	13.26	9.80
	C.D. (5%)	0.587	0.533	0.527		0.575
	C.V. (%)	1.568	1.469	1.123		1.568

The probable reasons may be the fungicidal properties of *Allium sativum* due to the presence of allicin and di-allyl sulphide. Allicin is the most important biologically active substance of garlic, it is formed from its precursor, allin, by the action alliinase enzyme. Antifungal activities of allicin can be attributed to its interaction with the thiol group of proteins and amino acids and that, especially with the latter, allicin forms S-allyl derivatives. These results were similar to the findings of Shivpuri and Gupta [16], Chattopadhyay *et al.* [5], Yadav [22], Sharma *et al.* [14] and Bharti *et al.* [4].



**Figure 2. Effect of treatments on disease incidence (%), number of sclerotia and yield (q/ha) of mustard**

#### 4. CONCLUSION

The present study concludes that the treatment with garlic bulb extract at 10% significantly inhibited the radial growth of *Sclerotinia sclerotiorum* (stem rot of mustard). The minimum disease incidence in stem rot of mustard (*Sclerotinia sclerotiorum*) was observed with garlic bulb extract at 10% (foliar spray). The yield and cost-benefit ratio were superiorly recorded in the treatment of garlic bulb extract at 10% (foliar spray) + carbendazim (seed treatment) when compared to the treated check and the untreated check. The present investigation was limited to one crop season (Rabi), under the climatic conditions of Prayagraj (U.P.). Therefore, to substantiate the present result, more such trials are required for further recommendation.

#### Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### REFERENCES

1. Aala, F., Yusuf, U. K., Nulit, R. and Rezaie, S. (2014). Inhibitory effect of allicin and garlic extracts on growth of cultured hyphae. *Iranian Journal of Basic Medical Sciences*, 17(3), 150.
2. Bairwa, S.K., Godara, S.L. and Meena, S. (2015). *In vitro* efficacy of fungicides on *Sclerotinia sclerotiorum* and their potential for control of stem rot in Indian mustard (*Brassica juncea* L.). *International Journal of Bio-Resource and Stress Management*. 6(4):497-502.

3. **Baribeau, B. (1923).** Stem wilt and rot of sunflower in the province of Quebec. *Science et Agriculture*. 3(11): 397-400.
4. **Bharti, O. P., Pandya, R. K., Gupta, J. C., Singh, R. K. and Tripathi, A. K. (2021).** Management of stem rot of mustard incited *Sclerotinia sclerotiorum* (Lib.) de Bary. *International Journal of Agriculture Sciences*. 13(1): 10578-10581.
5. **Chattopadhyay, C., Kumar, V. R. and Meena, P. D. (2007).** Bio management of sclerotinia rot of *Brassica juncea* in india- a case study. *Phytomorphology*. 57(1/2): 71-83.
6. **Collmer, A. and Keen, N. T. (1986).** The role of pectic enzymes in plant pathogenesis. *Annual Review of Phytopathology*. 24: 383-409.
7. **El-Saber Batiha, G., Beshbishy, A. M., Wasef, L. G., Elewa, Y. H., Al-Sagan, A., Abd El-Hack, M. E. and Prasad Devkota, H. (2020).** Chemical constituents and pharmacological activities of garlic (*Allium sativum* L.): A review. *Nutrients*, 12(3), 872.
8. <https://gain.fas.usda.gov>
9. **Joshi, S. D. (1924).** The wilt disease of safflower. *Botanical Series*. 13: 39-46.
10. **Kevwate, B., Singh, D., Singh, V. D., Malik, N. P. and Kumar, R. (2020).** *In-vitro* effect evaluation of botanicals against *Sclerotinia sclerotiorum* (Lib.) de Bary, caused stem rot disease in rapeseed- mustard. *International Journal of Current Microbiology and Applied Sciences*. 9(8): 3733-3741.
11. **Kolte, S. J. (1988).** Diseases of annual edible oilseed crops. *Rapeseed-Mustard and Sesame Diseases*. CRC Press, Inc., Florida. 2: 35-39.
12. **Reddy, T. Y. and Reddi, G. H. S. (2004).** Principles of Agronomy III<sup>rd</sup> edition. *Kalyani Publisher*, pp: 527.
13. **Roy, A. K. and Saikia, U. N. (1976).** White blight of mustard and its control. *Indian Journal of Agricultural Sciences*. 46: 274-277.
14. **Sharma, J., Godika, S., Ghasolia, R. P., Meena, S. and Yadav, A. L. (2016).** Fungitoxicity of plant extracts against sclerotinia rot of indian mustard incited by *Sclerotinia sclerotiorum*. *Indian Phytopathology*. 69(4): 387-390.
15. **Shaw, F. J. F. and Ajrekar, S. L. (1915).** The genus *Rhizoctonia* in India. Department of Agriculture, Indian Botanical Survey. 7: 177-194.
16. **Shivpuri, A. and Gupta, R. B. L. (2001).** Evaluation of different fungicides and plant extracts against *Sclerotinia sclerotiorum* causing stem rot of mustard. *Indian Phytopathology*. 54(2): 272-274.
17. **Sikarwar, N. S. (1987).** Studies on white blight of mustard caused by *Sclerotinia sclerotiorum* (Lib.) De bary. Ph.D. Thesis, Jiwaji University, Gwalior, MP. 238-249.
18. **Tripathi, D. K., Singh, R. and Prajapati, A. K. (2023).** Evaluation of fungicide and plant extracts against stem rot of mustard caused by *Sclerotinia sclerotiorum* *in-vitro* and *in-vivo* condition. *Journal of Agri search*. 10(1): 65-70.
19. **Tuite, J. (1969).** Plant pathological methods: fungi and bacteria. Burgess Publishing. *Minnesota*. pp: 239.
20. **Upadhyay, P. and Tewari, A. K. (2019).** Evaluation of botanical extracts, animal wastes, organic and inorganic salts, micronutrients and bio-agents against *Sclerotinia sclerotiorum* (lib) de Bary a cause of sclerotinia rot of rapeseed mustard under field conditions. *Bulletin of Environment, Pharmacology and Life Sciences*. 8(12): 60-65.
21. **Vincent, J. M. (1947).** Distortion of fungal hyphae in the presence of certain inhibitors. *Nature*. 159(4051): 850.
22. **Yadav, M. S. (2009).** Bio-pesticidal effect of botanicals on the management of mustard diseases. *Indian Phytopathology*. 62(4): 488-492.

UNDER PEER REVIEW