

A review on Role of Root Exudates in Shaping Plant-Microbe-Pathogen Interactions

Abstract

Root exudates are diverse compounds secreted by plant roots that significantly influence the rhizosphere-where complex interactions among plants, microbes, and pathogens occur. These exudates include sugars, amino acids, organic acids, phenolics, and volatile organic compounds, which collectively shape the microbial community structure in the soil. By attracting beneficial microbes like rhizobia, arbuscular mycorrhizal fungi (AMF), and plant growth-promoting rhizobacteria (PGPR), root exudates enhance nutrient acquisition, promote plant growth, and improve resistance to environmental stresses. Exudates can directly inhibit pathogens through the release of antimicrobial compounds such as benzoxazinoids and phenolics or indirectly by priming plant immune responses and inducing systemic resistance. Environmental factors like nutrient availability, soil pH, texture, and abiotic stressors (e.g., drought, salinity, and heavy metals) profoundly impact the composition and quantity of root exudates, thereby influencing their ecological functions. Phosphorus and iron deficiencies trigger the secretion of organic acids to mobilize these nutrients, while drought and salinity alter exudation patterns to recruit drought-tolerant microbes. Despite their potential, studying root exudates presents technical challenges, such as isolating exudates in natural soil systems and analyzing them with sufficient precision. Knowledge gaps also exist regarding the temporal dynamics of exudation and how these compounds influence multi-species plant-microbe interactions under field conditions. Future research should focus on breeding crops with optimized root exudate profiles for enhanced nutrient uptake, pathogen resistance, and resilience to climate change. Leveraging root exudates in developing biofertilizers, biostimulants, and integrated pest management (IPM) strategies can reduce the reliance on chemical fertilizers and pesticides, promoting sustainable agriculture. By manipulating root exudates, we can improve soil health, increase crop productivity, and support sustainable farming practices in a changing climate. As global agriculture faces challenges from soil degradation, climate variability, and food security demands, root exudates offer an eco-friendly approach to enhancing soil fertility, boosting plant resilience, and reducing agricultural inputs, thus contributing to more sustainable and resilient agroecosystems.

Keywords: *Root exudates, rhizosphere, plant-microbe interactions, biofertilizers, nutrient acquisition*

I. Introduction

A. Root Exudates and Their Significance

Root exudates are a diverse group of chemical compounds released by plant roots into the surrounding soil environment, collectively known as the rhizosphere. These compounds include sugars, amino acids, organic acids, phenolics, flavonoids, fatty acids, nucleotides, and various secondary metabolites [1]. The secretion of root exudates is a dynamic and continuous process, which is vital for plant growth and development as well as for the maintenance of soil health. Root exudates serve multiple functions in the soil ecosystem. They act as nutrient sources for microorganisms, influence soil structure, and mediate interactions between plants and soil organisms. Exudates play a critical role in nutrient acquisition, such as phosphorus solubilization, by altering the chemical environment in the rhizosphere. The chemical composition of root exudates varies according to plant species, developmental stages, and environmental conditions. This variability is important for determining the structure and diversity of microbial communities in the rhizosphere [2]. The secretion of root exudates can be divided into several categories based on their mode of release: passive diffusion, active secretion, and root cell lysis. Passive diffusion is responsible for releasing low-molecular-weight compounds like amino acids and sugars, while active secretion involves energy-dependent processes for more complex molecules, such as secondary metabolites.

B. The Rhizosphere as a Dynamic Zone of Interactions

The rhizosphere is defined as the narrow region of soil that is directly influenced by root secretions and associated soil microorganisms [3]. It is a highly dynamic zone where plant roots, soil microbes, and pathogens interact continuously. The rhizosphere can extend a few millimeters to centimeters from the root surface and is characterized by high microbial density and activity compared to the bulk soil. Microorganisms in the rhizosphere include bacteria, fungi, archaea, and other soil organisms that form complex communities. These microbial communities are influenced not only by the types of root exudates released but also by soil properties, such as pH, temperature, and nutrient content. Beneficial microbes, such as Plant Growth-Promoting Rhizobacteria (PGPR) and mycorrhizal fungi, enhance plant health by improving nutrient uptake, producing phytohormones, and protecting against pathogens. The rhizosphere is also a battleground where pathogenic microorganisms attempt to colonize roots, often leading to plant diseases [4]. Pathogens like *Fusarium*, *Pythium*, and *Ralstonia* spp. exploit root exudates as signals to locate their host plants. Conversely, plants can modify their exudate profiles in response to pathogen attack, thereby recruiting beneficial microbes to outcompete or inhibit pathogens.

C. Importance of Plant-Microbe-Pathogen Interactions in the Sustainable Agriculture

In recent years, there has been a growing recognition of the importance of understanding plant-microbe-pathogen interactions in sustainable agriculture. These interactions are important for developing agricultural practices that reduce the reliance on synthetic fertilizers and pesticides, which have adverse environmental impacts. Root exudates serve as a bridge between plants and their microbial partners, facilitating beneficial symbioses that can enhance nutrient availability, improve soil health, and increase crop resilience to stress [5]. The application of beneficial microbes, such as PGPR and mycorrhizal fungi, as bioinoculants has shown promise in increasing crop productivity while reducing the environmental footprint of agriculture. For example, root exudates from legumes stimulate the growth of nitrogen-fixing *Rhizobia*, which reduce the need for nitrogen fertilizers. The release of strigolactones by crop roots enhances phosphorus uptake through mycorrhizal associations. By understanding how root exudates shape the rhizosphere microbiome, researchers can develop strategies to manipulate these interactions in favor of plant health. This can lead to innovations in sustainable farming practices, such as the use of microbial inoculants, crop rotations, and exudate-enhancing cultivars to suppress soil-borne diseases and enhance nutrient cycling [6].

D. Objectives of the Review and Its Relevance to Current Research

The primary objective of this review is to explore the multifaceted roles of root exudates in shaping plant-microbe-pathogen interactions. This includes an in-depth analysis of how different types of exudates influence the recruitment of beneficial microbes and the suppression of soil-borne pathogens. The review will cover the impact of environmental factors on root exudation patterns and the potential applications of root exudate management in sustainable agriculture. Recent advances in molecular biology and analytical chemistry have enabled a deeper understanding of root exudate composition and function [7]. There are still significant gaps in our knowledge regarding the specific mechanisms by which root exudates mediate complex interactions in the rhizosphere. This review aims to bridge these knowledge gaps by synthesizing current research and identifying future directions for studies on root exudates. By elucidating the mechanisms by which plants, microbes, and pathogens interact through root exudates, we can leverage this knowledge to enhance crop productivity and resilience in the face of climate change and other global challenges. Understanding these interactions is critical for developing eco-friendly agricultural practices that support long-term soil health and food security.

II. Composition and Diversity of Root Exudates

Root exudates are a complex blend of compounds released by plant roots into the surrounding soil, significantly influencing soil microbial communities, nutrient availability, and plant health [8]. These compounds are diverse, encompassing a wide range of chemical structures and functions. The composition of root exudates varies across plant species, developmental stages, and environmental conditions, making them an important factor in the interactions between plants, soil microbes, and pathogens.

A. Classification of Root Exudates

Root exudates can be broadly classified into four categories: low molecular weight compounds, high molecular weight compounds, secondary metabolites, and volatile organic compounds (VOCs) (Table 1). These categories capture the diversity of the chemical structures and biological functions of the exudates. Low molecular weight compounds are one of the most significant components of root exudates and include sugars, amino acids, and organic acids. Sugars such as glucose, fructose, and sucrose are released into the rhizosphere and are quickly utilized by soil microorganisms, serving as an energy source that enhances microbial proliferation [9]. These sugars not only fuel microbial metabolism but also shape the composition of microbial communities, promoting the growth of beneficial bacteria like *Pseudomonas* and *Bacillus* species that enhance plant health. Amino acids, such as glutamine and asparagine, play a dual role in providing nitrogen to soil microbes and in signaling beneficial bacteria to colonize root surfaces. Organic acids, including citric, malic, and oxalic acids, are important for nutrient solubilization. The release of malic acid by maize roots has been shown to increase the availability of phosphorus by chelating soil-bound phosphates. This nutrient mobilization is especially important in nutrient-poor soils where phosphorus availability is limited. High molecular weight compounds, such as polysaccharides, mucilage, and proteins, also play an essential role in root exudation. Polysaccharides secreted as mucilage help in soil aggregation, thereby enhancing soil structure and water retention [10]. This mucilage forms a protective layer around roots, reducing water loss and protecting roots from desiccation, particularly in arid environments. Proteins, on the other hand, include enzymes and signaling molecules that facilitate interactions with soil microbes. Enzymes like phosphatases secreted by roots enhance the breakdown of organic phosphorus compounds, making them accessible for plant uptake. Lectins and other root-secreted proteins promote microbial adherence to root surfaces, fostering beneficial symbiotic relationships. Secondary metabolites encompass a wide range of bioactive compounds such as phenolics, terpenoids, flavonoids, saponins, and alkaloids [11]. These compounds are not directly involved in plant growth but serve important roles in plant defense and microbial signaling. Phenolic compounds like tannins and lignins act as antimicrobial agents, reducing the presence of harmful pathogens in the rhizosphere. For example, the secretion of catechin by *Centauria maculosa* roots has been shown to inhibit fungal pathogens. Terpenoids and flavonoids are particularly important in symbiotic interactions. In legumes, flavonoids induce the expression of *nod* genes in *Rhizobia*, leading to the formation of nitrogen-fixing nodules on the roots [12]. This specific signaling ensures that only compatible rhizobial strains initiate symbiosis, enhancing nitrogen fixation efficiency. Volatile organic compounds (VOCs) represent a unique category of root exudates because they are small, volatile molecules that can travel through the soil atmosphere, influencing both microbial communities and neighboring plants. For example, terpenes and aldehydes released by tomato roots can suppress the growth of pathogenic fungi like *Rhizoctonia solani*. These VOCs serve as long-distance chemical signals that modulate the behavior of soil organisms, thus playing an important role in root defense.

Table: 1 Types of Root Exudates and Their Functions

Exudate Type	Primary Functions	Examples
Sugars	Serve as a readily available energy source for	Glucose, sucrose, fructose.

	soil microbes.	
Amino Acids	Attract beneficial microbes; involved in nitrogen cycling.	Glycine, asparagine, glutamate.
Organic Acids	Solubilize soil nutrients (e.g., phosphorus) and modulate pH levels.	Citric acid, malic acid, oxalic acid.
Phenolics	Antimicrobial properties; act as signaling molecules for symbionts.	Flavonoids, tannins, lignin precursors.
Secondary Metabolites	Function in plant defense and microbial signaling.	Terpenoids, alkaloids, phytoalexins.

B. Factors Influencing Root Exudate Composition

The diversity and concentration of root exudates are influenced by a variety of factors, including plant species, developmental stage, environmental conditions, and biotic stresses (Table 2). Plant species have distinct exudate profiles that contribute to the establishment of specific microbial communities in the rhizosphere [13]. Legumes are known for their high secretion of flavonoids, which are critical for establishing symbiotic relationships with *Rhizobia*. In contrast, cereals like maize release a higher proportion of organic acids that enhance nutrient solubilization in nutrient-poor soils. The developmental stage of the plant also affects exudate composition. Seedlings tend to release more sugars and amino acids to support rapid root growth and the establishment of beneficial microbial communities. In mature plants, the focus shifts towards the secretion of secondary metabolites that protect against soil-borne pathogens. Environmental factors, such as soil type, pH, and nutrient availability, significantly influence root exudation patterns. Acidic soils can promote the exudation of organic acids like citric acid to solubilize bound minerals, enhancing nutrient availability. Phosphorus deficiency triggers increased exudation of organic acids to mobilize unavailable phosphorus from soil complexes. Soil texture and moisture content also affect the diffusion and persistence of root exudates in the rhizosphere, influencing the microbial community structure [14]. Biotic stresses, including the presence of pathogens and beneficial microbes, can alter the exudation profile of plants. For example, Arabidopsis plants secrete antimicrobial compounds like camalexin in response to pathogen attack to inhibit the growth of pathogens such as *Pythium*. Conversely, the presence of beneficial mycorrhizal fungi stimulates the release of strigolactones, which enhance fungal colonization and nutrient exchange.

Table: 2 Factors Influencing Root Exudate Composition

Factor	Details	Examples
Plant Species/Genotype	Root exudate profiles vary across species and cultivars due to genetic differences.	Legumes release flavonoids, while grasses exude phenolics.
Plant Age/Development	Younger plants release more exudates, with composition changing over the plant's lifecycle.	Seedlings exude sugars and amino acids, while mature plants shift to phenolics.
Soil Type	Soil characteristics (texture, pH, organic matter) affect the quantity and nature of exudates.	Acidic soils may promote the release of organic acids.
Microbial	Beneficial microbes in the rhizosphere	Rhizobia increase flavonoid release

Communities	can influence root exudation patterns.	in legume roots.
Nutrient Availability	Plants alter exudate composition in response to nutrient levels in soil.	Phosphorus deficiency enhances organic acid exudation to mobilize P.
Abiotic Stress	Environmental stresses (drought, salinity, heat) can increase or modify exudate release.	Drought stress induces the release of osmoprotectants like proline.
Biotic Stress	Herbivory, pathogen attacks, and plant competition can trigger defensive exudates.	Pathogen infection can increase exudation of antimicrobial compounds.
Root Architecture	The spatial distribution of roots influences exudate release patterns and rhizosphere effects.	Fine roots release more exudates than thicker, woody roots.
Photosynthetic Activity	Higher carbon fixation rates lead to more carbon allocation to root exudates.	Elevated CO ₂ levels can boost exudate production, impacting soil microbes.
Circadian Rhythm	Some plants exhibit diurnal rhythms in exudate release linked to light/dark cycles.	Exudation of sugars may peak during the day to support microbial symbionts.

C. Methods for Studying Root Exudates

Studying root exudates is challenging due to their diverse chemical nature and the complexity of soil environments. Various methods have been developed to collect and analyze root exudates, including hydroponics, soil leaching, and the use of root chambers. Hydroponic systems allow for the collection of root exudates in a liquid medium, thus avoiding soil interference, but may not fully capture the natural exudation profiles seen in soil [15]. Soil leaching involves washing soil with solvents to extract exudates, providing a more field-relevant representation, though it can be affected by soil properties. To analyze the collected exudates, advanced analytical techniques like Gas Chromatography-Mass Spectrometry (GC-MS), Liquid Chromatography-Mass Spectrometry (LC-MS), and Nuclear Magnetic Resonance (NMR) spectroscopy are used. GC-MS is highly effective for identifying volatile compounds, while LC-MS is better suited for non-volatile and high molecular weight compounds like phenolics and flavonoids. NMR spectroscopy provides detailed structural information, making it invaluable for characterizing complex secondary metabolites [16].

III. Root Exudates and Beneficial Microbial Interactions

Root exudates play an important role in mediating interactions between plants and beneficial soil microorganisms. These interactions are important for plant health, nutrient acquisition, stress resilience, and overall growth. By selectively releasing specific chemical compounds, plant roots create a favorable environment in the rhizosphere that supports the recruitment and colonization of beneficial microbes.

A. Role of Root Exudates in Recruiting Beneficial Microbes

Root exudates serve as powerful signals and chemoattractants that influence the behavior of soil microbes, enhancing the establishment of mutualistic relationships that benefit the plant. Three

prominent examples include the recruitment of *Rhizobia* by legumes, the stimulation of arbuscular mycorrhizal fungi (AMF), and the attraction of plant growth-promoting rhizobacteria (PGPR) [17].

1. Rhizobia-Legume Symbiosis: Role of Flavonoids in Nod Gene Induction

Legumes engage in a well-studied mutualistic interaction with nitrogen-fixing bacteria known as *Rhizobia*. The establishment of this symbiosis is mediated by the secretion of flavonoids from legume roots, which act as signaling molecules. Flavonoids, such as genistein and luteolin, are specifically recognized by *Rhizobia*, leading to the induction of bacterial *nod* genes responsible for the production of Nod factors. These Nod factors trigger the formation of root nodules, specialized structures where *Rhizobia* fix atmospheric nitrogen into ammonia, which is then utilized by the host plant. This symbiotic relationship significantly enhances nitrogen availability in soils, reducing the need for synthetic nitrogen fertilizers [18]. The specificity of the flavonoid-Nod factor signaling ensures that legumes selectively interact with compatible *Rhizobia* strains, optimizing the efficiency of nitrogen fixation. For example, soybean roots release specific flavonoids that attract *Bradyrhizobium japonicum*, a highly effective nitrogen-fixing bacterium. This selective recruitment has profound implications for agricultural practices, as it can improve soil fertility and reduce the environmental impact of nitrogen fertilizers.

2. Arbuscular Mycorrhizal Fungi (AMF): Stimulation by Strigolactones

Arbuscular mycorrhizal fungi (AMF) form symbiotic associations with the roots of over 80% of terrestrial plant species. The establishment of this symbiosis is facilitated by the release of strigolactones from plant roots, which stimulate spore germination and hyphal branching in AMF. Strigolactones are sesquiterpene lactones that act as signaling molecules, enhancing the colonization of roots by AMF. The AMF-plant symbiosis is particularly beneficial for phosphorus uptake, as these fungi extend the root system's absorptive surface area, enabling better access to phosphate ions that are otherwise immobile in the soil [19]. In nutrient-poor soils, plants that form symbiotic relationships with AMF show enhanced growth and resilience due to improved phosphorus and water uptake. Maize plants colonized by AMF demonstrate increased resistance to drought stress, highlighting the role of root exudates in modulating plant responses to environmental stressors.

3. Plant Growth-Promoting Rhizobacteria (PGPR): Attractants and Signaling Molecules

PGPR are a diverse group of bacteria that colonize plant roots and promote growth by various mechanisms, including nutrient solubilization, phytohormone production, and pathogen suppression. Root exudates rich in sugars, amino acids, and organic acids attract PGPR such as *Pseudomonas*, *Bacillus*, and *Azospirillum* species. These microbes use exudates as carbon sources, which in turn enhance their colonization of root surfaces. For example, *Pseudomonas fluorescens* responds to malic acid exuded by Arabidopsis roots, which enhances bacterial chemotaxis and biofilm formation, ultimately promoting plant growth and systemic resistance [20]. PGPR produce secondary metabolites that suppress soil-borne pathogens, thereby contributing to plant health.

B. Mechanisms by Which Beneficial Microbes Utilize Root Exudates

Beneficial microbes utilize root exudates to support their metabolism, promote plant growth, and protect the host from biotic and abiotic stresses. The utilization of exudates is linked to several key processes that improve plant fitness.

1. Nutrient Acquisition (Phosphorus Solubilization, Nitrogen Fixation)

Root exudates play an important role in nutrient cycling in the rhizosphere. Organic acids such as citric and oxalic acid chelate minerals, making phosphorus more available to plants. *Bacillus* and *Pseudomonas* species utilize organic acids to solubilize phosphate, enhancing its uptake by the plant.

Nitrogen-fixing bacteria like *Rhizobia* convert atmospheric nitrogen into ammonia, which plants can readily assimilate, significantly boosting nitrogen availability in agricultural systems [21].

2. Production of Phytohormones (Auxins, Gibberellins, Cytokinins)

PGPR can synthesize phytohormones such as indole-3-acetic acid (IAA), gibberellins, and cytokinins in response to root exudates. These hormones influence root architecture, leading to increased root surface area and enhanced nutrient uptake. For example, IAA production by *Azospirillumbrasilense* has been shown to stimulate lateral root formation, improving nutrient acquisition in maize.

3. Enhancement of Stress Tolerance (Drought, Salinity, Heavy Metals)

Microbial interactions mediated by root exudates can enhance plant resilience to abiotic stresses. PGPR like *Bacillus* and *Pseudomonas* spp. produce osmoprotectants and antioxidant enzymes that protect plants from oxidative stress caused by drought and salinity. Certain PGPR can sequester heavy metals, reducing their toxicity to plants and enabling growth in contaminated soils [22].

C. Impact of Microbial Colonization on Plant Health and Growth

The colonization of roots by beneficial microbes, facilitated by root exudates, has a profound impact on plant health and productivity. Enhanced nutrient uptake, increased tolerance to environmental stress, and protection against pathogens are some of the key benefits conferred by microbial symbioses. For example, tomato plants inoculated with *Pseudomonas fluorescens* exhibit increased resistance to *Fusarium wilt*, a common soil-borne disease. Beneficial microbes modulate plant immune responses through induced systemic resistance (ISR). This form of resistance is primed by root exudates that stimulate microbial production of ISR-inducing metabolites, thereby preparing the plant to better respond to future pathogen attacks [23].

IV. Root Exudates and Plant Pathogen Defense

Plants face constant threats from soil-borne pathogens, including fungi, bacteria, and nematodes. In response, they have evolved sophisticated defense mechanisms, many of which are mediated by root exudates. These exudates not only serve as signals to beneficial microbes but also play an important role in the direct and indirect defense against pathogens. By modulating the chemical environment of the rhizosphere, root exudates help plants suppress pathogen growth and activate their immune responses, thus contributing to enhanced plant resilience and health [24].

A. Direct Defense Mechanisms via Exudates

Root exudates can exert direct antimicrobial effects, either by releasing compounds that inhibit pathogen growth or by altering the rhizosphere environment to make it hostile to potential invaders.

1. Allelopathy: Release of Antimicrobial Compounds (Benzoxazinoids, Saponins)

Allelopathy refers to the ability of plants to release biochemicals that suppress the growth of competing organisms, including pathogens. Several root exudates serve as allelopathic agents with potent antimicrobial properties. For example, benzoxazinoids, a class of secondary metabolites produced by maize roots, have been shown to inhibit the growth of soil-borne fungi such as *Fusarium* and *Pythium*. These compounds are particularly effective in reducing the incidence of seedling diseases, thereby promoting crop establishment. Saponins, another group of allelopathic compounds, are released by a wide range of plant species, including legumes and oats. These compounds disrupt the cell membranes of pathogenic fungi and bacteria, leading to cell lysis and death [25]. Saponin-producing plants have been found to have reduced levels of root rot caused by pathogens like *Phytophthora* spp.. The secretion of these allelopathic substances not only protects the roots from infection but also enhances the competitive advantage of the plant in pathogen-rich soils.

2. Inhibition of Pathogen Growth (Phenolics, Alkaloids)

Phenolic compounds such as flavonoids, tannins, and lignin derivatives are commonly exuded by plant roots and have been shown to possess antimicrobial properties. Phenolics can interfere with the cell wall synthesis of fungal pathogens, thereby inhibiting their growth and spore germination. Camalexin, a phytoalexin released by *Arabidopsis thaliana*, is known to be highly effective against *Pythium* spp., a notorious pathogen that causes root rot [26]. Alkaloids, such as nicotine and berberine, are another class of antimicrobial compounds secreted by certain plant species. These compounds can inhibit bacterial and fungal pathogens by disrupting their protein synthesis and enzyme activity. For example, berberine has been shown to be effective against *Ralstonia solanacearum*, a bacterial pathogen that causes wilt diseases in solanaceous crops.

B. Indirect Defense Mechanisms

In addition to direct antimicrobial action, root exudates can also modulate plant immune responses, enhancing the plant's ability to resist infections. This is achieved through mechanisms such as induced systemic resistance (ISR) and immune priming.

1. Induced Systemic Resistance (ISR) by Root Exudates

Induced systemic resistance (ISR) refers to a plant's enhanced defensive capacity against a broad spectrum of pathogens, triggered by beneficial rhizosphere microbes such as PGPR. Root exudates like malic acid, which attract beneficial microbes like *Pseudomonas fluorescens*, play an important role in ISR activation [27]. These bacteria colonize the root surface and secrete metabolites that induce ISR, leading to the systemic activation of plant defense genes. Plants with activated ISR often exhibit increased levels of pathogenesis-related proteins, phytoalexins, and other defensive enzymes that inhibit pathogen colonization. This mechanism is particularly important for protecting plants against foliar pathogens, as ISR provides systemic protection even in parts of the plant that are not in direct contact with the pathogen.

2. Priming of Plant Immune Responses Through Microbial Metabolites

Root exudates can also enhance the effectiveness of plant defenses through a process known as immune priming. Primed plants exhibit a faster and stronger defensive response when exposed to pathogens. For example, exudates such as organic acids and amino acids released in response to microbial colonization can prime the plant immune system by enhancing the accumulation of defensive metabolites [28]. Microbial metabolites produced in response to root exudates, such as lipopeptides from *Bacillus* species, can act as elicitors that trigger immune responses in plants. This interaction not only improves the plant's resistance to pathogens but also enhances its ability to cope with abiotic stressors like drought and salinity.

C. Case Studies of Specific Pathogens and Root Exudate Responses

To better understand the role of root exudates in plant defense, it is useful to examine specific cases where exudates have been shown to affect interactions with pathogens.

1. Interaction with Fungal Pathogens (*Fusarium*, *Pythium*)

Fungal pathogens such as *Fusarium oxysporum* and *Pythium ultimum* are major threats to crop production, causing diseases like wilt and root rot. Studies have shown that maize roots release benzoxazinoids, which inhibit the growth of *Fusarium* spores, reducing infection rates [29]. *Arabidopsis* plants secrete phenolic compounds that are toxic to *Pythium*, limiting its ability to colonize the root system. Certain crops, like wheat, can exude antifungal compounds in response to pathogen attack, a process that is enhanced by microbial inoculants like *Trichoderma* spp., which in turn boost plant defenses.

2. Effects on Bacterial Pathogens (*Ralstonia*, *Pseudomonas*)

Bacterial pathogens such as *Ralstonia solanacearum* and *Pseudomonas syringae* can cause devastating wilt and blight diseases in economically important crops. Root exudates play a critical role in controlling these bacterial pathogens. Tomato plants release phenolics and alkaloids in response to *Ralstonia* infection, which inhibits the pathogen's virulence. The recruitment of beneficial PGPR through specific root exudates can suppress *Pseudomonas* infections by inducing systemic resistance [30]. The application of *Pseudomonas fluorescens* as a bioinoculant has been shown to reduce bacterial speck disease in tomatoes by enhancing root exudation of organic acids that support the growth of protective microbial communities.

V. Influence of Environmental Factors on Root Exudation

Root exudation is a dynamic process that plants use to interact with their surrounding soil environment. The composition and quantity of root exudates are highly sensitive to various environmental factors, including soil nutrient levels, pH, soil texture, and abiotic stressors like temperature, drought, and heavy metals. These factors can significantly alter the exudate profile, influencing microbial communities in the rhizosphere, plant growth, and resilience to stress [31].

A. Soil Nutrient Levels and Their Impact on Exudate Composition

Plants adjust their root exudation patterns based on the availability of essential nutrients in the soil. This adjustment helps optimize nutrient uptake and recruit beneficial soil microorganisms that can enhance nutrient availability.

1. Phosphorus and Iron Availability

Phosphorus (P) is a critical but often limiting nutrient in many soils due to its low mobility and strong fixation to soil particles. In response to P deficiency, plants increase the secretion of organic acids such as citric, malic, and oxalic acids to mobilize bound phosphates. For example, *Arabidopsis thaliana* roots release higher levels of malic acid under P-deficient conditions, which chelates calcium-bound phosphates, making them more available for root uptake [32]. This strategy is particularly important in acidic soils, where phosphorus is tightly bound to aluminum and iron oxides. Iron (Fe) deficiency induces the release of exudates like phenolics and flavonoids that can chelate Fe^{3+} ions, reducing them to the more soluble Fe^{2+} form, which is easier for plant uptake. Studies on *Strategy I* plants (non-graminaceous species) like tomato have shown increased secretion of phenolic compounds such as caffeic acid under iron-limited conditions, enhancing iron solubility and absorption.

2. Drought and Salinity Stress

Drought and salinity are major abiotic stressors that can significantly impact root exudation patterns. Under drought conditions, plants increase the secretion of osmoprotective compounds like proline, glycine betaine, and certain sugars to protect root cells from osmotic stress. These compounds not only help in maintaining cell turgor but also attract beneficial microbes like *Bacillus* spp. that enhance drought resilience [33]. Salinity stress leads to the accumulation of sodium ions in the soil, which can be toxic to plants. To cope with high salinity, plants modify their root exudate profiles by increasing the release of organic acids and amino acids that help in ion detoxification and stress mitigation. Increased exudation of organic acids like citric acid has been observed in barley under saline conditions, which aids in the chelation of toxic ions and supports beneficial microbial colonization.

B. Influence of Soil pH and Texture

Soil pH and texture are key determinants of the bioavailability of nutrients and the behavior of root exudates in the soil. Acidic soils (pH < 5.5) often lead to the increased release of organic acids by

plant roots to neutralize the soil environment and mobilize nutrients like phosphorus and iron. In contrast, alkaline soils (pH > 7.5) may enhance the exudation of amino acids and phenolics to cope with nutrient imbalances and metal toxicity [34]. Soil texture influences the diffusion rate of root exudates. Sandy soils, with larger pore spaces, allow for faster diffusion of exudates, whereas clayey soils, with smaller pores, tend to retain exudates near the root zone, leading to higher microbial activity in the rhizosphere. For example, the retention of root exudates in clay-rich soils can enhance the growth of mycorrhizal fungi, promoting nutrient uptake and plant health.

C. Impact of Abiotic Stressors (Temperature, Heavy Metals) on Root Exudate Profiles

Abiotic stressors such as temperature fluctuations and heavy metal contamination can drastically alter root exudation, influencing plant-microbe-pathogen interactions. High temperatures can increase the metabolic rate of plants, leading to a surge in the release of sugars, amino acids, and other low molecular weight compounds [35]. These compounds can enhance the activity of rhizosphere microbes that contribute to soil nutrient cycling, although prolonged heat stress may deplete root energy reserves, reducing exudation over time. Heavy metal stress, such as exposure to cadmium (Cd) and lead (Pb), prompts the secretion of chelating agents like organic acids and phenolics to detoxify the soil environment. The exudation of oxalic acid by maize roots increases in the presence of cadmium, which forms complexes with Cd²⁺ ions, reducing their toxicity. Flavonoid release can be triggered in response to heavy metal stress, aiding in the sequestration of metals and mitigating oxidative damage to plant cells.

D. Effect of Plant-Microbe-Pathogen Interactions Under Changing Climate Conditions

Climate change, characterized by rising temperatures, altered precipitation patterns, and increased atmospheric CO₂, has significant implications for root exudation and subsequent plant-microbe interactions [36]. Elevated CO₂ levels can enhance root biomass and increase the quantity of exudates released, thereby promoting the growth of beneficial soil microbes like mycorrhizae and nitrogen-fixing bacteria. Enhanced microbial activity under elevated CO₂ can lead to improved nutrient cycling, but it may also exacerbate the competition between beneficial and pathogenic microbes in the rhizosphere. Drought conditions induced by climate change can alter root exudate composition, leading to the enhanced release of osmoprotective substances that support drought-resistant microbial communities. The increased abundance of certain exudates can also attract soil-borne pathogens, such as *Fusarium* and *Pythium*, which can exploit weakened plants during prolonged droughts [37].

VI. Applications of Root Exudate Research in Agriculture

Root exudates are increasingly recognized for their potential to revolutionize sustainable agricultural practices by enhancing crop productivity, improving soil health, and reducing dependency on chemical inputs. The intricate interactions mediated by root exudates in the rhizosphere can be leveraged to develop strategies for optimizing nutrient availability, controlling pathogens, and fostering beneficial plant-microbe relationships.

A. Enhancing Crop Productivity Through Root Exudate Management

Harnessing the power of root exudates can significantly boost crop productivity. By understanding and manipulating exudate profiles, researchers and farmers can develop innovative techniques to enhance nutrient uptake, stimulate beneficial microbes, and optimize soil health [38].

1. Development of Biofertilizers and Biostimulants

One promising application of root exudate research is the development of biofertilizers and biostimulants that can improve plant growth and productivity in a sustainable manner. Biofertilizers are formulations containing beneficial microbes like Plant Growth-Promoting Rhizobacteria (PGPR)

and mycorrhizal fungi that utilize root exudates for colonization and activity. These biofertilizers enhance nutrient availability, particularly nitrogen and phosphorus, which are often limiting in agricultural soils. Root exudates rich in sugars, organic acids, and amino acids stimulate microbial populations that are essential for nutrient cycling. For example, PGPR such as *Bacillus*, *Azospirillum*, and *Pseudomonas* spp. are known to increase nutrient availability by solubilizing phosphorus and fixing atmospheric nitrogen, thereby reducing the need for chemical fertilizers [39]. The exudation of specific compounds like malic acid has been shown to attract beneficial microbes, which can improve root growth and nutrient uptake. Biostimulants, on the other hand, include compounds or formulations that directly enhance plant physiological processes, improving stress tolerance and yield. These may include seaweed extracts, humic substances, and microbial inoculants that interact with root exudates to boost plant growth. The application of biostimulants can enhance root exudation, which in turn promotes beneficial microbial colonization, leading to better nutrient absorption and increased resilience to drought and salinity stress.

2. Use of Cover Crops and Crop Rotation to Optimize Root Exudation

Another practical approach to enhancing crop productivity involves using cover crops and crop rotation to modulate root exudation patterns. Cover crops such as legumes and clover release nitrogen-rich exudates that enhance soil fertility by stimulating nitrogen-fixing bacteria [40]. These crops also exude organic acids that mobilize phosphorus, making it more accessible to subsequent crops. Crop rotation, especially with deep-rooted species, promotes the secretion of diverse exudates that enrich the soil microbiome, reducing pathogen loads and increasing nutrient cycling. For example, rotating cereal crops with legumes has been shown to reduce soil-borne diseases and improve soil structure due to increased microbial diversity and activity in the rhizosphere.

B. Biological Control of Soil-Borne Pathogens Using Exudate-Mediated Microbial Recruitment

Root exudates play an important role in attracting beneficial microbes that can suppress soil-borne pathogens. By leveraging this natural defense mechanism, agricultural practices can be developed to control diseases more sustainably [41].

1. Formulation of Biocontrol Agents (PGPR, Mycorrhizal Fungi)

Biocontrol agents such as PGPR and mycorrhizal fungi are increasingly used to manage soil-borne diseases by exploiting root exudate-mediated interactions. These beneficial microbes can outcompete pathogens by colonizing root surfaces, producing antimicrobial compounds, and inducing plant systemic resistance. *Pseudomonas fluorescens* responds to exudates like malic acid, leading to the production of lipopeptides that inhibit pathogenic fungi such as *Fusarium* and *Pythium*. Arbuscular mycorrhizal fungi (AMF) are also effective in protecting plants against soil pathogens. The release of strigolactones by plant roots not only promotes AMF colonization but also enhances plant resistance to diseases by improving nutrient uptake and activating plant immune responses [42]. By formulating bioinoculants that are compatible with specific crop exudate profiles, it is possible to maximize the effectiveness of biocontrol strategies.

2. Breeding Crops with Desirable Root Exudate Profiles for Disease Resistance

Recent advances in plant breeding and biotechnology have opened new possibilities for selecting or genetically engineering crops with optimized root exudate profiles to enhance resistance to pathogens. Crops can be bred to increase the secretion of specific antimicrobial exudates or compounds that recruit beneficial microbes, thereby reducing the incidence of soil-borne diseases. For example, maize varieties that exude higher levels of benzoxazinoids are more resistant to root rot diseases caused by *Fusarium* [43]. Breeding tomato plants for enhanced exudation of phenolics has been shown to suppress the growth of *Ralstonia solanacearum*, a bacterial pathogen that causes wilt. These strategies

can lead to the development of crop varieties that are more resilient to biotic stresses, reducing the need for chemical pesticides.

C. Potential for Sustainable Agriculture and Soil Health Improvement

The integration of root exudate research into agricultural practices holds great promise for advancing sustainable agriculture. By optimizing root exudation, it is possible to improve soil health, reduce dependence on synthetic inputs, and enhance crop resilience to environmental stressors. One of the most promising applications is the use of root exudate-mediated processes to enhance soil organic matter and nutrient cycling. As plants exude a diverse array of compounds, they enrich the soil microbiome, which in turn improves soil structure, enhances nutrient availability, and increases soil carbon sequestration [44]. Cover crops that exude high levels of polysaccharides and organic acids can improve soil aggregation and water retention, which is critical for maintaining soil fertility in dry regions. The use of bioinoculants that interact with root exudates to promote nutrient cycling can reduce the need for chemical fertilizers, leading to a decrease in greenhouse gas emissions and soil pollution. For example, inoculating legumes with nitrogen-fixing bacteria can reduce the reliance on synthetic nitrogen fertilizers, which are a major source of environmental pollution. In climate change, promoting root exudation to support beneficial microbial communities can enhance crop resilience to drought, salinity, and heat stress, thereby contributing to food security [45]. By leveraging the natural interactions between plants and soil microbes, agricultural systems can become more resilient, sustainable, and productive in the face of global environmental challenges.

VII. Challenges and Future

While research on root exudates has provided substantial insights into their role in modulating soil ecosystems, numerous challenges and knowledge gaps remain. These limitations hinder the full application of root exudate studies in agriculture and environmental management.

A. Technical Challenges in Studying Root Exudates

Researching root exudates is fraught with technical complexities, especially when attempting to isolate, characterize, and analyze these compounds in natural soil systems.

1. Difficulties in Isolating and Characterizing Root Exudates in Soil Systems

Root exudates are continuously released into the rhizosphere, a highly dynamic zone influenced by numerous biotic and abiotic factors. Isolating root exudates from soil is challenging due to the rapid degradation and transformation of these compounds by soil microbes [46]. Traditional methods, such as hydroponic and sand culture systems, often do not accurately represent the conditions in natural soils, leading to potential discrepancies in exudate composition. Distinguishing between compounds that are truly secreted by roots versus those that are derived from microbial metabolism remains a significant obstacle. Soil organic matter and microbial secretions can contaminate exudate samples, making it difficult to isolate pure root-derived compounds. The complexity of soil matrices also makes it challenging to study root exudates in situ, where interactions with microbes and soil particles can alter the original composition of exudates [47].

2. Need for Advanced Analytical Techniques and Bioinformatics Tools

The identification and quantification of root exudates require advanced analytical techniques due to their chemical diversity and low concentrations in soil environments. Techniques such as gas chromatography-mass spectrometry (GC-MS), liquid chromatography-mass spectrometry (LC-MS), and nuclear magnetic resonance (NMR) spectroscopy are commonly used but have limitations related to sensitivity, resolution, and the ability to analyze complex mixtures. The use of bioinformatics and metabolomics is becoming increasingly essential to process large datasets generated from these analytical methods. The lack of standardized protocols for sample collection, processing, and data

analysis complicates the reproducibility of results across different studies. There is a growing need for the development of integrated databases and bioinformatics tools that can facilitate the identification of novel exudates and their associated microbial interactions [48].

B. Knowledge Gaps in Root Exudate-Microbe-Pathogen Interactions

Despite significant advances in understanding the role of root exudates in plant-microbe interactions, several knowledge gaps limit our ability to harness these interactions effectively for agricultural applications.

1. Lack of the Temporal Dynamics of Exudation

Root exudation is not a static process; it varies with the plant's developmental stage, environmental conditions, and microbial interactions. Much of the current research has focused on static snapshots of exudation profiles, often ignoring the temporal dynamics that influence these processes. The secretion of organic acids and phenolics may fluctuate in response to diurnal cycles, nutrient availability, and biotic stress [49].

2. Interactions Between Multiple Plant Species and Soil Microbiomes

Most studies on root exudates have been conducted on single plant species grown under controlled conditions, which does not reflect the complexity of natural ecosystems where multiple plant species coexist. In diverse cropping systems or natural plant communities, interactions between the root exudates of different species can influence soil microbiome composition and function. For example, intercropping systems can alter root exudation patterns, leading to synergistic or antagonistic effects on microbial communities that impact nutrient cycling and disease suppression [50].

C. Potential Future Research Areas

Addressing the current challenges and knowledge gaps in root exudate research will require interdisciplinary approaches and innovative strategies. The following are potential areas where future research could make significant contributions.

1. Engineering Plants for Optimized Root Exudate Production

Biotechnological advances offer the potential to genetically engineer plants with optimized root exudate profiles to enhance nutrient uptake, attract beneficial microbes, and suppress pathogens. For example, increasing the secretion of specific organic acids could improve phosphorus availability in soils with low phosphorus levels. Engineering crops to produce higher levels of antimicrobial compounds, like benzoxazinoids, could enhance resistance to soil-borne pathogens [51]. CRISPR-Cas9 and other genome editing technologies can be employed to modify the expression of genes involved in root exudate biosynthesis, allowing researchers to develop crops that are more resilient to environmental stressors and diseases.

2. Root Exudates in the Context of Climate Change Adaptation

Climate change is expected to alter plant physiology, soil properties, and microbial dynamics, which could have profound effects on root exudation patterns [52]. Elevated CO₂ levels can increase root biomass and alter exudate composition, potentially enhancing soil carbon sequestration and nutrient cycling. It remains unclear how these changes will impact plant-microbe-pathogen interactions under different climate scenarios, such as increased drought or extreme temperatures. Future research should focus on how plants can be managed or bred to optimize root exudation in changing environments to improve agricultural sustainability.

3. Integrated Pest Management (IPM) Strategies Leveraging Root Exudates

Root exudates can play an important role in integrated pest management (IPM) by enhancing natural disease suppression. Leveraging root exudates to recruit beneficial microbes that outcompete pathogens could reduce the need for chemical pesticides [53]. For example, crop varieties that release higher amounts of malic acid or phenolic compounds can attract PGPR that induce systemic resistance, thereby reducing disease incidence. Using bioinoculants that interact synergistically with crop exudates can create a healthier rhizosphere environment that supports plant growth while minimizing pest pressures. Developing IPM strategies that integrate the manipulation of root exudates with other sustainable practices, such as crop rotation and organic amendments, could offer a holistic approach to disease management, improving agricultural productivity while reducing environmental impacts.

VIII. Conclusion

Root exudates play an important role in shaping plant-microbe-pathogen interactions, offering significant potential for advancing sustainable agricultural practices. These exudates not only enhance nutrient acquisition and promote beneficial microbial associations but also serve as natural defenses against soil-borne pathogens. The complexities of root exudate dynamics is challenged by technical limitations, knowledge gaps, and environmental variability. Future research focusing on optimizing exudate profiles, engineering stress-resilient crops, and leveraging root exudates in integrated pest management could transform crop productivity and soil health. As agriculture faces pressures from climate change and declining soil fertility, harnessing the power of root exudates offers a promising path toward more resilient and eco-friendly farming systems, reducing the need for chemical inputs and contributing to global food security.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

1. Alamgir, A. N. M., & Alamgir, A. N. M. (2018). Secondary metabolites: Secondary metabolic products consisting of C and H; C, H, and O; N, S, and P elements; and O/N heterocycles. *Therapeutic use of medicinal plants and their extracts: volume 2: phytochemistry and bioactive compounds*, 165-309.
2. Prashar, P., Kapoor, N., & Sachdeva, S. (2014). Rhizosphere: its structure, bacterial diversity and significance. *Reviews in Environmental Science and Bio/Technology*, 13, 63-77.
3. Prashar, P., Kapoor, N., & Sachdeva, S. (2014). Rhizosphere: its structure, bacterial diversity and significance. *Reviews in Environmental Science and Bio/Technology*, 13, 63-77.
4. Sharma, P., Sharma, M. M. M., Malik, A., Vashisth, M., Singh, D., Kumar, R., ... & Pandey, V. (2021). Rhizosphere, rhizosphere biology, and rhizospheric engineering. In *Plant growth-promoting microbes for sustainable biotic and abiotic stress management* (pp. 577-624). Cham: Springer International Publishing.
5. Ma, W., Tang, S., Dengzeng, Z., Zhang, D., Zhang, T., & Ma, X. (2022). Root exudates contribute to belowground ecosystem hotspots: A review. *Frontiers in Microbiology*, 13, 937940.
6. Singh, D. P., Singh, H. B., & Prabha, R. (Eds.). (2016). *Microbial inoculants in sustainable agricultural productivity* (Vol. 2, p. 308). New Delhi: Springer.
7. Vranova, V., Rejsek, K., Skene, K. R., Janous, D., & Formanek, P. (2013). Methods of collection of plant root exudates in relation to plant metabolism and purpose: a review. *Journal of Plant Nutrition and Soil Science*, 176(2), 175-199.

8. Ma, W., Tang, S., Dengzeng, Z., Zhang, D., Zhang, T., & Ma, X. (2022). Root exudates contribute to belowground ecosystem hotspots: A review. *Frontiers in Microbiology*, *13*, 937940.
9. Gunina, A., & Kuzyakov, Y. (2015). Sugars in soil and sweets for microorganisms: review of origin, content, composition and fate. *Soil Biology and Biochemistry*, *90*, 87-100.
10. Di Marsico, A., Scranò, L., Labella, R., Lanzotti, V., Rossi, R., Cox, L., ... & Amato, M. (2018). Mucilage from fruits/seeds of chia (*Salvia hispanica* L.) improves soil aggregate stability. *Plant and Soil*, *425*, 57-69.
11. Alamgir, A. N. M., & Alamgir, A. N. M. (2018). Secondary metabolites: Secondary metabolic products consisting of C and H; C, H, and O; N, S, and P elements; and O/N heterocycles. *Therapeutic use of medicinal plants and their extracts: volume 2: phytochemistry and bioactive compounds*, 165-309.
12. Dong, W., & Song, Y. (2020). The significance of flavonoids in the process of biological nitrogen fixation. *International journal of molecular sciences*, *21*(16), 5926.
13. Huang, X. F., Chaparro, J. M., Reardon, K. F., Zhang, R., Shen, Q., & Vivanco, J. M. (2014). Rhizosphere interactions: root exudates, microbes, and microbial communities. *Botany*, *92*(4), 267-275.
14. Adeniji, A., Huang, J., Li, S., Lu, X., & Guo, R. (2024). Hot viewpoint on how soil texture, soil nutrient availability, and root exudates interact to shape microbial dynamics and plant health. *Plant and Soil*, 1-22.
15. Meuser, H., & Meuser, H. (2013). Soil Decontamination. *Soil Remediation and Rehabilitation: Treatment of Contaminated and Disturbed Land*, 201-278.
16. Mahrous, E. A., & Farag, M. A. (2015). Two dimensional NMR spectroscopic approaches for exploring plant metabolome: A review. *Journal of advanced research*, *6*(1), 3-15.
17. Liu, A., Ku, Y. S., Contador, C. A., & Lam, H. M. (2020). The impacts of domestication and agricultural practices on legume nutrient acquisition through symbiosis with rhizobia and arbuscular mycorrhizal fungi. *Frontiers in Genetics*, *11*, 583954.
18. Abd-Alla, M. H., Al-Amri, S. M., & El-Enany, A. W. E. (2023). Enhancing Rhizobium-Legume Symbiosis and Reducing Nitrogen Fertilizer Use Are Potential Options for Mitigating Climate Change. *Agriculture*, *13*(11), 2092.
19. Etesami, H., Jeong, B. R., & Glick, B. R. (2021). Contribution of arbuscular mycorrhizal fungi, phosphate-solubilizing bacteria, and silicon to P uptake by plant. *Frontiers in Plant Science*, *12*, 699618.
20. Yu, K., Stringlis, I. A., van Bentum, S., de Jonge, R., Snoek, B. L., Pieterse, C. M., ... & Berendsen, R. L. (2021). Transcriptome signatures in *Pseudomonas simiae* WCS417 shed light on role of root-secreted coumarins in Arabidopsis-mutualist communication. *Microorganisms*, *9*(3), 575.
21. Dashora, K. (2011). Nitrogen yielding plants: the pioneers of agriculture with a multipurpose. *American-Eurasian Journal of Agronomy*, *4*(2), 34-37.
22. Zainab, N., Amna, Khan, A. A., Azeem, M. A., Ali, B., Wang, T., ... & Chaudhary, H. J. (2021). PGPR-mediated plant growth attributes and metal extraction ability of *Sesbania sesban* L. in industrially contaminated soils. *Agronomy*, *11*(9), 1820.
23. Choudhary, D. K., Prakash, A., & Johri, B. N. (2007). Induced systemic resistance (ISR) in plants: mechanism of action. *Indian Journal of Microbiology*, *47*, 289-297.
24. Olanrewaju, O. S., Ayangbenro, A. S., Glick, B. R., & Babalola, O. O. (2019). Plant health: feedback effect of root exudates-rhizobiome interactions. *Applied microbiology and biotechnology*, *103*, 1155-1166.
25. Ramsdale, M. (2008). Programmed cell death in pathogenic fungi. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research*, *1783*(7), 1369-1380.

26. Mansfield, J. W. (2000). Antimicrobial compounds and resistance: the role of phytoalexins and phytoanticipins. In *Mechanisms of resistance to plant diseases* (pp. 325-370). Dordrecht: Springer Netherlands.
27. Choudhary, D. K., Prakash, A., & Johri, B. N. (2007). Induced systemic resistance (ISR) in plants: mechanism of action. *Indian Journal of Microbiology*, *47*, 289-297.
28. Mhlongo, M. I., Piater, L. A., Madala, N. E., Labuschagne, N., & Dubery, I. A. (2018). The chemistry of plant–microbe interactions in the rhizosphere and the potential for metabolomics to reveal signaling related to defense priming and induced systemic resistance. *Frontiers in plant science*, *9*, 112.
29. Cen, Z., Hu, B., Yang, S., Ma, G., Zheng, Y., & Dong, Y. (2024). Mechanism of benzoxazinoids affecting the growth and development of *Fusarium oxysporum* f. sp. *fabae*. *Plant Molecular Biology*, *114*(3), 1-17.
30. Doornbos, R. F., van Loon, L. C., & Bakker, P. A. (2012). Impact of root exudates and plant defense signaling on bacterial communities in the rhizosphere. A review. *Agronomy for Sustainable Development*, *32*, 227-243.
31. More, S. S., Shinde, S. E., & Kasture, M. C. (2020). Root exudates a key factor for soil and plant: An overview. *Pharma Innov. J*, *8*, 449-459.
32. Tan, Z., Liu, F., Wan, Y., Zhu, S., Zhang, J., Zhang, K., & Luo, L. (2023). Morphological and Physiological Mechanism of Activating Insoluble Inorganic Phosphorus of Different Peanut (*Arachis hypogaea* L.) Varieties under Low Phosphorus. *Agriculture*, *13*(12), 2270.
33. Tsotetsi, T., Nephali, L., Malebe, M., & Tugizimana, F. (2022). Bacillus for plant growth promotion and stress resilience: what have we learned?. *Plants*, *11*(19), 2482.
34. George, E., Horst, W. J., & Neumann, E. (2012). Adaptation of plants to adverse chemical soil conditions. In *Marschner's mineral nutrition of higher plants* (pp. 409-472). Academic press.
35. Guy, C., Kaplan, F., Kopka, J., Selbig, J., & Hinch, D. K. (2008). Metabolomics of temperature stress. *Physiologia plantarum*, *132*(2), 220-235.
36. Aamir, M., Rai, K. K., Dubey, M. K., Zehra, A., Tripathi, Y. N., Divyanshu, K., ... & Upadhyay, R. S. (2019). Impact of climate change on soil carbon exchange, ecosystem dynamics, and plant–microbe interactions. In *Climate change and agricultural ecosystems* (pp. 379-413). Woodhead Publishing.
37. Naik, A., Jogi, M., & Shreenivas, B. V. (2024). Assessing the impact of climate change on global crop yields and farming practices. *Archives of Current Research International*, *24*(5), 696-712.
38. Qiu, Z., Egidi, E., Liu, H., Kaur, S., & Singh, B. K. (2019). New frontiers in agriculture productivity: optimised microbial inoculants and in situ microbiome engineering. *Biotechnology advances*, *37*(6), 107371.
39. Hayat, R., Ahmed, I., & Sheirdil, R. A. (2012). An overview of plant growth promoting rhizobacteria (PGPR) for sustainable agriculture. *Crop production for agricultural improvement*, 557-579.
40. Kebede, E. (2021). Contribution, utilization, and improvement of legumes-driven biological nitrogen fixation in agricultural systems. *Frontiers in Sustainable Food Systems*, *5*, 767998.
41. Akanmu, A. O., Babalola, O. O., Venturi, V., Ayilara, M. S., Adeleke, B. S., Amoo, A. E., ... & Glick, B. R. (2021). Plant disease management: leveraging on the plant-microbe-soil interface in the biorational use of organic amendments. *Frontiers in Plant Science*, *12*, 700507.
42. Boyno, G., Rezaee Danesh, Y., Demir, S., Teniz, N., Mulet, J. M., & Porcel, R. (2023). The Complex Interplay between Arbuscular Mycorrhizal Fungi and Strigolactone: Mechanisms, Sinergies, Applications and Future Directions. *International Journal of Molecular Sciences*, *24*(23), 16774.

43. Duan, S., Jin, J., Gao, Y., Jin, C., Mu, J., Zhen, W., ... & Ma, J. (2022). Integrated transcriptome and metabolite profiling highlights the role of benzoxazinoids in wheat resistance against Fusarium crown rot. *The Crop Journal*, 10(2), 407-417.
44. Chaparro, J. M., Sheflin, A. M., Manter, D. K., & Vivanco, J. M. (2012). Manipulating the soil microbiome to increase soil health and plant fertility. *Biology and Fertility of Soils*, 48, 489-499.
45. Zhao, J., Yu, X., Zhang, C., Hou, L., Wu, N., Zhang, W., ... & Tian, J. (2023). Harnessing microbial interactions with rice: Strategies for abiotic stress alleviation in the face of environmental challenges and climate change. *Science of The Total Environment*, 168847.
46. Casas, M. E., & Matamoros, V. (2021). Analytical challenges and solutions for performing metabolomic analysis of root exudates. *Trends in Environmental Analytical Chemistry*, 31, e00130.
47. Oburger, E., & Jones, D. L. (2018). Sampling root exudates—mission impossible?. *Rhizosphere*, 6, 116-133.
48. Mishra, S., Lin, Z., Pang, S., Zhang, W., Bhatt, P., & Chen, S. (2021). Recent advanced technologies for the characterization of xenobiotic-degrading microorganisms and microbial communities. *Frontiers in Bioengineering and Biotechnology*, 9, 632059.
49. Iqbal, Z., Iqbal, M. S., Hashem, A., Abd_Allah, E. F., & Ansari, M. I. (2021). Plant defense responses to biotic stress and its interplay with fluctuating dark/light conditions. *Frontiers in Plant Science*, 12, 631810.
50. Li, C., Lambers, H., Jing, J., Zhang, C., Bezemer, T. M., Klironomos, J., ... & Zhang, F. (2024). Belowground cascading biotic interactions trigger crop diversity benefits. *Trends in Plant Science*.
51. Niu, B., Wang, W., Yuan, Z., Sederoff, R. R., Sederoff, H., Chiang, V. L., & Borriss, R. (2020). Microbial interactions within multiple-strain biological control agents impact soil-borne plant disease. *Frontiers in Microbiology*, 11, 585404.
52. Lamichhane, J. R., Barbetti, M. J., Chilvers, M. I., Pandey, A. K., & Steinberg, C. (2024). Exploiting root exudates to manage soil-borne disease complexes in a changing climate. *Trends in Microbiology*, 32(1), 27-37.
53. Ali, S., & Glick, B. R. (2024). Root Exudate Metabolites Alter Food Crops Microbiomes, Impacting Plant Biocontrol and Growth. *Crops*, 4(1), 43-54.