

ENHANCED ITERATIVE METHODS USING MAMADU-NJOSEH POLYNOMIALS FOR SOLVING THE HESTON STOCHASTIC PARTIAL DIFFERENTIAL EQUATION

Abstract

In this paper, a modified form of the Homotopy Perturbation Method (HPM) and the Variational Iteration Method (VIM), both developed by J.H. He, is presented using the newly constructed Mamadu-Njoseh orthogonal polynomial(MNPs)as modifier and basis function. The HPM combines principles from the field of topology and the usual perturbation techniques, while the goal of the VIM is to construct a correction functional for nonlinear systems. These two analytical techniques are modified through the orthogonal collocation method using the MNPs as basis function. The modified methods are employed to determine which approximates the Heston Stochastic Partial Differential Equation (HSPDE) faster to its exact solution. The Heston SPDE is a volatility model for determining the European bond and currency options as determined by stock pricing. While other methods exist in literature in determining the numerical solution to the HSPDE, our numerical schemes, the MHPM and the MVIM being presented as a new technique by the presence of the MNPs is noticed by comparison, to possess a faster approximation to the exact solution. In this work, we observe that the Modified VIM approximates faster to the exact solution of the HSPDE than the modified HPM due to its highly effective use of orthogonal polynomials, greater chances of handling of nonlinearities, superiority in the convergence properties, and having to adapt better to the boundary and initial conditions of the problem. This new approach is highly beneficial in handling large and complex nonlinear PDEs due to the presence of the MNPs and the iterative nature of the VIM.

Keyword: Homotopy Perturbation Method (HPM), Variational Iteration Method (VIM), orthogonal polynomials, Mamadu-Njoseh Polynomials (MNPs), Heston Stochastic Partial Differential Equation (HSPDE).

AMS CLASSIFICATION: 65F10

1. Introduction

Based on the limitations of most perturbation methods which served as an- analytical techniques for solving nonlinear problems, Ji-Huan He in 1998 developed the Homotopy perturbation method (HPM), a special technique which combines principles from the field of topology and the usual perturbation techniques. It is efficient in obtaining approximate solutions to Nonlinear DEs, BVPs, PDEs, Biology and Epidermiology, Astrophysics and Cosmology, Stability Analysis, Control theory, etc where other methods like direct integration, perturbation theory, or separation of variables may not be function properly. Gupta [1] used the HPM to derive an exact solution for the coupled one-dimensional time fractional nonlinear shallow water system which is a system of PDEs involved with the flow of fluids in riverine areas. [2] employed the HPM to find the solution of some linear and non- linear PDEs, while [3] employed the HE's HPM in their article to derive a solution for a nonlinear system of two-dimensional Volterra-Fredholm integral equations, while [4],[5], and [6] employed various modified HPM in finding approximate solutions to a wide range of problems. A year later, [7] developed an iterative method for solving differential and integral equations. This method, the Variational Iteration Method (VIM) was devoid of the limitations of the Grid point techniques, the Spline solution, the Perturbation method and the Adomian method [8]. The method has the ability to treat linear and non-linear equations alike without any unrealistic assumptions. It is able to produce approximations to solutions that converge rapidly due to the coupling of the Lagrange Multiplier Method and iteration schemes, and sometimes obtains an exact solution in a finite number of iterations. [9] applied the VIM to derive the analytic solution of the space fractional diffusion equation, while [10] employed the method on the non-linear free vibration of conservative oscillator. A modified form of the VIM was applied by [11] to solve a fourth order parabolic partial differential equation with variable coefficients and [12] in same vain applied a modified VIM to obtain the solution of some non-linear, nonhomogeneous differential equations. In [13], a comparison between the VIM, the ADM and HPM was carried out on the numerical solutions of the Heston partial differential equation where the authors concluded that the VIM is much easier, more convenient, more stable and efficient than the other two iterative methods. In [14],a modified VIM was employed to find the approximate solution for the Time Fractional Nagumo Equation, while [15] and [16] carried out comparisons on the solutions of the Heston stochastic Partial Differential Equation between a MVIM and other numerical methods. Other works on the VIM can be

obtained in [17], [18], [19] and [20]. Mamadu-Njoseh Polynomials (MNPs) was developed and employed on the numerical solutions of fifth order boundary value problems by [21]. The MNPs, a $C^{[a,b]}$ orthogonal polynomial was also used to derive the numerical solutions of the Volterra equation using Galerkin method [22]. In [23], the authors worked on the space Discretization of Time-Fractional Telegraph equation using the MNPs as basis factor, while [24] combined the Gauss Quadrature with the MNPs to obtain the Gauss-Mamadu-Njoseh Quadrature and used it to obtain solutions for Numerical Integration Interpolation. This paper is geared towards using the MNPs as a modifier and basis function to modify the HPM and the VIM in arriving at the approximate solution of Heston stochastic partial differential equation via the orthogonal collocation method and comparisons made to determine which approximates the HSPDE faster to its exact solution. The swift convergence of the MVIM is represented graphically as against the MHPM.

2. Heston Stochastic Partial Differential Equation (HSPDE)

According to [25], the Heston model is a typical stochastic volatility model of the form $\alpha(t, S(t), V(t)) = (a - bV(t))$ and $\beta(t, S(t), V(t)) = \sigma\sqrt{V(t)}$, while [13] gives the Heston model as

$$\begin{aligned}\frac{dS(t)}{S(t)} &= rdt + \sigma\sqrt{V}d\widehat{W}_1t \\ \frac{dV(t)}{d(t)} &= (a - bV(t)) + \sigma\sqrt{V}d\widehat{W}_2t\end{aligned}\tag{1}$$

where α is the option price, β is the price of the volatility risk, r is the interest rate, $S(t)$ is the asset price at time t , $V(t)$ is the volatility of the asset price at time t with \sqrt{V} as the variance of the volatility, a is the long-run mean, b is the speed of the mean reversion, σ is the volatility of the of the variance process, while $d\widehat{W}_1(t)$ and $d\widehat{W}_2(t)$ are correlated Brownian motions under the risk-neutral measure with the correlation coefficient $\rho \in (-1, 1)$ such that

$$\widehat{W}_1(t)d\widehat{W}_2(t)\tag{2}$$

The risk-neutral price of a call expiring at time $t \leq T$ in the Heston stochastic volatility model is given as

$$c(t, S(t), V(t)) = \widehat{E} \left[e^{-r(T-t)} (S(T) - K)^+ \right], 0 \leq t \leq T\tag{3}$$

The equation

$$\frac{\partial c}{\partial t} + rs\frac{\partial c}{\partial s} + (a - bv)\frac{\partial c}{\partial v} + \frac{1}{2}s^2v\frac{\partial^2 c}{\partial s^2} + \rho\sigma sv\frac{\partial^2 c}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c}{\partial v^2} - rc = 0\tag{4}$$

is the Heston partial differential equation (PDE) for the fair values of European style options forming a time dependant convection diffusion reaction equation with mixed spatial derivative terms. The Heston PDE (2.4) has the initial and boundary conditions given as

$$\begin{aligned}C(S, v, t) &= \max(0, s - K) \\ C(0, v, t) &= 0\end{aligned}\tag{5}$$

where $k > 0$ is the given strike price.

3. He's Homotopy Perturbation Method (HPM)

According to [26], the applications of the homotopy perturbation method mainly cover in nonlinear differential equations, nonlinear integral equations, nonlinear differential-integral equations, difference differential equations, and fractional differential equations. He [12] noted that the earlier perturbation methods were limited based on what was called the "small parameter assumption" where it was assumed that a small parameter whose appropriate choice that leads to ideal results must exist in an equation. Should these so called "small parameters" be chosen without care and suitability, the given result will become inappropriate. With the existence of this assumption, it became difficult to explore greatly many nonlinear problems as most of them have big parameters. The work of the HPM was to eliminate this limitation existing in the traditional perturbation methods.

By the homotopy technique, a homotopy $V(r, \rho) : \infty \rightarrow [0, 1] \rightarrow \mathfrak{R}$ is constructed satisfying

$$x(V, \rho) = (1 - \rho)[L(V) - L(U_0)] + \rho[(V) - f(r)] = 0, \quad \rho \in [0, 1], \quad r \in \infty\tag{6}$$

or

$$x(V, \rho) = L(V) - L(U_0) + \rho L(U_0) + \rho[N(V) - f(r)] = 0\tag{7}$$

4. He's variational Iteration Method (VIM)

Ji Huan He in 1999 [7] developed a Variational Iteration Method (VIM) in solving linear and nonlinear equations alike without any unrealistic assumptions. [27] concluded that the method converges faster to the exact solution by successive approximation. Unlike the Adomian Decomposition Method (ADM) developed by George Adomian in 1982, the VIM does not require any form of polynomials to obtain its approximate solution. According to [8], the basic concept of the VIM is to construct a correction functional for nonlinear systems. This is given as

$$U_{n+1}(x, t) = U_n(x, t) + \int_0^t \lambda(LU_n(\tau) + N\widehat{U}_n(\tau)g(\tau))d\tau \quad (8)$$

where λ is a general Langrange multiplier, which can be identified optimally via the variational theory. $NU\widehat{n}(\tau)$ is considered as the restricted variations. If we set the Langrange multiplier $\lambda = 1$, then (8) will be given as

$$U_{n+1}(x, t) = U_n(x, t)(LU_n(\tau) + \int_0^t NU_n(\tau) - g(\tau))d\tau \quad (9)$$

$NU_n(\tau)$ is called the correction term and (9) can be solved iteratively using $U_0(x)$ as the initial approximation with possible unknowns.

5. Mamadu-Njoseh Polynomials (MNP)

The MNPs are a set of orthogonal polynomials having an interval of $[1, 1]$ and a weight function of $w(x) = (1 + x^2)$. It is given as

$$\int_{-1}^1 \varphi_m(x)\varphi_n(x)(1 + x^2)dx = 0 \quad (10)$$

with the first four MNPs given as

$$\begin{aligned} \varphi_0(x) &= 1 \\ \varphi_1(x) &= 0 \\ \varphi_2(x) &= \frac{1}{3}(5x^2 - 2) \\ \varphi_3(x) &= \frac{1}{5}(14x^3 - 9x) \end{aligned} \quad (11)$$

6. Modified HPM for Heston SPDE

The scheme for generating the initial approximation through the OCM with the Mamadu-Njoseh polynomials as basis function is described as follows:

Let the initial approximation be given as

$$C_O = \sum_{i=1}^N a_i \varphi_i(x) \quad (12)$$

where a_i are unknown constants to be determined and $\varphi_i(x)$ are the Mamadu-Njoseh Polynomials with interval of orthogonality $[-1, 1]$. According to [8], the Heston SPDE has a generalized initial condition

$$C(O) = 2s^2t^2 \quad (13)$$

Incorporating (12) and (13), we have

$$C(O) = \sum_{i=0}^N a_i \varphi_i(x) = 2s^2t^2 \quad (14)$$

Solving (14) at $N = 3$ (chosen arbitrarily) and substituting the $\varphi_i(x), i = 0, 1, 2, 3$, we have

$$a_0 + a_1s + a_2 \left(\frac{5}{3}s^2 - \frac{2}{3} \right) + a_3 \left(\frac{14}{5}s^3 - \frac{9}{5}s \right) = 2s^2t^2 \quad (15)$$

In [13], the value of t is defined within the range $0 \leq t \leq T$. Thus, collocating (6.4) at the zeroes of $\varphi_4(x)$, that is,

$$s = 0.3676425560, -0.3676425560, 0.8756710201, -0.8756710201$$

and writing the resulting linear algebraic equations in the form

$$\underline{AX} = \underline{b} \quad (16)$$

where,

$$A = \begin{bmatrix} 1 & 0.3676425560 & -0.4413982517 & -0.5226219309 \\ 1 & -0.3676425560 & -0.4413982517 & 0.5226219309 \\ 1 & 0.8756710201 & 0.6113328923 & 0.303892222 \\ 1 & -0.8756710201 & 0.6113328923 & -0.303892222 \end{bmatrix}$$

$$\underline{X} = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix}$$

$$\underline{b} = \begin{bmatrix} 52.98313121 \\ 52.98313121 \\ 300.5854963 \\ 300.5854963 \end{bmatrix}$$

Solving (16) using Gaussian elimination method, we obtain the following values for the constant a'_i 's

$$\begin{aligned} a_0 &= 156.800000 \\ a_1 &= 0.000000 \\ a_2 &= 235.200000 \\ a_3 &= 0.000000 \end{aligned}$$

Thus, substituting the above in (14), we obtain

$$c_0 = 392.00000s^2 \quad (17)$$

We use the initial approximation $c_0(s, v, 0) = 392.00000s^2$ as given in (17) satisfying the initial condition for the Heston SPDE. We introduce the structure of the HPM as relating to the HSPDE given as

$$H(s, v, t) = (1-p)(L(c) - L(v_0)) + p\left(\frac{\partial c}{\partial t} - rs\frac{\partial c}{\partial s} + (a-bv)\frac{\partial c}{\partial v} + \frac{1}{2}s^2v\frac{\partial^2 c}{\partial s^2} + \rho\sigma sv\frac{\partial^2 c}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c}{\partial v^2} - rc\right) = 0 \quad (18)$$

with given algorithm

$$\begin{aligned} P^0 &= \frac{\partial c_0}{\partial t} - \frac{\partial v_0}{\partial t} = 0 \\ P^1 &= \frac{\partial c_1}{\partial t} - rs\frac{\partial c_0}{\partial s} - (a-bv)\frac{\partial c_0}{\partial v} - \frac{1}{2}s^2v\frac{\partial^2 c_0}{\partial s^2} - \rho\sigma sv\frac{\partial^2 c_0}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c_0}{\partial v^2} + rc_0 + \frac{\partial v_0}{\partial t} = 0 \\ P^2 &= \frac{\partial c_2}{\partial t} - rs\frac{\partial c_1}{\partial s} - (a-bv)\frac{\partial c_1}{\partial v} - \frac{1}{2}s^2v\frac{\partial^2 c_1}{\partial s^2} - \rho\sigma sv\frac{\partial^2 c_1}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c_1}{\partial v^2} + rc_1 = 0 \\ P^3 &= \frac{\partial c_3}{\partial t} - rs\frac{\partial c_2}{\partial s} - (a-bv)\frac{\partial c_2}{\partial v} - \frac{1}{2}s^2v\frac{\partial^2 c_2}{\partial s^2} - \rho\sigma sv\frac{\partial^2 c_2}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c_2}{\partial v^2} + rc_2 = 0 \\ P^4 &= \frac{\partial c_4}{\partial t} - rs\frac{\partial c_3}{\partial s} - (a-bv)\frac{\partial c_3}{\partial v} - \frac{1}{2}s^2v\frac{\partial^2 c_3}{\partial s^2} - \rho\sigma sv\frac{\partial^2 c_3}{\partial s\partial v} + \frac{1}{2}\sigma^2v\frac{\partial^2 c_3}{\partial v^2} + rc_3 = 0 \\ &\vdots \end{aligned} \quad (19)$$

and the required approximate solution obtained is

$$c(s, v, t) = \sum_{n=0}^{\infty} P^n c_n \quad (20)$$

Executing the HPM methodology as described above, MAPLE 18 software is brought into play. Using the following parameters $a = 0.16, b = 0.055, \delta = 0.9, \rho = -0.5, T = 15, K = 100$, we have;

Table 1: Numerical results for MHPM

$C(t, s, v)$	HPM(Biazar et al , 2015)	MHPM	Error MHPM-HPM
C(1, 10, 0.1)	413.2583333	3.796325819 E5 10 ¹	409.462007
C(2, 50, 0.2)	81406.34666	5.476248101 E6 10 ⁶	81400.8704
C(4, 70, 0.3)	1.734519680 E6	4.40834821 E6 10 ⁶	2.6738285
C(6, 90, 0.4)	1.335877440 E7	-8.9567091 E5 10 ⁶	10.2925865
C(8, 120, 0.5)	7.485848702 E7	-1.24852579 E7 10 ⁶	8.73437449
C(10, 150, 0.6)	2.937374856 E8	-3.139821922 E7 10 ⁶	6.07719678
C(14, 200, 0.8)	2.198733197 E9	-7.467534155 E7 10 ⁷	9.66626735

7. Modified VIM for Heston SPDE

We now proceed to modify the He's VIM using the MNPs as modifier and basis function via the orthogonal collocation method (OCM). Given the general formulation of the VIM in (8), the correction function as related to the HSPDE is thus given as

$$c_{(n+1)}(s, v, t) = c_n(s, v, t) + \int_0^t \lambda(\xi) \left[\frac{\partial c}{\partial \xi} - rs \frac{\partial c}{\partial s} + (a - bv) \frac{\partial c}{\partial v} + \frac{1}{2} s^2 v \frac{\partial^2 c}{\partial s^2} + \rho \sigma s v \frac{\partial^2 c}{\partial s \partial v} + \frac{1}{2} \sigma^2 v \frac{\partial^2 c}{\partial v^2} + rc \right] d\xi \quad (21)$$

Hence, the initial approximation for the modified VIM is given by (17). From (8), we have that $\lambda(\zeta)$ is the general Langrange multiplier which can be obtained optimally via the variational theory. If we set

$$\lambda(\zeta) = -1 \quad (22)$$

and substituting (22) into (8) gives,


$$c_{(n+1)}(s, v, t) = c_n(s, v, t) - \int_0^t \left[\frac{\partial c}{\partial \xi} - rs \frac{\partial c}{\partial s} + (a - bv) \frac{\partial c}{\partial v} + \frac{1}{2} s^2 v \frac{\partial^2 c}{\partial s^2} + \rho \sigma s v \frac{\partial^2 c}{\partial s \partial v} + \frac{1}{2} \sigma^2 v \frac{\partial^2 c}{\partial v^2} + rc \right] d\xi \quad (23)$$

which is the MVIM with initial approximation $c_0(s, v, t) = 392.00000s^2$. Evaluating (23) with the aid of MAPLE 18 application software for $n \geq 0$ and with the following parametric values $a = 0.16, b = 0.055, \delta = 0.9, \rho = -0.5, T = 15, K = 100$ yields the following approximation


Table 2: Numerical results for MVIM

$C(t, s, v)$	VIM(Biazar et al , 2015)	MVIM	Error MVIM-VIM
C(1, 10, 0.1)	209.5038332	38456.89887	38247.395
C(2, 50, 0.2)	82474.33729	9.740327147 10 ⁶	82464.597
C(4, 70, 0.3)	1.147814970 E6	1.873262944 10 ⁶	0.725448
C(6, 90, 0.4)	1.950175983 E7	3.017060928 10 ⁶	1.0668849
C(8, 120, 0.5)	7.79650567 E7	5.276105707 10 ⁶	2.52039996
C(10, 150, 0.6)	2.317375113 E8	8.112385120 10 ⁶	5.79501
C(14, 200, 0.8)	2.576414314 E9	1.380834321 10 ⁷	1.19557999

Hence the approximate solution of the HSPDE given by the MVIM is thus



MHPM for HSPDE



MVIM for HSPDE

Figure 1: HSPDE given by the MVIM.

8. Conclusion

The MNPs being a new orthogonal polynomial was used as a modifier and basis function for the He's HPM and VIM via the OCM after which both modified methods were applied on the HSPDE. The compatibility of the MNPs and the numerical method created a new iterative method which was able to approximate the Heston Stochastic Partial Differential Equation to its exact solution with the MVIM standing out as a faster approximant for the HSPDE than the MHPM.

References

- 1 . Gupta,P.K.,Yidirim, A.,and Rai, K.N. (2012). Application of He's Homotopy Perturbation Method for Multi-Dimensional Fractional Helmholtz Equation. *International Journal of Numerical Methods for Heat and Fluid Flow* 22(4): 424-435
- 2 . Syed T. Mohyud-Din, Muhammed, A. N. (2009). Homotopy Perturbation Method for Solving Partial Differential Equations. *Semantic Scholar*
- 3 . Babolian, E., Dastani, N. (2011). Numerical Solutions of Two- Dimensional Linear and Nonlinear Volterra Integral Equations: Homotopy Perturbation Method and Differential Transform Method. *International Journal of Industrial Mathematics* 3(3): 157-167.
- 4 . Alaje, A.I., Olayiwole, M.O., Adedokun, K.A., Adedeji, J.A. and Oladapo, A.O. (2022). Modified Homotopy Perturbation Mwthod and its Application to Analytical solutions of Fractional - order Korteweg-de Vries Equation. *Beni-Suef University Journal of Basic and Applied Science*. 11:139(2022).
- 5 . Qayyum, M. and Oscar, I. (2021). Least Square Homotopy Perturbation Method for Ordinary Differential Equations. *Journal of Mathematics*. <https://doi.org/10.1155/2021/7059194>.
- 6 . Yadav, N., Singh, M., Singh, S., Singh, R. and Kumar, J. (2023). A note on homotopy perturbation approach for nonlinear coagulationequation to improve series solutions for longer times. *Chaos, Solitons and Fractals*. 173(2023). <https://doi.org/10.1016/j.chaos.2023.113628>.
- 7 . He, J. H (1999). Variational Iteration Method - A kind of Non-Linear Analytical Technique: some examples. *International Journal of Non-Linear Mechanics* . 34(4): 699-708
- 8 . He, J. H., Wazwaz, AM. and Xu, L. (2007). The Variational Iteration Method: Reliable, Efficient and Promising. *International Journal of Computers and Mathematics with Applications*. 54: 879-880
- 9 . Safari, M. (2011). Application of He's Variational Iteration Method for the Analytical Solution of Space Fractional Diffusion Equation. *Journal of Applied Mathematics* 2: 1091-1095
- 10 . Baghani, M., Fattahi, M. and Amjadan, A. (2012) Application of the Variational Iteration Method for Non-linear Free Vibration of Conservative Oscillators. *Scientia Iranica* 19(3): 513-518
- 11 . Elsheikh, A.M. and Elzaki, T.M. (2016) Modified Variational Iteration Method for Solving Fourth Order Parabolic PDEs with Variable Coefficients. *Global Journal of Pure and Applied Mathematics* 12(2): 1587-1592
- 12 . Abassy, T.A. (2012) Modified Variational Iteration Method (non-homogeneous initial value problem). *Mathematical and Computer Modelling* 55(3-4): 1222-1232
- 13 . Biazar, J., Goldoust, F. and Mehrdoust F. (2015) On the Numerical Solutions of Heston Partial Differential Equation. *Mathematical Science Letters* 4(1): 63-68

- 14 . Onyeoghane, J.N. and Njoseh, I.N. (2021): Modified Variational Iteration Method for the solution of the Time Fractional Nagumo Equation. *Nigerian Journal of Science and Environment*. 19(2).
- 15 . Onyeoghane, J.N and Njoseh, I.N. (2019): Modified Numerical Techniques for the Solution of Heston Stochastic Partial Differential Equation via Mamadu-Njoseh Polynomials. *Bulletin of the Science Association of Nigeria*. 30: 186-195.
- 16 . Onyeoghane, J.N. and Njoseh, I.N. (2020): Modified Variational Iteration Method for the solution of the Solution of Heston Stochastic partial Differential Equation using the Mamadu-Njoseh Orthogonal Polynomials. *Nigerian Journal of Science and Environment*. 18(1) (2020).
- 17 . Ajuhari, A., Maturi, D. and Alsheri, H. (2022). Variational Iteration Method for solving Boussinesg Equation using Maple. *Applied Mathematics*. 13(12). Doi:10.4236/am.2022.1312060.
- 18 . Islam, D.U. and Shirivastava, A. (2024). Fredholm Integral Equations by Variational Iteration Method. *International Journal of Scientific Research in Science Engineering and Technology*.11(4):215 - 217. doi:<https://doi.org/10.32622/IJSRET1104215>
- 19 . Tomar,S., Singh, M., Vajravelu,K. and Ramos. H. (2022). Simplifying the Variational Iteration Method: A new approach to obtain the Lagrange Multiplier. *Mathematics and Computer in Simulation*. 204(C). 640 - 644. <https://doi.org/10.1016/j.matcom.2022.09.003>.
- 20 .Vikash, K.S. and Maroju, B. (2023). New Development of Variational Iteration Method using Quasilinearization method for solving Nonlinear problems. *Mathematics* 2023. 11(4), 935. <https://doi.org/10.3390/math11040935>.
- 21 . Njoseh, I.N. and Mamadu, E.J. (2016) Numerical Solutions of Fifth Order Boundary Value Problems using Mamadu-Njoseh Polynomials. *Science World Journal* 11(4): 21-24
- 22 . Mamadu, J.E. and Njoseh, I.N. (2016) Numerical Solution s of Volterra Equations using Galerkin Method with certain Orthogonal Polynomials. *Journal of Applied Mathematics and Physics* 4:376-382. <http://dx.doi.org/10.4236/jam.2016.43047>
- 23 . Mamadu, E.J., Njoseh, I.N., and Ojarijkre, H.I. (2022): Space Discretization of Time-Fractional Telegraph Equation with Mamadu-Njoseh Basis Function. *Applied Mathematics*, 2022.13,760 - 773. <https://doi.org/10.4236/am.2022.13760>
- 24 . Mamadu, E.J., and Ojarijkre, H.I. (2023): Gauss-Mamadu-Njoseh Quadrature Formula for Numerical Integration Interpolation. *Journal of Advances in Mathematics and Computer Science*, 38(9):128 - 134.
- 25 . Alziary,B. and Takac, P. (2017) On the Heston Model with Stochastic Volatility: Analytic Solutions and Complete Markets. <https://arxiv.org/abs/1711.04536v1>, retrieved 20 March, 2018.
- 26 . He, J. H (1998) An Approximate Solution Technique Depending on an Artificial Parameter: A Special Example. *Communications in Nonlinear Science and Numerical Simulation* 3(2): 92-97. DOI: 10.1016/S1007-5704(98)90070-3
- 27 . He, J. H., Wu, GC, and Austin, F. (2010) The Variational Iteration Method which should be followed. *Non-linear science letters A. Mathematics, physics and mechanics* 2010, 1(1): 1-30