

## **Histopathological and Anti-hemorrhoid Effect of the Leaf Extract of *Ocimumgratissimum* in Croton oil-Induced Hemorrhoid**

### **ABSTRACT**

The use of medicinal plants in the treatment of various diseases dates back into centuries. Hemorrhoids are swollen veins in the lower rectum or anus, ranging from being acute to a complicated state; and affecting several millions of people worldwide. This study evaluates the histopathological and anti-hemorrhoid effect of the leaf extract of *Ocimumgratissimum* in croton oil-induced hemorrhoid in wistar rats. Hemorrhoids was induced to the experimental animals using croton oil, following standard methods. Thirty five (35) wistar rats were randomly grouped into five classes (A, B, C, D and E) of seven rats each. Groups A, B and C served as normal control (un-induced, untreated), negative control (induced, untreated) and standard control (induced, treated with standard drug; Daflon) respectively while groups D and E served as treatment groups and were administered 100 and 400 mg/kg dose of ethanol leaf extract of *Ocimumgratissimum* respectively. All the rats (except the group A) animals were induced with croton oil and were thereafter treated for fourteen days (two weeks). The body weight of the animals were monitored throughout the experimental period and percentage rectal water content, anorectal coefficient as well as histopathological assay were carried out at the end of the fourteen days, using standard methods. Results showed a significant decrease ( $p>0.05$ ) in body weight of the untreated group while there was a marked increase in the body weight of the animals in the extract treated groups from day 1 to the 14<sup>th</sup> day. The result of the percentage rectal water content as well as anorectal coefficient of the untreated group showed a significant increase ( $p<0.05$ ) (76%) when compared to the extract treated groups (67%), while the histopathological studies showed a reversal of the inflammatory and necrotic conditions of the induced anorectal portion for the extract treated animals. From the result of this study, it could be concluded that the leaf extract of *Ocimumgratissimum* could help to mitigate anorectal disorders and contribute significantly to improved health.

**Keywords:** *Ocimumgratissimum*, Hemorrhoid, Inflammation, Health

### **INTRODUCTION**

Hemorrhoid is one of the most common inflammatory disease which can be characterized by alteration in vasculature of the anal canal including blood vessels, supporting tissues, muscles and elastic fibers. According to the report of National Centre for Health Statistics, the prevalence of hemorrhoid is 3.82% in United States [1] and 0.36% in India that frequently appears in the age of 45 to 65 years [2]. Free radical generation is the primary reason for initiation of many physiological and pathological disorders like hemorrhoids. Involvement of free radicals in the precipitation of hemorrhoids is well documented in the literature [3, 4, 5]. It is well known fact that excess concentration of free radicals is a consequence of improper balance between reactive oxygen species and their metabolites. Antioxidants neutralize free radicals and thus play major role in the eradication of these free radicals and hence are involved in the management of diseases [6], including hemorrhoids. Unavoidable and serious adverse effects associated with commercial or synthetic antioxidants are becoming a major concern for the researchers to develop natural antioxidants. Herbal extracts rich in phytoantioxidants like polyphenols, flavonoids, tannins and other related compounds are known to possess positive health effects and eventually reduces the incidence of diseases [7]. Therefore much attention has been focused on the use of natural antioxidants that can provide more significant health benefits with minimal toxicities [8].

*Ocimum gratissimum* (*O. gratissimum*) is a herbaceous plant that belongs to the Lamiaceae family and is sometimes referred to as smell leaf, African basil, camphor basil, basil leaf, or ram tulsi [9]. This particular species of tropical plant is popularly known as "scent leaf," a term that Nigerians find particularly endearing. It is known as Nchanwu in Igbo, Daidoya in Hausa, and Efirin in Yoruba in native Nigeria [10, 11, 12]. It is a small to average-sized plant with leaves that is comparable to cloves-like

flavor and aroma, making it a significant herb in various cuisines [13]. In West Africa, the plant is commonly grown for both culinary and medicinal reasons [9], usually in gardens around community huts. *O. gratissimum* used as a culinary ingredient in salads, soups, pastas, vinegars, and jellies in many parts of the world. Nutritionally, the plants have been documented to be rich in microelements sufficient for improved health and vitality [12].

Additionally, the plant has been documented to contain a number of phytochemicals such as flavonoids and polyphenols [14, 15] which has been reported to be responsible for its many pharmacological activities such as hypoglycaemic activities [16], anti-inflammatory activities [17], anti-anaemic, hepatoprotective [18], anti-hypertensive [19], antibacterial [15], antifungal [20] as well as exhibits many other pharmacological activities. Although there have been several reports on its other pharmacological activities, its histopathological effects, especially in hemorrhoids disease remain to be explored. Hence, this study was designed to evaluate the histopathological effect of the leaf extract of *Ocimumgratissimum* in Croton oil-Induced Hemorrhoid.

## **MATERIAL AND METHODS**

### **Sample Collection and Preparation of Extract**

Fresh leaves of *Ocimumgratissimum* were purchased from the daily market, Mgbakwu in Awka North Local Government Area of Anambra State, Nigeria and were authenticated by a Taxonomist, MrIroka Finan of the Department of Botany, Nnamdi Azikiwe University, Awka, and a voucher specimen was deposited with Herbarium no NAUH 35B. The leaves were then detached from the stalk, rinsed with distilled water and were oven-dried at 40 °C and weighed (300 g). The weighed powdered sample (300 g) was then used for the extraction with a solvent combination

of ethanol and water (7:3) (2500 ml) for 48 hr via maceration in an unheated medium. The mixture was decanted and filtered using sterile Whatman paper No. 1. The filtrate was there after evaporated to dryness with the aid of a rotary evaporator set at 50 °C to obtain crude ethanol extract which was carefully preserved for further analysis.

## **ANIMAL STUDIES**

### **Procurement of Study Animals**

Wistar albino rats (35) weighing approximately 150g were purchased from Chris Farm Ltd Mgbakwu. The rats were kept in standard cages with saw dust as bedding, and at standard room temperature as well as standard housing conditions of 12:12 light: dark cycles and fed with standard rat pellets and water *ad libitum*. The animals were allowed to acclimatize for seven days.

### **Dose Preparation and Treatment**

The hydro-ethanolic leaf extract of *Ocimum gratissimum* was prepared with distilled water in two doses (100 and 400) mg / kg, Daflon(1 g/kg) was used as a reference drug and distilled water was used as a vehicle for the untreated group. The animals were administered the extract and drug orally for fourteen consecutive days with water *per os* and feed *ad libitum* [21].

### **Experimental Design**

The animals were randomly grouped into five, with seven animals in each group, and the treatment was as follows: Groups A served as normal control and was administered distilled water, group B served as negative control and was induced with hemorrhoid without any treatment, group C served as the standard control and was induced and treated with standard drug (Daflon), groups D and E served as the treatment groups and was administered 100 and 400 mg/kg ethanol extract of *O. gratissimum*.

### **Induction of Hemorrhoid**

By administering a croton oil formulation, hemorrhoids were caused in all subjects except group A (Normal Control). A cotton swab (4mm in diameter) was placed into the anus and left there for 10seconds after being soaked in Croton oil preparation (0.16ml). Within 7-8 hours of the introduction of croton oil, a linear progression of oedema was seen. Body weight was checked weekly while treatment was given daily for 14 days [21].

### **Monitoring and Weight Evaluations**

The rats were physically monitored and body weights were measured weekly. Data was collected throughout the duration of the experiment and recorded.

### **Treatment and Anorectal Evaluation**

After 24 hours of induction, two rats were taken from group A and used for pilot studies; while two rats each were taken from groups B to D and used for microscopy, histology and wet weight determination respectively as baseline studies. Treatment was then given to all the groups daily for 14 days. The standard drug used for group C was Daflon at the dose of 1g (3x per day) for 4 days, then 1g (2x per day) subsequently while 100mg/kg of extract was used to treat group D and 400mg/kg of extract was used for group E. The control group A was given distilled water only. On day 14, all animals were euthanized under anesthesia. The anorectal portions were subsequently cut and evaluated [21].

### **Determination of Wet Weight and Anorectal Coefficient**

Twenty millimeter of the anorectal portion of the first set was cut and the wet weight recorded before drying. After drying, the dry weight was also recorded. The values were used to calculate the % rectal water content by the method of Liu *et al.* [21].

The % rectal water content was calculated using the formula,

$$\% \text{ rectal water content} = \frac{\text{Wet weight} - \text{Dry weight}}{\text{Wet weight}} \times \frac{100}{1}$$

The Anorectal Co-efficient (ARC) was calculated using the formular,

$$\frac{\text{Water Content (ml)}}{\text{Body weight (g)}}$$

### **Histologic and Microscopic Analysis**

Twenty millimeter of the anorectal portion of the second set of rats was cut and fixed in 10% formal saline and used for microscopic and histological evaluation by the method of Carleton [22] and Ajibade *et al.*, [23] respectively.

Histological observation was performed for inflammation, congestion, hemorrhage, vasodilation and necrosis.

Microscopy was done, and the section's permanent micrographs and morphological observations were recorded.


### **Data Analysis**

The values obtained and the significance between the treated and control group was analyzed by one-way ANOVA using the SPSS version 17 and  $P < 0.05$  was considered to be statistically significant.

## **RESULTS**

The result showing the effect of *O. gratissimum* leaf extract in Croton oil-Induced Hemorrhoid on body weight of Wistar rats is presented in figure 1.0. Result showed an increase in the weight of extract treated groups (D and E) after 14 days while there


was a significant decrease in the weight of the untreated animals (group B) after 14 days. From the bar chart, a total of about 1.9% weight loss was calculated for the group B animals after the period of the experiment while 18% weight gain was observed in the normal control (group A). However, extract treated groups (D and E) had 12.3% and 13.2% weight increase respectively.



**Figure 1.0: Effect of Oral Administration of the leaf extract of *Ocimumgratissimum* in Croton oil-Induced Hemorrhoid on body weight of Wistar Rat**

The result showing the effect of *O. gratissimum* leaf extract in Croton oil-Induced Hemorrhoid of percentagerectal water content in Wistar rat is presented in figure 2.0. Result showed a significant increase in the percentagerectal water content of group B ( $76.00 \pm 1.00$ ) (untreated) animals compared to the extract treated groups D and E


( $67.00 \pm 2.34$ ) while the standard control group (group C) has percentage rectal water content of ( $67.75 \pm 1.14$ ).




**Figure 2.0: Effect of Oral Administration of the leaf extract of *Ocimum gratissimum* in Croton oil-Induced Hemorrhoid on percentage rectal water content of Wistar Rat**

The result showing the effect of *O. gratissimum* leaf extract in Croton oil-Induced Hemorrhoid on anorectal coefficient in Wistar rat is presented in figure 3.0. Result showed a significant increase in the anorectal coefficient of group B (0.0056) (untreated) animals compared to the extract treated groups D and E (0.00415) and

(0.00416) respectively while the normal control group (group A) has anorectal coefficient of 0.003825.



**Figure 3.0: Effect of Oral Administration of the leaf extract of *Ocimumgratissimum* in Croton oil-Induced Hemorrhoid on anorectal coefficient in Wistar rat**



**Figure 4.0: Effect of Oral Administration of the leaf extract of *Ocimum gratissimum* in Croton oil-Induced Hemorrhoid on microscopy and histological analysis in Wistar rat**

## DISCUSSION

From time immemorial, medicinal plants have contributed significantly to the prevention and treatment of diseases including hemorrhoids. Plants are known to contain bioactive compounds which help them to perform these functions [24]. Assessing the impacts of plants' administration on hemorrhoid disorder remains crucial in evaluating their therapeutic index against inflammatory diseases including hemorrhoid. Thus, the possible ameliorative effects of leaf extract of *Ocimum gratissimum* on hemorrhoid were carried out.

Overall, the results of this finding showed that extract of *Ocimum gratissimum* could help to ameliorate inflammatory disorders especially hemorrhoids. Figure 1.0 showed the effect of *O. gratissimum* leaf extract in croton oil-induced hemorrhoid on body weight of wistar rats. From the result of the study, there was significant increase in the body weight of the experimental animals in the extract treated groups from day 1 to day 14 of the experiment while a decline in body weight was observed for the untreated control (group B). This is consistent with the findings of Anyanwu and Ogbadu, [25]. It is worthy of note that hemorrhoids are a very common anorectal condition that has been defined as the symptomatic enlargement and distal displacement of the normal anal cushions and characterized with abnormal dilatation and distortion of the vascular channel, together with destructive changes in the supporting connective tissue within the anal cushion [26] and consequently leading to loss of water in a biological system.

Furthermore, the result of this study is also in agreement with previous finding which indicated that inflammation demands the convergence of more cells at the site of activity by way of secreting inflammatory cytokines and chemokines which can cause the animals to lose weight due to high energy expended in the process [27]. Earlier

reports have indicated that phytochemicals help to reduce cytokine gene expression of IL-1 $\beta$ , IL-6 and TNF- $\alpha$  which are pro-inflammatory cytokines and increase IL-10 which is anti-inflammatory cytokine in LPS-activated cells [28]. The anti-inflammatory response of the extract could be vital in averting the weight loss among the treated animals (Group D and E) and possibly enhanced its anti-hemorrhoid capacity. The response of the extract is supported by Sorhuet *al.* [29] who reported that *Ocimumgratissimum* increased IL-10 expression while simultaneously decreasing IL-1 $\beta$  expression in experimental animals.

As depicted in figures 2.0 and 30.0, there was significant increase in the percentage rectal water content and anorectal coefficient of untreated group (group B) compared to the extract treated groups respectively. Percentage rectal water content refers to the amount of water present in the stool within the rectum, expressed as a percentage of the total stool weight. Since hemorrhoids occur when veins in the rectum or anus become swollen, percentage rectal water content therefore plays a significant role in hemorrhoid development and prevention. Induction of hemorrhoid through the administration of croton oil in the experimental animals caused a significant increase in the rectal water content of the animals above the reference range of 60-70%. However, following the administration of the plant extract over the treatment period, the percentage rectal water content was able to be brought to normal (67%). This is an important observation as this could serve as a potential anti-hemorrhoid agent for clinical practice. This result is consistent with the reports of Anusmitha *et al.*, [30].

Similarly, anorectal coefficient of the untreated group was observed to be significantly high compared to the extract-treated groups. Anorectal coefficient is said to be the ratio of the anal canal length to the rectal length, typically measured during anorectal manometry and it helps to diagnose and assess anorectal disorders, including

hemorrhoids. Since a high coefficient (>0.8) has been implicated in the pathogenesis of hemorrhoids, the ability of the plant extract to significantly reduce this coefficient in the treatment group is suggestive of its potency to mitigate anorectal disorders. Huiyi, [31] made similar assertion.

The morphological Observations and the histopathology assessments (figure 4.0) displayed reversal of the inflammatory and necrotic conditions of the induced anorectal portion for the Group E animals that were treated with 400mg/kg dose of extract since the photomicrograph showed that the cytoarchitecture of their induced anorectal tissue was restored and normalised. This is a possible indication of the hemorrhoid healing effect of the extract and is supported by Lohsiriwat [32], Faujdar *et al.* [1] and Margetis [33] who identified inflammatory and necrotic conditions of the anorectal tissue as characteristic pathologic presentations of hemorrhoid. Also, Faujdar *et al.* [1] noted that the degree of restoration or normalisation of the anorectal cytoarchitecture after treatment is indicative of the degree of hemorrhoid healing.

## **CONCLUSION**

From the observations in this study, it can be concluded that the ethanol leaf extract of *Ocimumgratissimum* showed significant medicinal properties via the reparation of anorectal tissue and therefore could help to mitigate anorectal disorders especially hemorrhoids and serve as potential agent for inflammatory disorders.

### **Disclaimer (Artificial intelligence)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

### **Ethical Approval**

Animal Ethic committee approval has been collected and preserved by the author(s)

## REFERENCES

1. Faujdar, S., Sati, B., Sharma, S., Pathak, A.K. and Paliwal, S.K. (2019). Phytochemical evaluation and anti-hemorrhoidal activity of bark of *Acacia ferruginea*. *Journal of Traditional and Complementary Medicine*, 9(2): 85-89.
2. Sowndhararajan, K., Hong, S., Jhoo, J. W., Kim, S. and Chin, N. L. (2015). Effect of acetone extract from stem bark of *Acacia species* (*A. dealbata*, *A. ferruginea* and *A. leucophloea*) on antioxidant enzymes status in hydrogen peroxide induced HepG2 cells. *Saudi Journal of Biological Science*, 22(6):685-691.
3. Ojo, O., Oloyede, O., Olarewaju, O., Ojo, A., Ajiboye, B. and Onikanni, S. (2013). *In-vitro* antioxidant and free radical scavenging activities of *Ocimum gratissimum*. *World Journal of Pharmaceutical research*, 2(6): 1899-1912.
4. Scaldaferrri, F., Ingravalle, F., Zinicola, T., Holleran, G. and Gasbarrini, A. (2018). Medical Therapy of Hemorrhoidal Disease. *Hemorrhoids*, 2(3): 319-356.
5. Jorge, B. C., Júnior, A. J., Stein, J., Reis, A.C., Moreira, S., Manoel, B., Mota, J., Kassuya, C. A. and Arena, A.C. (2022). Safety assessment of the ethanolic extract from *Piper vicosanum* leaves in male rats. *Phytomedicine*, 2: e100165.

6. Onwubuya, E. I. and Oladejo, A. A. (2022). Possible Effects of Free Radical Scavengers on Dissolution of Inflammatory Granuloma: An Experience with Wistar Rats. *Curr Res Cmpl Alt Med* 6: 149.
7. Kalu, O. A., Onwubuya, E. I., Oladejo, A. A., Okeke, C. B., and Ezenwelu, C. O. (2024). "Investigating the CardioProtective Activity of *Volvariella volvaceain* Isoproterenol-Induced Myocardial Infarction Using Animal Model: An *In Vivo* Approach". *Journal of Complementary and Alternative Medical Research*, 25 (7):73-83.
8. Oladejo, A.A., Onwubuya, E.I., Kalu, O. A., Okeke,C.B., Okeke,C.M. and Ikemefuna, P.N. (2024). "Investigating the Cardio-Protective Activity of *Agaricus bisporus* in Wistar Rats Induced With Myocardial Infarction". *Cardiology and Angiology: An International Journal*, 13 (3):57-66.
9. Kalita, M., Devi, N. and Narzary, D. (2023). *Ocimumgratissimum L. ssp. gratissimum var. macrophyllum Briq.* (Lamiaceae: Nepetoideae: Ocimeae) a new record from northeastern India. *Journal of Threatened Taxa*, 15(10): 24086-24091.
10. Zaku, S.G., Emmanuel, S., Tukur, A.A. and Kabir, A. (2015). *Moringa oleifera*: An underutilized tree in Nigeria with amazing versatility: A review. *AfricanJournal of Food Science*, 9(9): 456-461.
11. Imosemi, I.O. (2020). A review of the medicinal values, pharmacological actions morphological effects and toxicity of *Ocimumgratissimum* Linn. *Eur J Pharm Med Res*, 7(7): 29-40.
12. Moneme, E. C., Nwaka, A. C., Ajakpofo, F. O. and Anyanwu, R. O. (2024). Evaluation of Nutritional Composition of *Ocimumgratissimum* Leaf Extract. *International Journal of Biochemistry Research and Review*, In-press.

13. Puneet, S. and Monika, S. (2022). Exploration of antifungal potential of extracts of leaves of *Ocimumgratissimum* against *Microsporungypseum*. *Research Journal of Pharmacy and Technology*, 4(2): 234.250.
14. Irondi, E.A., Agboola, S.O., Oboh, G. and Boligon, A.A., (2016). Inhibitory effect of leaves extracts of *Ocimumbasilicum* and *Ocimumgratissimum* on two key enzymes involved in obesity and hypertension *in vitro*. *J. Intercult. Ethnopharmacol*, 5: 396–402.
15. Melo, R.S., Azevedo, A.M.A., Pereira, A.M.G., Rocha, R.R., Cavalcante, R.M.B., Matos, M.N.C., Lopes, P.H.R., Gomes, G.A., Rodrigues, T.H.S., Dos-Santos, H.S., Ponte, I.L., Costa, R.A., Brito, G.S., Júnior, F.E.A.C. and Carneiro, V.A. (2019). Chemical composition and antimicrobial effectiveness of *Ocimumgratissimum* L. Essential oil against multidrug-resistant isolates of *Staphylococcus aureus* and *Escherichia coli*. *Molecules*, 3864:2–17.
16. Shittu, S.T., Shittu, S.A., Olatunji, A.A. and Oyeyemi, W.A. (2019). *Ocimum gratissimum* leaf extract may precipitate infertility in male diabetic Wistar rats. *JBRA Assisted Rep.* 23:34–44.
17. Ajayi, A.M., Umukoro, S., Ben-Azu, B., Adzu, B. and Ademowo, O.G. (2017c). Toxicity and protective effect of phenolic-enriched ethylacetate fraction of *Ocimumgratissimum* (linn.) leaf against acute inflammation and oxidative stress in rats. *Drug Dev. Res.* 78: 135–145.
18. Akara, E.U., Okezie, E., Ude, V.C., Uche-Ikonne, C., Eke, G. and Ugbogu, A.E. (2021). *Ocimumgratissimum* leaf extract ameliorates phenylhydrazine induced anaemia and toxicity in Wistar rats. *Drug Metab. Pers. Ther.* 10(3):342-354.

19. Shaw, M., Wu, J. and Wang, M. (2017). Antihypertensive effects of *Ocimum gratissimum* extract: angiotensin-converting enzyme inhibitor in vitro and in vivo investigation. *Journal of Functional Foods*, 35:68–73.
20. Mohr, F.B., Lermen, C., Gazim, Z.C., Gonçalves, J.E. and Alberton, O. (2017). Antifungal activity, yield, and composition of *Ocimumgratissimum* essential oil. *Genet. Mol. Res*, 11(6): 23-31.
21. Liu, H., Yang, M. and Qiu, G.P. (2012). Aquaporin 9 in rat brain after severe traumatic brain injury. *Arq.Neuropsiquiatr.*, 20(3): 214-220.
22. Carleton's Histological Techniques, (1967). 4<sup>th</sup> ed. Oxford University Press, NY. U.S.A. P. 279-280.
23. Ajibade, A.J., Esho, J.O., Kehinde, B.D. and Adeleye, O.O. (2019). Histological and Biochemical effects of mercury chloride on the kidney of adult Wister rats. *EAS J. Pharm. Pharmacol*,1(1): 21-27.
24. Onwubuya, E. I., Oladejo, A. A., Kalu, O. A. and Ndubueze, I. K. (2023). Possible Ameliorative and Preventive Effect of Leaf Extract of *Andrographis paniculata* on Atherosclerosis: An Experience with Isoproterenol Induced Wistar Rats. *Journal of Complementary and Alternative Medical Research*, 23(1): 1-8.
25. Anyanwu, G. O. and Ogbadu, D. E. (2024). Anti-obesity effect of *Ocimum gratissimum* leaf powder supplementation onhigh fat diet-induced obesity in male Wistar rats. *Trop J Nat Prod Res*, 8(6):7397-7402.
26. Lohsiriwat, V.(2017). Hemorrhoids: From basic pathophysiology to clinical management. *World J Gastroenterol*,18(17):20–27.

27. Huang, C.M. and Lee, T.T. (2018). Immunomodulatory effects of phytochemicals in chickens and pigs; a review. *Asian-Australasian Journal of Animal Sciences*, 31:617-627.
28. Lee, M., Lin, W., Wang, S., Lin, L., Yu, B. and Lee, T. (2018). Evaluation of potential antioxidant and anti-inflammatory effects of *Andropogon squarrosus* powder and the underlying molecular mechanisms via Nrf2 and NF- $\kappa$ B dominated pathways in broiler chickens. *Poultry science*, 97:2419-2434.
29. Sorhue, U.G., Onainor, E.R., Moemeka, A.M., Peterson, I.E. (2021). Dietary inclusion of Scent leaf meal (*Ocimum gratissimum*) affects immune genes expression in chicken spleen at 28 and 56 days. *Animal Review*.8 (1): 10-19.
30. Anusmitha, K. M., Aruna, M., Job, J. T., Narayanankutty, A., Benil, P. B., Rajagopal, R., Alfarhan, A. and Barcelo, D. (2022). Phytochemical analysis, antioxidant, anti-inflammatory, anti-genotoxic, and anticancer activities of different *Ocimum* plant extracts prepared by ultrasound-assisted method. *Physiological and Molecular Plant Pathology*, 117: 101746.
31. Huiyi, J. (2013). Theoperamide treats neonatal hypoxic-ischaemic encephalopathy by postsynaptic H1 receptors. *Neural regeneration research*. 8:1814-1822.
32. Lohsiriwat, V. (2012). Hemorrhoids: From basic pathophysiology to clinical management. *World J. Gastroenterol*, 18(17): 2009-2017.
33. Margetis, N. (2019). Pathophysiology of internal hemorrhoids. *Ann Gastroenterol*, 32(3): 264-272.