

PHYTOTHERAPY ANTI-DIABETIC: ETHNOBOTANICAL SURVEYS AMONGIN THE DISTRICT OF MBOUR (SENEGAL)

ABSTRACT

Background: Despite the promises of a wide range of conventional medications, the prevalence of diabetes mellitus is steadily rising. Therefore, scientific research on the antidiabetic effects of plants are required.

Aim: The objective of this study is to evaluate the diversity of medicinal plants used to treat diabetes mellitus.

Methodology : Ethnobotanical surveys were carried out among 30 randomly selected traditional practitioners using semi-structured interviews in the Mbour district.

Results: A total of 14 plant species belonging to 10 families was established. The most commonly used plant parts were leaves (44.20%), followed by barks (31%), roots (20.90%) and seeds (3.90%). The most common preparation techniques are infusion (61.20%), decoction (31%) and maceration (7.80%). The plants most frequently cited by traditional practitioners were the following : *Neocaryamacrophylla*, *Moringa oleifera*, *Terminaliaavicennioides* and *Sclerocaryabirrea*. And the least cited plants are : *Garcinia kola*, *Anacardium occidentale*, *Vahliadichotoma*, *Streptogynegerontogaea*, *Combretumlecardii*, *Chrozophora senegalensis*, *Boscia senegalensis*, *Chrysobalanusicaco*, *Allium cepa* and *Jatropha curcas*.

Conclusion: These results may be a database for the discovery of new molecules with **antimalarial potential** and the development of improved traditional medicines (ITM).

Keywords: Ethnobotany, medicinal plants, diabetes mellitus, traditional practitioners, Mbour, Senegal.

1. INTRODUCTION

Diabetes is a public health problem. In 2014, the WHO estimated that 8.5% of the adult population (aged 18 and over) had diabetes. In 2015, diabetes was the direct cause of 1.6 million deaths, and in 2012 hyperglycaemia caused a further 2.2 million deaths [1]. Diabetes mellitus (DM) is a group of metabolic diseases characterized by hyperglycemia resulting from defects in insulin secretion and/or insulin action [2]. The chronic hyperglycemia in diabetes is associated with long-term damage, dysfunction, and failure of different organs, especially the eyes, kidneys, nerves, heart, and blood vessels. Failure of these organs is frequently associated with development of macro- and microvascular diseases [3]. Now a days, DM is becoming a disease of major concern both globally and regionally and is a leading cause of death in most countries [4]. It is one of the four major non-communicable diseases comprising, cardiovascular diseases, cancers and chronic respiratory diseases, which jointly contributes to 63% according to epidemiological predictions by the World Health Organisation (WHO), diabetes could be the seventh leading cause of death by 2030, due to its increasing prevalence. In 2015, approximately 5.0 million deaths were attributed to diabetes, albeit in the same year, more than 12% of the global health expenditure was dedicated to coping with the disease and its complications [5]. Diabetes complications are common among patients with type 1 or type 2 diabetes but, at the same time, are responsible for significant morbidity and mortality. The chronic complications of diabetes are broadly divided into microvascular and macrovascular, with the former having much higher prevalence than the latter [6].

In Senegal, an estimated 400,000 people have diabetes, representing a prevalence of 3.4%, but only a small minority of the population is diagnosed. This prevalence varies according to geographical area, reaching around 10% in Saint-Louis, the region most affected by diabetes. [7]

Treating diabetics requires considerable resources and is not always within the reach of most of the African population. In Africa, 80% of the population relies on traditional medicine for primary healthcare. Many plant-based recipes are used in traditional African medicine to treat diabetes [8]. In addition to their standard therapies, many diabetes patients are known to take herbal medications with antidiabetic qualities. This can be advantageous or potentially harmful to the efficient management of their disease.

Despite the availability of many conventional prescription medications, numerous side effects, which are intolerable for many patients is on the rise. In recent years, the search for alternative therapeutic agents in the treatment of diabetes has been the focus of scientific research, as medicinal plants with diverse actions have been used traditionally for the control, management and/or treatment of DM in many parts of the world [9].

Many medicinal plants with hypoglycaemic properties have been used since ancient times to treat hyperglycaemia [10]. Nowadays, a large number of medicinal plants are believed to possess antidiabetic properties and have been utilised to manage diabetes [4,5,11]. For example, many herbal medicines possess antioxidant properties which could be beneficial for reducing oxidative stress, a key pathogenic factor of diabetes [12,13,14]. Several pharmaceutical agents effective in

reducing diabetic mortalities (e.g., 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors) have also been shown to have antioxidant activities [12]. Antioxidants including ascorbic acid have shown to have a prospect in the treatment of DM [15]

Africa is the world's leading region for undiagnosed diabetes, according to the WHO. Only 46% of people with diabetes on the African continent are aware of their condition, which increases the risk of serious illness and death. For this reason, we have set out to research medicinal plants with hypoglycaemic properties, which will provide practitioners with an overview of the scientific aspect of medicinal plants used clinically to treat diabetes.

2. Material and Methods

2.1. Study area

Mbour is a town in western Senegal, located on the Petite-Côte, about 80 km south of Dakar and bordering the seaside resort of Saly. It is the capital of the department of Mbour. The nearest towns are Saly Niakhiakhal, Saly-Portudal, Malikounda Sas, Malicoundangoukhoudji, Falokh, Sintiou Mbadane, Nianing and Warang.

Located between 14° 24' 42" north, 16° 57' 57" west and the altitude is 0m. Mbour covers an area of 1,607 km². Mbour has a semi-arid steppe climate, with an average annual temperature of 26.2°C and low rainfall of around 585 mm per year.

According to the 2023 report by the National Agency of statistics of demography (NASD), the department of Mbour has 215 facilities comprising: 01 hospital, 04 health center, 74 health posts and 136 health huts for an estimated population of 937,189 inhabitants. Fig. 1 shows the map of study area.



Fig.1. Map of study area

2.2. Study design

Surveys based on direct questioning about the uses of anti-diabetic plants were conducted over 3 months in two markets visited in the town of Mbour.

2.3. Study population

The target population consisted of 30 traditional healers interviewed without distinction as to sex, age, social status or religion.

2.4. Sampling

Using a questionnaire, surveys were carried out in the field with 30 traditional practitioners.

2.5. Data collection instruments

The questionnaire consists of two parts first part on the respondent, in which identification parameters are entered (sex, age, education, marital status, length of service, etc.) A second part which collects information about the plants used to treat diabetes (name of the plant, part used, method of preparation).

2.6. Data processing

To process the data collected, we used the method of calculating percentages or citation frequencies (CF), which is the ratio of the number of times the species was cited (n) to the total number of players (N).

$$CF = n/N$$

3. RESULTS

The main data collected in the localities of the town of Mbour were the socio-demographic characteristics of the people surveyed and information on the plants used for diabetics by traditional practitioners.

3.1. Distribution by locality

Most of the traditional healers, we met were in the central market, with a total of 25 out of 30, i.e. 83.33%. Fig. 2 shows the distribution according to locality.




Fig. 2. The distribution according to locality.

3.2. Distribution by gender

A total of 30 traditional practitioners were surveyed, including 22 men (72%) and 08 women (28%), giving a sex ratio of 0.36. Fig. 3 shows the distribution by sex.




Fig. 3. The distribution by sex.

3.3. Distribution according to marital status

The majority of the traditional healers were married with a rate of 90% (i.e. 27 out of 30). Fig. 4 shows the distribution according to marital status.

TRADITIONAL PRACTITIONERS




Fig. 4. The distribution according to marital status.

3.4. Age distribution

The most represented age group is [40-60] with 46.66%, followed by [20-40] and [60-80] with 26.66% each. The fig. 5 shows the age distribution.




Fig. 5. The age distribution.

3.5. Distribution by number of years in the business

86.66% of those surveyed had been in the trade for more than 5 years. The age group most represented was [10-15] with 33.33%, followed by [5-10] with 30%, then [15-20] with 16.66%, then [0-5] with 13.33% and finally [20-25] with 6.66%. Fig. 6 shows the Distribution by number of years in the business.




Fig. 6. The distribution by number of years in the business.

3.6. Distribution by level of education

Most of the traditional healers interviewed (56.6%, or 17 out of 30) had attended school, while 44.4% had not. Fig. 7 shows the distribution according to level of education.



Fig.7. The distribution according to the level of education.

3.7. Inventory of plants cited in the treatment of diabetes

At the end of the survey, 14 plants used to treat diabetes were cited. These plants are divided into 10 families, the most represented being the Euphorbiaceae, Combretaceae and Anacardiaceae. They will be translated into a table showing their scientific names, families, vernacular names, parts used and methods of use. Table 1 shows inventory of plants cited.

Table 1. The inventory of plants cited.

Scientific names	Families	Vernacular Names	Parts used	Modes of use
<i>Sclerocarya birrea</i> (Höchst)	Anacardiaceae	Beer (W) Arik (s) KoutenDiao (S)	Leaves, Barks	Infusion, Decoction
<i>Garcinia kola</i> (Heckel)	Clusiaceae	Bitikola (W)	Seeds	Maceration
<i>Anacardium occidentale</i> (Köhler)	Anacardiaceae	Darkassé (W) Darkassou (s) Finza (b)	Barks	Infusion
<i>Streptogyne rontogaea</i> (Hook. F.)	Poaceae	Fékh	Roots	Infusion
<i>Vahlia dichotoma</i> (Murray)	Vahliaceae	Mbélam (W)	Leaves	Décoction
<i>Combretum lecardii</i> (Engl. Et Diels)	Combretaceae	Ndadél (W) Piroriem (B)	Leaves	Infusion
<i>Chrozophora senegalensis</i> (A.Juss.)	Euphorbiaceae	Ndiamat (W) Mbélo (S)	Leaves	Infusion
<i>Boscia senegalensis</i> (Poir.)	Capparaceae	Ndiandam (W)	Leaves	Infusion
<i>Moringa oleifera</i> (Lam)	Moringaceae	Nebeday (W) Némédayo (S)	Leaves, Barks, Roots and Seeds	Decoction, Infusion
<i>Neocarya macrophylla</i> (F.White)	Chrysobalanaceae	Neew (W) Niamoui (P) Néoudi (P)	Leaves, Barks and Roots	Decoction, Infusion
<i>Chrysobalanus icaco</i> (L.)	Chrysobalanaceae	Radji (W) Moholo (P)	Leaves, Roots	Maceration
<i>Terminalia avicennioides</i> (Guill. Et Perr.)	Combretaceae	Reubreub (W) Bori billel (P)	Leaves, Barks and Roots	Infusion, Decoction
<i>Allium cepa</i> (L.)	Amaryllidaceae	Soblé (W)	Seeds	Macération
<i>Jatropha curcas</i> (L.)	Euphorbiaceae	Timitimi (W) Tabanano (S) Tabanani (s)	Leaves	Infusion

W : Wolof ; **P** : Pulaar ; **S** : Socé ; **s** : Sérère ; **B** : Balante and **b** : Bambara

3.8. Frequency of plant citations

Table 2 shows that the most frequently cited plant among the 14 was *Neocaryamacrophylla* with 21 citations (70%), followed by *Moringa oleifera* with 53.33%, *Terminaliaavicennioides* (46.66%) and *Sclerocaryabirrea* (40%).

Table 2. Frequency of plant citations

ScientificsNames	Number of citation	Percentage of citation
<i>Sclerocaryabirrea</i> (Höchst, 1844)	12	40%
<i>Garcinia kola</i> (Heckel, 1883)	2	6.66%
<i>Anacardium occidentale</i> (Köhler)	3	10%
<i>Streptogynerontogaea</i> (Hook. F.)	1	3.33%
<i>Vahliadichotoma</i> (Murray)	2	6.66%
<i>Combretumlecardii</i> (Engl. Et Diels)	1	3.33%
<i>Chrozophasenegalensis</i> (A.Juss.)	1	3.33%
<i>Bosciasenegalensis</i> (Poir., 1819)	2	6.66%
<i>Moringaoleifera</i> (Lam, 1785)	16	53.33%
<i>Neocaryamacrophylla</i> (F.White, 1976)	21	70%
<i>Chrysobalanusicaco</i> (L., 1753)	2	6.66%
<i>Terminaliaavicennioides</i> (Guill. Et Perr.)	14	46.66%
<i>Alliumcepa</i> (L., 1753)	1	3.33%
<i>Jatrophacurcas</i> (L., 1753)	1	3.33%

3.9. Frequency of cited plant parts used

The leaf of the tree is the part most frequently cited by actors, with a frequency of 44.2%, followed by bark (31%), roots (20.9%) and seeds (3.9%). Fig. 8 shows the frequency with which the parts used on the plant are cited.




Fig. 8. The frequency with which the parts used on the plant are cited.

3.10. Frequency of cited methods in use

Use after infusion is cited most frequently (61.2%), followed by decoction (31%) and maceration (7.8%). Fig. 9 shows the frequency with which methods of use are cited.




Fig. 9. The frequency with which methods of use are cited.

4. DISCUSSION

Our Survey was carried out in the Community of Mbour, with the aim of listing plants used in the treatment of diabetes. This study was initiated with 30 traditional practitioners.

During our study, almost all the traditional healers we met told us that they had inherited their knowledge. The study revealed that 72% were men and 28% were women. This large predominance can be explained by the fact that, it is rare for women to inherit traditional knowledge.

All age groups were represented, with a minimum age of 22 and a maximum age of 78. The most represented age group was [40-60] with 46.66%, followed by [20-40] and [60-80] with 26.66% each. The low rate of representation in the over-60s may be explained by the old age and activity of older subjects.

The level of education among participants is relatively low, with only 56.66% having attended school. We also noted that 70.58% of those attending school had left at elementary level.

The majority of our participants are married, representing 90% (27 out of 30), compared with 10% who are single. This may be due to the fact that, they earn for their living.

The longest periods of practice were [5-10] and [10-15], with 30 and 33.33% respectively. In other words, 93.33% of actors have been in practice for [0-20] years, and 6.66% for more than 20 years. This can be explained by a more marked start to practice after the age of 40 and retirement in old age.

With regard to the anti-diabetic plants cited by traditional practitioners, this work enabled us to identify 14 species divided into 14 genera in 10 families, the most common of which are: Anacardiaceae, Combretaceae, Chrysobalanaceae and Eupobiaceae. This low number of species can be explained by the fact that the people surveyed live in the same locality and cite the same plants for the treatment of diabetes. The number of citations for each plant shows that the species most frequently cited by the in this study to treat diabetes are: *Neocaryamacrophylla*, *Moringa oleifera*, *Terminaliaavicennioides* and *Sclerocaryabirrea*.

The anti-diabetic effect of *Neocaryamacrophylla* is very well known, even from Diatta's work in 2022 on the anti-diabetic properties of *Neocaryamacrophylla* [16]. These studies revealed that the flavonoids contained in *N. macrophylla* leaves prevent diabetic cataracts by inhibiting aldose reductase in the lens. In fact, myricetin has hypoglycaemic effects in diabetic animals. This confirms its anti-diabetic effect.

As regards *Moringa oleifera*, studies according to Gupta R *et al.*, 2012 investigated the anti-diabetic and antioxidant effects of methanolic extracts of *M. oleifera* pods in streptozotocin-induced diabetic albino rats. These results showed that the progression of diabetes was significantly reduced after treatment with *M. oleifera*, confirming that the latter has anti-diabetic properties [17].

The studies of Yahaya SF *et al.*, 2019 on the hypoglycaemic effects of *Terminaliaavicennioides*. This study evaluated *Terminaliaavicennioides* stem bark extracts for their effect on alloxan-induced diabetes mellitus in male wistar rats

[18]. These results obtained in the course of this work demonstrate the hypoglycaemic effects of *T. avicennioides* cited in our survey.

Previous work by Ojewole JAO, 2004 reported a hypoglycaemic action of the aqueous extract of *Sclerocaryabirrea* stem bark, studied in rats. These results confirm our work, that *Sclerocaryabirrea* contains anti-diabetic properties [19].

The studies according to Diouf P, 2016 on the contribution to the study of *Anacardium occidentale* in the city of Thiès, shows that this plant is used in the treatment of diabetes [20].

There are studies showing the antihyperglycaemic property of certain species of Chrysobalanaceae, such as *Chrysobalanus icaco*, confirming its use in traditional medicine [21].

Niang K (2022) in his studies on the antioxidant activity of the hydroethanolic extract of *Boscia senegalensis* leaves showed an anti-hyperglycaemic activity of *Boscia senegalensis* on albinorabbits [22].

Studies by Pradeep SR *et al.*, 2017 indicated *Allium cepa* amplified the amelioration of diabetic hyperglycaemia and related metabolic abnormalities in experimental rats [23]. The present study evaluates the enhanced benefits of onion seed addition on oxidative stress in diabetic rats.

5. CONCLUSION

At the end of this study, a repertory of 14 plant species belonging to 10 families was established. The most commonly used plant parts were leaves (44.20%), followed by barks (31%), roots (20.90%) and seeds (3.90%). The most common preparation techniques are infusion (61.20%), decoction (31%) and maceration (7.80%).

In the course of our work, the plants most frequently cited by traditional practitioners were the following: *Neocaryamacrophylla*, *Moringa oleifera*, *Terminalia avicennioides* and *Sclerocaryabirrea*. And the least cited plants are: *Garcinia kola*, *Anacardium occidentale*, *Vahlia dichotoma*, *Streptogynegerontogaea*, *Combretum lecardii*, *Chrozophora senegalensis*, *Boscia senegalensis*, *Chrysobalanus icaco*, *Allium cepa* and *Jatropha curcas*. The scientific literature has shown that these plants have anti-diabetic properties and maybe an alternative treatment for diabetes.

Phytochemical, toxicological and pharmacological studies need to be carried out to help validate their traditional use and find new plants with anti-diabetic potential.

6. DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

7. REFERENCES

1. OMS. (2023). Diabetes [Internet]. Disponible sur:<https://www.who.int/news-room/fact-sheets/detail/diabetes>
2. Riddle MC. Standards of medical care in diabetes. American Diabetes Association. 2019 ; 42:S1-193.PMID: 30559228. DOI: [10.2337/dc19-S002](https://doi.org/10.2337/dc19-S002)
3. Papatheodorou K, Banach M, Bekiari E, Rizzo M, Edmond M. Complications of diabetes 2017. *J Diabetes Res*;2018:3086167. doi: 10.1155/2018/3086167
4. Qi LW, Liu EH, Chu C, Peng YB, Cai HX, Li P. Anti-diabetic agents from natural products—an update from 2004 to 2009. *Curr Top Med Chem*. 2010;10(4):434–57.PMID: 20180758. DOI: [10.2174/156802610790980620](https://doi.org/10.2174/156802610790980620)
5. Ghorbani A. Clinical and experimental studies on polyherbal formulations for diabetes: current status and future prospective. *J Integr Med*. 2014;12(4):336–45.PMID: 25074883. DOI: [10.1016/S2095-4964\(14\)60031-5](https://doi.org/10.1016/S2095-4964(14)60031-5)
6. Samad A, Shams MS, Ullah Z, Wais M, Nazish I, Sultana Y, Aqil M. Status of herbal medicines in the treatment of diabetes: a review. *Curr Diabetes Rev*. 2009;5(2):102–11.PMID: 19442095.DOI: [10.2174/157339909788166837](https://doi.org/10.2174/157339909788166837)
7. Zimmer P, Magliano DJ, Herman WH, Shaw J. Diabetes a 21st century challenge. *The Lancet Diabetes Endocrinology*. 2014 ;2(1):56-64.PMID: 24622669.DOI: [10.1016/S2213-8587\(13\)70112-8](https://doi.org/10.1016/S2213-8587(13)70112-8)
8. Ogurtsova K, da Rocha Fernandes JD, Huang Y, Linnenkamp U, Guariguata L, Cho NH, Cavan D, Shaw JE, Makaroff LE. IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res Clin Pract*. 2017 Jun;128:40-50. PMID: 28437734.doi: 10.1016/j.diabres.2017.03.024.
9. Deshpande A. D., Harris-Hayes M., Schootman M. Epidemiology of diabetes and diabetes-related complications. *Physical Therapy*. 2008;88(11):1254–1264. doi: 10.2522/ptj.20080020.
10. LVL Medical. (2023). Our committed collaborators, beyond borders | LVL Medical [Internet]. Available at: <https://www.lvlmedical.com/nos-collaborateurs-engages-au-dela-des-frontieres>
11. Dia L. (2006). Antidiabetic phytotherapy: Bibliographic synthesis on antidiabetic research. Thesis pharma. PhD, Cheikh Anta Diop University of Dakar. 129p.
12. Qi LW, Liu EH, Chu C, Peng YB, Cai HX, Li P. Anti-diabetic agents from natural products—an update from 2004 to 2009. *Curr Top Med Chem*. 2010;10(4):434-57. doi: 10.2174/156802610790980620. PMID: 20180758.
13. Khiari M, Boukhatia F, Zahra H, Mizouri R, Zribi S, Temessek A, & Mami F B. Ketotic decompensation between type 1 and type 2 diabetes. In *Annales d'Endocrinologie*. 2018; Vol. 79, No. 4: p 486. Elsevier Masson. <https://doi.org/10.1016/j.ando.2018.06.962>

14. Sowndharajan K, Kang S. Free radical scavenging activity from different extracts of leaves of *Bauhinia vahlii* Wight & Arn. *Saudi J Biol Sci.* 2013;20(4):319-25. PMID: 24235867 PMCID: PMC3824142 DOI: 10.1016/j.sjbs.2012.12.005
15. Ramachandran S, Rajasekaran A, Manisenthilkumar K. Investigation of hypoglycemic, hypolipidemic and antioxidant activities of aqueous extract of *Terminalia paniculata* bark in diabetic rats. *Asian Pac J Trop Biomed.* 2012;2(4):262. PMID: 23569911 PMCID: PMC3609298 DOI: 10.1016/S2221-1691(12)60020-3
16. Diara DIATTA. Pharmacochemistry, structural analysis and biological activities of metabolites of *Raphionacmevignei* and *Neocaryamacrophylla*. Unique Doctoral Thesis, Assane Seck University of Ziguinchor (Senegal); 2022: 224 pp.
17. Gupta R, Mathur M, Bajaj VK, Katariya P, Yadav S, Kamal R, et al. Evaluation of antidiabetic and antioxidant activity of *Moringa oleifera* in experimental diabetes. *Journal of Diabetes.* 2012;4(2):164-71. PMID: 22103446 DOI: 10.1111/j.1753-0407.2011.00173.x
18. Yahaya SF, Suleiman MM, Mohammed A, Ibrahim NDG. Antidiabetic potentials of stem-bark extracts of *Terminalia avicennioides* on alloxan-induced diabetic rats. *Sokoto Journal of Veterinary Sciences.* 26 Nov 2019;17(2):33-44. DOI: 10.4314/sokjvs.v17i2.5
19. Ojewole JAO. Evaluation of the analgesic, anti-inflammatory and antidiabetic properties of *Sclerocarya birrea* (A. Rich.) Hochst. stem-bark aqueous extract in mice and rats. *Phytotherapy Research.* 2004;18(8):601-8. doi: 10.1002/ptr.1503. PMID: 15476310
20. Diouf P. (2016). Contribution to the study of *Anacardium occidentale*. L. Ethnobotanical survey of traditional practitioners in the city of Thiès. Thesis Doctorate in pharmacy, Cheikh Anta Diop University of Dakar. 116 pp.
21. Feitosa EA, Xavier HS, Randau KP. *Chrysobalanaceae*: traditional uses, phytochemistry and pharmacology. *Rev arm farmacogn.* Oct 2012;22:1181-6. <https://doi.org/10.1590/S0102-695X2012005000080>
22. Niang MK. (2022). Studies on the antioxidant activity of the hydroethanolic extract of the leaves of *Boscia senegalensis* (PERS) LAM (CAPPARACEAE) and fractions by the ABTS method. Pharma thesis. PhD, Cheikh Anta Diop University of Dakar. 98p.
23. Pradeep SR, Srinivasan K. Amelioration of oxidative stress by dietary fenugreek (*Trigonella foenum-graecum* L.) seeds is potentiated by onion

(*Allium cepa* L.) in streptozotocin-induced diabetic rats. *Appl Physiol Nutr Metab*.
Aug 2017;42(8):816-28. <https://doi.org/10.1139/apnm-2016-0592>

UNDER PEER REVIEW