

## **“A Review on Importance Nutritional Value and Uses of Water Chestnut (*Trapanatans var. bispinosaRoxb.*)”**

### **ABSTRACT**

Water chestnut (*Trapanatansvar. bispinosaRoxb.*), an aquatic perennial plant. This research aims to elucidate the health benefits of water chestnut, focusing on its nutritional composition, medicinal properties, and potential roles in promoting overall wellness. Water chestnuts are a rich source of essential nutrients, including vitamins B6, C, and E, as well as minerals such as potassium, phosphorus, and magnesium. Antioxidants play a critical role in reducing inflammation, which is linked to numerous health issues, including heart disease and arthritis. The anti-inflammatory properties of water chestnuts may help alleviate symptoms associated with these conditions, improving overall quality of life. The potassium content helps regulate blood pressure, while dietary fiber aids in cholesterol management. These factors contribute to a healthier cardiovascular system, potentially reducing the risk of heart disease and stroke. Water chestnuts are also renowned for their hydrating properties, as they contain a high water content, making them an excellent choice for maintaining hydration, especially in hot climates or during physical activities.

**Keywords:** Water chestnuts, nutritional value, uses, health benefit

### **1. INTRODUCTION**

Water chestnut (*Trapanatans var. bispinosaRoxb.*) is an important aquatic fruit crop in India, and it belongs to the 'Trapaceae' family. Water chestnut, also known as Singhara in Hindi, is said to have originated in India, China, and Eurasia. It is a free-floating annual aquatic plant that grows in shallow fields, ponds, and swampy areas. It prefers nutrient-rich water with a pH range of 6.7 to 8.2 (Anonymous, 2002; Bhatiwalet al., 2012). Water chestnuts are widely grown in India, particularly in Madhya Pradesh, Uttar Pradesh, Bihar, Odisha, West Bengal, Jharkhand, Karnataka, and Jammu & Kashmir. The plant is commercially farmed in tropical places around the world, including Pakistan, Sri Lanka, Indonesia, and Africa. It's common in Indonesia, Southeast Asia, southern China, and the eutrophic waterways of Japan, Italy, and tropical America. It has also naturalized in several places of the Eastern United States, where it thrives in tropical and subtropical conditions (Takano and Kadono, 2005). Fresh water chestnuts have the highest water content (80%) (Puste, 2004), starch (52%), protein (1.87%), and total soluble solids (TSS) (7–8%) (Singh et al., 2010). They are also high in fiber, vitamin B, calcium (Ca), potassium (K), iron (Fe), and zinc (Adkar et al. 2014) and (Alfasaneet al. 2011), fresh chestnuts contain 4.40% protein and 22.3% carbohydrate, respectively. Water chestnuts are also a good source of crude fiber, with the green variety having 2.13% and the red type having 2.27% (Faruk et al., 2012). Water chestnut flour contains a considerable amount of potassium and fiber. Each cup of water chestnut slices contains around 130 calories. In India, water chestnut flour is consumed during fasting days (Singh, 2017). Water chestnuts' protein and carbs include a significant quantity of flavonoid and antioxidants, making them potential sources of nutrition (Mann et al., 2012). Water

chestnuts have a strong antiulcer effect, considerably lowering ulcer indices in both pyloric and aspirin-induced ulceration instances (Kar *et al.*, 2010). Furthermore, both red and green water chestnuts have antibacterial properties (Razvyet *et al.*, 2011).

## **2. BOTANICAL DESCRIPTION**

A floating aquatic plant with two leaf types is called a water chestnut. One grows along the length of the stem, split like a feather and submerged. Undivided floating leaves that form a rosette close to the water's surface make up the other (Jana, 2016). The floating leaves are fan-shaped, serrated on the edges, and rhomboid. Their diameter ranges from 2 to 6.5 cm, with their width being more than their length (Zhu, 2016). According to (Chandana *et al.*, 2013), the undersides of the leaves are reddish-purple, while the uppers are dark green. The plant's spongy, buoyant, cord-like stems can grow up to 16 feet in length, though most of them are usually between 6 and 8 feet. According to (Adkar *et al.*, 2014), a channel of branching roots anchors the stems to the water's bed. White flowers have four petals that are around 8 mm long. It starts to bloom in August and is situated in the middle of the rosette (Singh *et al.*, 2018; Jana, 2016). The fruit features a bony or woody nut that is 2.5 to 5 cm in diameter and has two or four robust horns or spines. While the fruit of *Trapa bicorn* resembles that of *Trapanatans*, it bears two spines instead of four (Jana, 2016; Adkar *et al.*, 2014).

### **2.1 Habitat**

Worldwide, under full sun and in freshwaters rich in nutrients but low in energy, *Trapanatans* can be found (Hummel and Kiviat, 2004). It tends to avoid waters that are high in calcium (PFAF, 2000) and is typically found in waters with an alkalinity of 12 to 128 mg/L of calcium carbonate (O'Neill, 2006). The average depth of water that *Trapanatans* dwell has been reported in a variety of ways. According to some sources (Van Driesche, 2002), *T. natans* can grow in water as deep as five meters, according to other sources (Hummel and Kiviat, 2004), and yet other sources (PFAF, 2000) state that the maximum depth is 0.6 meters. The species is most common in water that is about two meters deep and on soft terrain, according to (Hummel and Kiviat, 2004).

### **2.2 Flowering and Fruiting**

Following the start of the rainy season, which is usually in July, the creepers begin their great time of growth, and by the end of July, flower bud initiation occurs. After being transplanted, the plant takes approximately one month to begin flowering. The plants don't stop blossoming once they get going, even as they become older and eventually die (Khanna, 1963). The process of flower opening and anther dehiscence practically happen at the same time, which guarantees self-pollination (Khanna, 1963). There are roughly 30 to 35 leaves on each stalk, and every leaf has a bud that appears from its axil and grows over the course of four to six days. While fruit development occurs beneath the water's surface, flower opening and fertilization occur outside of it. It is possible for a bloom to turn into a fruit within 10 to 15 days of pollination (Chakor, 1974). But after 21 days of anthesis, (Purewal and Kong, 1958) that the fruit was ready to be harvested. The observations show that the time between anthesis and fruit maturity varies, which could be caused by variations in the climate or in the varieties.



time,  
35  
single  
over  
occur  
turn  
days  
noted



**Image1- Flowering**

**Image 2- Fruiting**

### 3. NUTRITIONAL VALUE OF WATER CHESTNUT

*Trapabispinosa* fruits have a biochemical makeup that has been examined. The results indicate that the fruit is suitable for inclusion in a human diet and may include significant amounts of protein, carbohydrates, and minerals (Alfasaneet *al.*,2011). In terms of nutrients, water chestnuts have the following composition: 62.5 percent moisture, 1.04 percent ash, 2.13% crude fiber, 0.92% total soluble sugar, 0.33% reducing sugar, 0.59% nonreducing sugar, 8.7% starch, and 0.84% fat. Beta-carotene was 60 micrograms, vitamin C was 1.1 mg,

water soluble protein was 0.275 mg, and total phenol was 0.5 mg in one hundred grams of green variety. Iron, copper, manganese, and zinc were found in 200, 430, 90, and 600 parts per milligram, respectively, while potassium was found in 5.22%, sodium in 0.64%, calcium in 0.25%, phosphorus in 6.77%, and sulphur in 0.38% of the green variety's mineral components. Moisture (62.7%), ash (1.30%), crude fiber (2.27%), total soluble sugar (0.90%), reducing sugar (0.30%), nonreducing sugar (0.60%), starch (8.2%), and fat (0.83%) were all present in the red variant. Within 100 grams, the red type had 0.251 mg of water-soluble protein, 92 microg of beta-carotene, 0.9 mg of vitamin C, and 0.60 mg of total phenol. Potassium (5.32%), sodium (0.59%), calcium (0.26%), phosphorus (6.77%), sulphur (0.32%), iron (200 ppm), copper (450 ppm), manganese (110 ppm), and zinc (650 ppm) were all present in the red variety. Both types frequently contained the free amino acids glutamic acid, tryptophan, tyrosine, alanine, lysine, and leucine. Furthermore, cysteine, arginine, and proline, as well as glutamine and asparagine, were present in the green and red variants, respectively. As a result, the current study clarifies the nutritional differences between the two types of water chestnuts and raises the possibility that they are essential for human nutrition (Faruk *et al.*, 2012).

**List 1 :Nutrient profile of 100g of Water Chestnut (IFCT 2017).**

<b>Nutrient</b>	<b>Amount Per 100g</b>
Protein	0.86 g
Carbohydrates	21.46 g
Total Fat	0.37 g
Energy	95.6 kcal
Calcium	37.15 mg
Phosphorous	62.83 mg
Iron	0.77 mg
Vitamin C	5.26 mg
Total Fiber	3.02 g

#### **4. MEDICINAL VALUE**

Medicinal value of the whole herb and fruit have been recognized in folklore medicine as a cure for various diseases (Rani *et al.*, 2016). The whole herb has been reported for hepato - protective activity, antimicrobial activity, antibacterial activity (Razvy *et al.*,2011), antitumor activity, antioxidant activity and free radical scavenging activity (Chandana *et al.*, 2013; Das

*et al.*, 2011; Jha, 1999 & Jana *et al.*, 2020). The fruits have been used as an anti-inflammatory, anti-diarrhea, intestinal astringent, aphrodisiac antileprotic agent and in urinary discharges, fractures, bronchitis, and anemia. The fruits of *T. bispinosa*. have been identified as the Ayurvedic drug Shurangataka. It is also said to have cancer-preventing properties. Stem juice is used in ophthalmic preparations. The dimension of its starch particles ranges between 15 and 39  $\mu$  which is very near to those of potato's starch particles.

#### **4.1 Reduce the risk of heart disease**

With modernization, the risk of chronic diseases such as heart disease has increased, making it one of the leading causes of death worldwide. Heart disorders are caused by a variety of causes, including excessive blood pressure, high blood cholesterol levels, and high blood triglycerides. One of the most important aspects of the water chestnut as a fruit is that it aids in blood pressure regulation due to its high potassium content. Research suggests that lowering the risk of stroke and blood pressure can lower the likelihood of developing heart disease. A study found that those with high blood pressure who drank more potassium had lower systolic and diastolic blood pressure by 3.49 mmHg and 1.96 mmHg, respectively, and had a lower risk of heart stroke. Another review of 11 research found that persons who consumed more potassium had a 21% decreased risk of heart attack and heart disease (Rani *et al.*, 2016).

#### **4.2 Anti-oxidant property of water chestnut**

Water chestnuts contain a high level of total phenol, flavones, and flavonoids. The water chestnut fruit seed extract was found to contain carbohydrates, saponins, phytosterols, fixed oils, and fat, but the fruit pericarp extract contained tannins, flavonoids and glycosides, alkaloids, saponins, steroids, and phenol compounds (Hussain *et al.*, 2018).

#### **4.3 Anti-diabetic property of water chestnut**

Water Chestnut fruit peel extract shown anti-diabetic effects in streptozotocin-induced diabetic rats. One of the most noteworthy advantages is that the ethanolic extract of Water Chestnut roots and fractions can slow the progression of hyperglycaemia while preventing hypoglycaemia (Das *et al.*, 2011). The presence of ferulic acid and caffeic acid in Water Chestnut explains its anti-hyperglycaemic properties. Ferulic acid is known to minimize oxidative stress and hyperglycemic response, although caffeic acid has a generative effect on Langerhans islets as a result of decreased expression of glucose transporter 2 in the liver, which lowers hyperglycaemia (Corovicet *et al.*, 2021).

#### **4.4 Anti-ulcer activity of water chestnut**

The study used Wistar rats to test the antiulcer capabilities of *Trapabispinosa* fruits. The antiulcer activity of 50% ethanolic extract was evaluated at two dosage levels using the aspirin plus pyloric ligation and pyloric ligation plus pyloric ligation models. The test extract had significant antiulcer effect, possibly due to increased carbs and alterations in the stomach's mucosal barrier. The research suggests that *Trapabispinosa* fruit extract (ethanolic) may have antiulcer potential (Kar *et al.*, 2010).

#### **4.5 Anti-microbial property of water chestnut**

Water Chestnut leaf extract has the potential to be used in microbial control because it is a rich source of biologically active chemicals. Antimicrobial activity was specifically demonstrated by the bacteria under examination. *Pseudomonas aeruginosa* and other gram-negative bacteria were especially susceptible to this effect (Radojevic *et al.*, 2016). The anti-quorum sensing activity of Water Chestnut leaf extracts inhibits the manufacture of many virulence factors required for the progression of infection in *P. aeruginosa*, a pathogen (Aleksic *et al.*, 2018). As a reference, kanamycin was utilized by (Razvyet *et al.*, 2011). To evaluate the antibacterial activity of fruit extract from two varieties of water chestnuts: green and red. The disc diffusion method was employed. The green variety of water chestnuts exhibited the highest antibacterial activity (12 mm) against both *Shigella sonnei* and *Staphylococcus aureus*, whereas the extract from the red variety exhibited a strong antibacterial potential (31 mm) against *Bacillus subtilis*.

#### **4.6 Anti-inflammatory property of water chestnut**

Water chestnut extracts contained phenolic compounds such as gallic acid. Gallic acid appears to have antioxidant, anti-inflammatory, antiviral, and anti-cancer activities (Chuang *et al.*, 2010). Water chestnut extract contains gallic acid, which has a high anti-inflammatory effect and has been utilized in the treatment of skin inflammatory conditions (Kim *et al.*, 2015). Trapabispnosahydroalcoholic extract (500 mg/kg peroxidase) reduced fluorescence product and lipid peroxidation in the cerebral cortex of female albino mice while restored glutathione peroxidase and catalase function (Ambikaret *et al.*, 2010). Using pyloric ligation and aspirin plus pyloric ligation models, (Kar *et al.*, 2010), Examined the antiulcer impact of a 50% ethanolic extract of *Trapabispinosa* (Trapaceae) fruits on Wistar rats and discovered that the plant possesses potential antiulcer activity. When rats were exposed to sheep red blood cells (SRBC) as an antigen, (Patelet *et al.*, 2010), examined the humoral immune response, cell-mediated delayed-type hypersensitivity reaction (DTH), and the percentage change in neutrophil count. They found that an aqueous preparation of *T. bispinosa* fruit increased the animal's humoral and cellular responses.

## **5. USES OF WATERCHESTNUT**

### **5.1 AS A FOOD**

In the Indian subcontinent, water chestnuts, are used as a substitute for grains during fasting. In the case of wheat flour (WF), which is associated with a problem with the digestion of gluten (a wheat protein), water chestnuts may be a preferable option. (Gulet *et al.*, 2014 and Miret *et al.*, 2015). In India, on Navratri and other fasting days, water chestnut cookies are served as a specialty meal. Water chestnut flour (WCF) holds great potential for the creation of gluten-free bread goods due to its high starch content and lack of gluten. Its varied nutritious qualities and high fiber content further validate its use in the dishes. There is a rise in demand for water chestnut flour cookies because of the higher spreading ratio of water chestnut cookies when compared to WF cookies.



**Image 3,4 Water Chestnut Kernel**

## **5.2 VALUE ADDED PRODUCTS**

Nutritious and gastronomic benefits can be obtained from value-added products made from *Trapanatans*, also called water chestnut. The plant's edible parts, the seeds or corms inside the spiky fruit, are usually processed into these goods. This list includes a few Water Chestnut items with extra value.

### **a. Production of Gluten-Free Products**

Chestnut flour is gluten-free and has additional health benefits, therefore it can be used by patients with celiac disease. Some scientists investigated the possibility of using chestnut flour instead of cow milk for making kid-friendly soups and desserts because lactose in cow milk might trigger allergic reactions in young children. Additionally, flakes, pasta, bread, baby formula, and milky puddings can be made with chestnut flour, per (Crow, *et al.*, 2000 and Parekh *et al.*, 2007). High in dietary fiber (4–10%), high in sugar (20–32%), high in starch (50–60%), high in protein (4-5%), high in essential amino acids (4-6%), and low in fat (2-4%) are all characteristics of chestnut flour. It also has high levels of magnesium, calcium, potassium, and the vitamins B, C, and E. It is thought that utilizing chestnut flour will help because most gluten-free products currently lack vitamin B, iron, and fiber. Because chestnut flour has so many nutritional benefits, it is also used to make gluten-free bread (Gani *et al.*, 2010).

### **b. Water Chestnut Flour**

Many researchers have previously investigated the physico-chemical characteristics of water chestnut flour and starch (Shafi *et al.*, 2016; Gani *et al.*, 2010). Water chestnut flour hasn't been thoroughly studied in terms of its functional, viscosity, thermal, and structural characteristics, which could indicate why it's a good fit for developing new products. Through a variety of biochemical interactions, starch interacts with other main and minor ingredients during food processing, determining the final product's overall desirability.



**Image 5,6- Water Chestnut Flour**

**c. Water Chestnut Snacks**

To make crispy, wholesome snacks, water chestnuts can be cut and dried. To improve the flavor of these foods, seasoning or flavoring is frequently added. Crisp and delicious, they're a healthier substitute for regular potato chips.

**d. Canned or Preserved Water Chestnuts**

Water chestnuts with additional value are frequently canned or preserved. The water chestnuts are cut, peeled, and frequently kept in a light syrup or water. They give a crisp texture and a mild, somewhat sweet flavor to salads, stir-fries, and other foods, making them handy to use.

**e. Water Chestnut Chips**

Thinly sliced water chestnuts that have been deep-fried or dried to produce crispy chips are called water chestnut chips. Once seasoned, these chips can be eaten as a snack or added to salads and appetizers as a garnish.

**f. Water Chestnut Syrup or Concentrate**

The juice or essence of water chestnuts is extracted to make water chestnut syrup or concentrate. It adds a distinct flavor to drinks, sweets, and mixed drinks when used as a sweetener.

**g. Water Chestnut Paste**

Water chestnut flesh is used to make a purée known as water chestnut paste. Particularly in Asian cuisines, it serves as a foundation for a variety of sweets and desserts. Dumplings, buns, and pastry fillings can be made with it by flavoring and sweetening (Mondal *et al.*, 2024).

## **6. DISCUSSION AND CONCLUSION**

In conclusion, water chestnut emerges as a highly nutritious food with multiple health benefits. Its rich nutritional profile, combined with antioxidant, anti-inflammatory, and hydrating properties, positions it as a functional food that can contribute to disease prevention and overall health enhancement. As awareness of these benefits grows, further research is warranted to explore the mechanisms behind their health effects and to promote water chestnuts as a valuable component of a balanced diet. This exploration may lead to broader recognition and incorporation of water chestnuts in health-conscious culinary practices.

#### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

The authors hereby state that no generative A.I. tools, such as Large Language Models (COPILOT, ChatGPT, etc.) or text-to-image generators, were utilized during creation and editing of this work.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist

#### **REFERENCES**

1. Adkar, P., Dongare, A., Ambavade, S., & Bhaskar, V. H. (2014). *TrapabispinosaRoxb.*: a review on nutritional and pharmacological aspects. *Advances in Pharmacological and Pharmaceutical Sciences*, 2014(1), 959830.
2. Aleksic, I., Ristivojevic, P., Pavic, A., Radojević, I., Čomić, L. R., Vasiljevic, B., & Senerovic, L. (2018). Anti-quorum sensing activity, toxicity in zebrafish (*Danio rerio*) embryos and phytochemical characterization of *Trapanatans* leaf extracts. *Journal of ethnopharmacology*, 222, 148-158.
3. Alfasane, M. A., Khondker, M., & Rahman, M. M. (2011). Biochemical composition of the fruits of water chestnut (*TrapabispinosaRoxb.*). *Dhaka University Journal of Biological Sciences*, 20(1), 95-98.
4. Ambikar, D. B., Harle, U. N., Khandare, R. A., Bore, V. V., & Vyawahare, N. S. (2010). Neuroprotective effect of hydroalcoholic extract of dried fruits of *TrapabispinosaRoxb* on lipofuscinogenesis and fluorescence product in brain of D-galactose induced ageing accelerated mice.
5. Anonymous (2002). Addressing water chestnut in CNY. Oneida Lake and Watershed Management Plan Project News (Central New York Regional Planning and Development Board, Syracuse, NY) (fall):3-4.
6. Bhatiwala, Shalabh., Jain, A. and Chaudhary, J. (2012). *Trapanatans* L. (water chestnut): an overview. *Int. J. Res. Pharm.; Sci.*3: 31-32.
7. Chakor, LS. (1974). *Indian Far. Dig*, 7:35.
8. Chandana, M., Rupa, M. and Chakraborty, G. S. (2013). A review on potential of plants under *Trapa* species. *International Journal of Research in Pharmacy and chemistry*, 3 (2): 502 - 508.
9. Chuang, C. Y., Liu, H. C., Wu, L. C., Chen, C. Y., Chang, J. T., & Hsu, S. L. (2010). Gallic acid induces apoptosis of lung fibroblasts via a reactive oxygen species-

- dependent ataxia telangiectasia mutated-p53 activation pathway. *Journal of agricultural and food chemistry*, 58(5), 2943-2951.
10. Corovic, R. C., Bradic, J., Tomovic, M., Dabanovic, V., Jakovljevic, V., Zarkovic, G., & Rogac, Z. (2021). Chemical Composition and Biological Activity of *Trapa Natans* L. *Experimental and Applied Biomedical Research (EABR)*.
  11. Crow, G. E., & Hellquist, C. B. (2000). Aquatic and Wetland Plants of Northeastern North America: Angiosperms: Monocotyledons (Volume 2).
  12. Das, P. K., Bhattacharya, S., Pandey, J. N. and Biswas, M. (2011). Antidiabetic activity of *Trapanatans* fruit peel extract against streptozotocin induced diabetic rats. *Global Journal of Pharmacology*, 5 (3): 186 - 190.
  13. Faruk, M.O., Amin, M.Z., Sana, N.K., Saha, R.K. and Biswas, R.K. (2012). Biochemical analysis of two varieties of water chestnuts (*Trapaspp*). *Pakistan Journal of Biological Sciences.*; 15 (21), 1019-1026.
  14. Gani, A., Haq, S. S., Masoodi, F. A., Broadway, A. A., & Gani, A. (2010). Physico-chemical, morphological and pasting properties of starches extracted from water chestnuts (*Trapanatans*) from three lakes of Kashmir, India. *Brazilian Archives of Biology and Technology*, 53, 731-740.
  15. Gul, K., Riar, C. S., Bala, A., & Sibian, M. S. (2014). Effect of ionic gums and dry heating on physicochemical, morphological, thermal and pasting properties of water chestnut starch. *LWT-Food Science and Technology*, 59(1), 348-355.
  16. Hummel, M., Kiviat, E. (2004). Review of world literature on water chestnut with implications for management in North America. *Journal of Aquatic Plant Management*, 42:17-28.
  17. Hussain, T., Subaiea, G. M., & Firdous, H. (2018). Hepatoprotective evaluation of *Trapanatans* against drug-induced hepatotoxicity of antitubercular agents in rats. *Pharmacognosy magazine*, 14(54), 180.
  18. Jana, B. R. (2020) Agronomic Management of Water chestnut (*Trapanatans* L.): A Review. *International Journal of Current Microbiology and Applied Sciences*, 9 (8): 2773 - 2777.
  19. Jana, H. (2016). Water Caltrop: A potential crop of waterbodies. *Rastriya Krishi*, 11(2), 7-11.
  20. Jha, Vidyanath (1999). Nutritional Evaluation of *Trapanatans* L. var. *Bispinosa* Roxb. (Singhara) and Scope of Raising the Crop Potential under Integrated Aquaculture. *Journal of Freshwater Biology*, 11 (1 - 2): 11 - 17.
  21. Kar, D., Maharana, L., Si, S. C., Kar, M. K., & Sasmal, D. (2010). Antiulcer activity of ethanolic extract of fruit of *Trapabispinosa* Roxb. in animals. *Der Pharmacia Lettre*, 2(2), 190-197.
  22. Khanna, A.N. (1963). *Horticultural Advance*, VII 24-37.
  23. Kim, B., Kim, J. E., Choi, B. K., & Kim, H. S. (2015). Anti-inflammatory effects of water chestnut extract on cytokine responses via nuclear factor- $\kappa$ B-signaling pathway. *Biomolecules & Therapeutics*, 23(1), 90.
  24. Mann, S., Gupta, D., Gupta, V. and Gupta, R. (2012). Evaluation of nutritional, phytochemical, and antioxidant potential of *Trapabispinosa* Roxb. <http://www.ijppsjournal.com/Vol-4Issue1/3050>.

25. Mir, N. A., Gul, K., & Riar, C. S. (2015). Technofunctional and nutritional characterization of gluten-free cakes prepared from water chestnut flours and hydrocolloids. *Journal of Food Processing and Preservation*, 39(6), 978-984.
26. Mondal, T., Sarkar, T., Sengupta, S., & Kundu, S. (2024). Nutritional and pharmacological aspects of *Trapanatans*: An underutilized boon crop of West Bengal.
27. O'Neill, C. R., & Eill Jr, J. S. S. (2006). Water Chestnut (*Trapanatans*) in the Northeast NYSG Invasive Species Factsheet Series: 06-1. *New York Sea Grant; Sea Grant Brockport: New York, NY, USA*, 4.
28. Parekh, J. and Chanda, S. (2007). In vitro antimicrobial activity of *Trapanatans* L. fruit rind extracted in different solvents. *African J. Biotechnol.* 6: 766-770. doi: 10.4314/AJB.V6I6.56900.
29. Patel, S., Banji, D., Banji, O. J. F., Patel, M. M., & Shah, K. K. (2010). Scrutinizing the role of aqueous extract of *Trapabispinosa* as an immunomodulator in experimental animals. *Int J Res Pharm Sci*, 1(1), 13-19.
30. Van Driesche, R. (2002). *Biological control of invasive plants in the eastern United States*. US Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team.
31. PFAF, (2000). *Trapanatans*. Blagdon Cross, Ashwater, Beaworthy, Devon, UK: Plantsfor a Future Database. <http://www.pfaf.org/user/Plant.aspx?LatinName=Trapa%20natans>.
32. Purewal, S.S. and Kong, U. (1958). *Bagwan*, 11:22-24.
33. Puste, A.M. (2004). Agronomic management of wetland crops. *Kalyani Publishers, India*.
34. Radojevic, I. D., Vasic, S. M., Dekic, M. S., Radulovic, N. S., Delic, G. T., Durdevic, J. S., & Comic, L. R. (2016). Antimicrobial and Antibiofilm Effects of Extracts from *Trapanatans* L., Evaluation of Total Phenolic and Flavonoid Contents and Gc-Ms Analysis. *Acta Poloniae Pharmaceutica*, 73(6), 1565-1574.
35. Rani, B., Verma, D., Bhati.; Chharang, H. and Maheshwari, R.K. (2016). Health Benefits of Scrumptious Water Chestnuts /Water Caltrop (*Trapanatans* L.). *Int. Arch. App. Sci. Technol*; Vol 7 [3]: 32-36.
36. Razvy, M.A., Faruk, M.O. and Hoque, M.A. (2011). Environment friendly antibacterial activity of water chestnut fruits. *JBES*.; 01: 26- 34;2011.
37. Shafi M, Baba WN, Masoodi FA, Bazaz R (2016). Wheat-water chestnut flour blends: effect of baking on antioxidant properties of cookies. *J Food Sci Technol* 53(12):4278–4288.
38. Singh, G.D., Singh, S.; Jindal, N., Bawa, A.S. and Saxena, D.C. (2010). Physico-chemical characteristics and sensory quality of Singhara (*Trapanatans* L.): An Indian water chestnut under commercial and industrial storage conditions. *African Journal of Food Science*.; Vol. 4(11), pp. 693 – 702.
39. Singh, H., Thakur, S.N., Wilson, I., Kishor, K., Rai, B.S. (2017). Studies on quality parameters of bun incorporated with wheat flour water chestnut flour and soya flour. *J. Pharm. Innov*.; 6: 119-124;2017.
40. Singh, I. S., Thakur, A. K., & Prakash, D. (2018). Scientific cultivation of water chestnut crop a boon for poor Fishermen. *Indian Farming*, 68(3).
41. Takano, A. and Kadono, Y. (2005). Allozyme variations and classification of *Trapa* (*Trapaceae*) in *Japan*. *Aqu Bot.*; 83: 108–118.

42. Zhu, F. (2016). Chemical composition, health effects, and uses of water caltrop. *Trends in Food Science & Technology*, 49, 136-145.

UNDER PEER REVIEW