

## Field bio efficiency of insect growth regulators and insecticides against Pod Fly, *Melanagromyza obtusa* in Pigeonpea (*Cajanus Cajan*, L.)

### ABSTRACT

The present study on the bio efficacy of insect growth regulators and insecticides on pod fly was carried out during *kharif* 2021 in pigeonpea using Co 7 variety. An experiment was laid out using a randomized block design (RBD) with seven treatments and three replications. Insecticidal treatments consisted of four insect growth regulators and two insecticides along with untreated check and applications were made twice. Among the seven treatments tested, T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> proved to be effective in reducing podfly population (maggots+pupae) (3.67 No/25 pods). The other treatments tested were found to be on par except T<sub>1</sub>: Buprofezin 25 SC @ 200 g a.i. ha<sup>-1</sup> treated plots (6.33 Nos), whereas untreated check reported with 14.67 Nos at 14 DAS. At the same time, pod damage was also calculated and the lowest pod damage was recorded in T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> and T<sub>5</sub>: Lufenuron 5.4 EC @ 30 g a.i. ha<sup>-1</sup> treated plots and were significantly superior over other treatments and found to be on par with each other with 10.00 and 11.33 per cent, respectively. After two applications of T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> the pod fly damage reduction over untreated check was upto 80.1 per cent and other treatments reported the damage reduction between 42.2-72.5-78.0 per cent over untreated check. The highest yield was reported in plots treated with T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> for its highest grain yield of 757.7 kg ha<sup>-1</sup> with 43.2 per cent increase over untreated check. The other treated plots reported between 402.3-615.3-478.3-680.0 kg of grain yield ha<sup>-1</sup> with 12.54-34.62 per cent increase over untreated check. The highest Benefit :Cost ratio was obtained in the plots treated with T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> (1:1.6) and other treatments reported between 1:1.0 - 1:1.4 whereas untreated plots reported with lowest benefit cost ratio of 1:0.9.

**Key words:** Insect growth regulators, insecticides, efficacy, Podfly, pod damage, pigeonpea

### INTRODUCTION

Redgram or Tur or Pigeonpea, or *Cajanus cajan* L. is a significant pulse crop that is grown on 4.65 million hectares worldwide. It produces 3.43 million tonnes annually and has an average productivity of 780 kg per hectare. Due in significant part to damage from insect pests, pigeonpea productivity has stagnated

Formatted: Centered, Space After: 0 pt, Line Spacing: Multiple 1.15 li

Comment [MB1]: \*Abstract can be simplified \*grain damage information in the abstract is missing that to be added

Comment [MB2]: SC

Comment [MB3]: Podfly should be written separately as Pod fly (As it is true fly)

Comment [MB4]: SC

Comment [MB5]: Numbers per how many pods should be specified

Comment [MB6]: Numbers per how many pods should be specified

Formatted: Strikethrough

Comment [MB7]: Cross check the value with table value

Comment [MB8]: Cross check the calculated value of per cent increase over untreated check and enter the correct value

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Not Highlight

Formatted: Font: Arial Narrow, 12 pt, Bold

Formatted: Left

Comment [MB9]: Re write the sentence in meaningful way

over the past forty years. Among the biotic constraints, there are nearly three hundred species of insect-pests known to infest at various growth stages of crop growth. In the recent years, among all, pod fly is the important pest of Pigeonpea[1]. Pod fly, *Melanagromyza obtusa* (Malloch) is a hidden pest occurring at pod formation stage without external symptom of damage and causes yield loss upto 60 to 80 per cent [2]. Pod flies alone cause 20–80% of seed damage, while India accounts for almost 25% of global pulse production. The female pod fly does not exhibit any outward signs of injury; instead, she oviposits separately in the growing pods. The infested immature pods do not show external evidence of damage until the fully-grown larvae make an exit hole in the pod walls which results in complexity of their management [3]. Hence, continuous monitoring and special management practices were required to overcome this problem.

The pod fly attack remains unnoticed by the farmers owing to the concealed mode of life and thus poses challenge for management, since the oviposition occurs at the inner surface of the podwall involving anatomical, morphological and biochemical basis of preference. In spite of application of three to four rounds of application of insecticides against other podborers also resulted in the failure of control of *M. obtusa* which requires timely application. In order to manage this hidden insect, evaluation of insect growth regulators and insecticide molecules with different modes of action were studied to identify the level of protection of each molecule against podfly in pigeonpea ecosystem.

## **MATERIALS AND METHODS**

To ascertain the bio-efficacy of insect growth regulators and insecticides with various modes of action molecules against pod-flies, a field experiment was carried out during the ~~Kharif~~ season of 2021–2022 at Agricultural Research Station, Virinjipuram, Vellore, Tamil Nadu. Using the CO 7 variety, an experiment was set up in a randomized block design (RBD) with three replications, ~~eight~~<sup>seven</sup> treatments with a plot size of 25 m<sup>2</sup> with 60 cm x 30 cm spacing. The crop was grown using rainfed circumstances and throughout the flowering stage, it received irrigation based on need. Two sprays were administered in total, starting at the pod initiation stage and spaced fifteen days apart. Using destructive sampling, the number of pod flies, *M. obtusa*, maggots and pupae from 25 randomly selected pods per plant was determined.

The pod and seed damage, per cent reduction over ~~control~~ untreated check and increased yield over untreated check were also worked out using the formula. Thus the data obtained on the population, pod and seed damage and grain yield in different treatments were analyzed statistically using AGRES [4].

**Comment [MB10]:** \*How many plants observed, and total number of pods selected for the observations to be clearly mentioned. \*Observation intervals should be mentioned in the methodology

**Comment [MB11]:** Font to be italic

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Not Highlight

$$\text{Pod damage (\%)} = \frac{\text{Number of damaged pods}}{\text{Total number of pods}} \times 100$$

$$\text{Seed damage (\%)} = \frac{\text{Number of damaged seeds}}{\text{Total number of seeds}} \times 100$$

$$\text{Pod damage reduction over untreated check (\%)} = \frac{\text{Mean pod damage in untreated check} - \text{Mean pod damage in treatment}}{\text{Mean pod damage in untreated check}} \times 100$$

$$\text{Seed damage reduction over untreated check (\%)} = \frac{\text{Mean seed damage in untreated check} - \text{Mean seed damage in treatment}}{\text{Mean seed damage in untreated check}} \times 100$$

$$\text{Increase over untreated check (\%)} = \frac{\text{Yield in treatment} - \text{Yield in untreated check}}{\text{Yield in untreated check}} \times 100$$

## RESULTS AND DISCUSSION

The data on the population of podfly maggots and pupae, pod and seed damage were presented in Table 1. The pre-count population of maggots and pupae ranged from ~~12.00~~ 11.67-12.67 Nos per 25 pods and found to be non-significant. After the first application of treatments, at pod initiation stage, at 14 DAS (Days after spray), there existed significant difference among the various treated plots. Among the seven treatments tested, T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> was found to be superior over other and recorded 3.67 Nos per 25 pods followed by other treatments viz., T<sub>5</sub>: Lufenuron 5.4 % EC @ 30 g a.i. ha<sup>-1</sup>, T<sub>3</sub>: Dinotofuran 20 % SG @ 40 g a.i. ha<sup>-1</sup>, T<sub>2</sub>: Diafenthiuran @ 350 g a.i. ha<sup>-1</sup>, T<sub>6</sub>: Thiamethoxam @ 50 g a.i.

Formatted: Strikethrough

Comment [MB12]: Correct the chemical name as "Dinotefuran"

Comment [MB13]: Correct the chemical name as "Diafenthiuron 50 % WP"

Comment [MB14]: Thiamethoxam 25 % WG - concentration and formulation is missing (25 % WG)

ha<sup>-1</sup> were found to be on par with each other and recorded with 4.33-5.33Nos per 25 pods, respectively. The highest population of podfly was recorded in T<sub>1</sub>: Buprofezin 25 % SC @ 200 g a.i.ha<sup>-1</sup> (6.33Nos./25 pods) treated plots, whereas the untreated plots recorded 14.67 Nos per 25 pods at 14 days after first spray. At the same time, pod damage was also calculated and the lowest pod damage was recorded in T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> and T<sub>5</sub>: Lufenuron 5.4 EC @ 30 g a.i. ha<sup>-1</sup> treated plots and were significantly superior over other treatments and found to be on par with each other with 10.00per cent and 11.33 per cent damages, respectively. Almost, similar trend of efficacy of different treatments was also observed with respect to seed damage and the highest seed damage was reported in untreated check (41.58 %). Before the second application of treatments, there existed significant difference in the pre-count population of podfly maggots and pupae and ranged between ~~6.776.67~~ - 16.00 Nos per 25 pods. Even after the second application of treatments also, T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> treated plots was significantly superior over other treatments and found to be on par with each other and recorded 3.15 Nos per 25 pods with 3.72 per cent pod damage and 11.88 per cent seed damage.

The mean per cent pod and seed damage was worked out after imposing the two applications of treatments and the mean per cent pod damage ranged between 6.90-34.6 with its lowest in T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> treated plots and the highest in untreated check. Likewise, the mean per cent seed damage was also ranged from 12.2-40.4 with its lowest in T<sub>4</sub>: Flubendiamide 480 Sc @ 30 g a.i. ha<sup>-1</sup> treated plots and the highest in untreated check.

After two applications of T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> the podfly damage reduction over untreated control was upto 80.1 per cent and other treatments reported the damage reduction between 42.2-~~72.5~~ 78.0 per cent over untreated check. The highest yield was reported in plots treated with T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> for its highest grain yield of 757.7 kg ha<sup>-1</sup> with 43.1 per cent increase over untreated check. The other treated plots reported between ~~431.6~~ 478.3 - 680 kg of grain yield per ha with 17.52 -36.53 per cent increase over untreated check. The highest Benefit :Cost ratio was obtained in the plots treated with T<sub>4</sub>: Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> (1:1.6) and other treatments reported between 1:~~1.01.1~~ - 1:1.4 whereas untreated plots reported with lowest benefit cost ratio of 1:0.9 (Table 2).

The current results corroborate those of [5] who found that the novel pesticide Chlorantraniliprole 18.5 SC @ 30 g a.i. ha<sup>-1</sup> reduced pod flies more effectively due to the similar diamide group. The results of [6] were likewise consistent with the timing of insecticide application, which was first applied during the pod start phases and then at a 15-day interval.

**Comment [MB15]:** With given CD these treatments also found at par with T<sub>4</sub>: Flubendiamide 480 SC, Cross check

**Formatted:** Not Highlight

**Formatted:** Strikethrough, Not Highlight

**Formatted:** Not Highlight

**Formatted:** Strikethrough

**Formatted:** Not Highlight

**Formatted:** Strikethrough

**Comment [MB16]:** Re calculate the value

**Formatted:** Highlight

**Formatted:** Strikethrough

**Comment [MB17]:** Re calculate the values (calculation error found for yield increase over untreated check)

**Comment [MB18]:** check

**Formatted:** Strikethrough

**Comment [MB19]:** Chlorantraniliprole

[7] also reported the efficacy of newer insecticides proved to be very effective against podfly in pigeonpea. Applying the insecticide dimethoate 30 EC @ 2 ml l<sup>-1</sup> at 2, 3, 4, and 5 weeks after 50% flowering was found to be beneficial in minimizing pod fly-induced damage to pods and grains while increasing yield [8]. The pod fly age was 13 to 25 days, and oviposition which produced the most eggs, was facilitated by 10 to 15 days, according to [9] and [10]. Therefore, it can be said that using Flubendiamide 480 SC @ 30 g a.i. ha<sup>-1</sup> twice throughout the pod-initiation stages was very helpful in reducing pod fly infestation and boosting grain yield in order to achieve the maximum Benefit:Cost ratio.

## CONCLUSION

In recent days, pod fly acts as the primary cause for decline in pigeon pea production, which has resulted in significant losses for farmers across India. The pod fly is a quiet pod killer that kills juvenile pods without causing visible harm until the fully grown larvae create an escape hole in the pod walls, making maintenance more difficult. Therefore, constant observation and extra care are always needed for the efficient control of this pest in order to prevent significant losses and insecticides should be used in rotation with different modes of action so as to prevent or delay the development of resistance towards insecticides and also to conserve natural enemies in pigeonpea ecosystem.

**Comment [MB20]:** Write the conclusion for the experiment conducted, don't give general statements in the conclusions. In the conclusion experimental outcomes like chemical and dosage which found superior in reducing the pod fly incidence can be mentioned.

## REFERENCES

1. Lal SS, Singh NB. In: Proceedings of National Symposium on Management of Biotic and Abiotic Stresses in Pulse Crops, 2018; Indian Institute for Pulse Research, Kanpur, India: 65-80.
2. Durairaj C. Evaluation of certain neem formulations and insecticides against Pigeonpea pod fly. *Indian Journal of Pulses Research*, 2006;19 (2): 269-270
3. Sharma OP, Gopali JB, Yelshetty, Suhas, Bambawale OM, Garg DK, Bhosle BB. Pests of pigeonpea and their management, 2010; NCIPM, LBS Building, IARI Campus, New Delhi-110012, India:4.
4. Gomez K A, Gomez A A. Statistical procedures for agricultural research, 1984; John Wiley and sons, New York 207-215
5. Chiranjeevi B, Sarnaik SV. Efficacy of different insecticidal treatments on population of pod fly, *Melanagromyza obtusa* (Malloch). *Journal of Entomology and Zoology studies*, 2014; 5(4): 1812-1815
6. Sreekanth M, Lakshmi MSM, Rao, YK. Efficacy of insecticides in the management of podfly, *Melanagromyza obtusa* on pigeonpea. *Indian Journal of Plant protection*, 2013;41(3):212-214.
7. Singh AK. Evaluation of new molecule of insecticides against pod fly *Melanagromyza obtusa* of Pigeonpea. *SAARC Journal of Agriculture*, 2014; 12(1):89-95

8. ReddyDVC, Sreekanth M, Sandhya Rani C, AdinarayanaM, Ramarao G. *International Journal of Chemical studies*, 2010;**6**(3):316-319
9. DasSB, OdakSC. Pigeonpea pod age and podfly *Melanagromyza obtusa* (Malloch) infestation on redgram (*Cajanus cajan*). *Agriculture Science Digest*, 1991; 11(2):63-64
10. DurairajC. Timing of insecticide application for the control of pigeonpeapodfly. *Madras Agricultural Journal*. 2000;**87**(10-12):628-631

UNDER PEER REVIEW

Comment [MB21]: \*For pre count data a ... [1]

Formatted ... [5]

Formatted ... [8]

Formatted ... [3]

Formatted ... [7]

Formatted ... [9]

Formatted ... [10]

Formatted ... [4]

Formatted ... [2]

Formatted ... [11]

Comment [MB22]: Data should be given ... [6]

Formatted ... [15]

Formatted ... [19]

Formatted ... [13]

Formatted ... [12]

Formatted ... [17]

Formatted ... [14]

Formatted ... [16]

Formatted ... [18]

Formatted ... [20]

Formatted ... [21]

Formatted Table ... [22]

Formatted ... [23]

Formatted ... [24]

Formatted ... [25]

Formatted ... [26]

Formatted ... [27]

Formatted ... [28]

Formatted ... [29]

Formatted ... [30]

Formatted ... [31]

Formatted ... [32]

Formatted ... [33]

Formatted ... [34]

Formatted ... [35]

Formatted ... [36]

Formatted ... [37]

Formatted ... [38]

Formatted ... [39]

Formatted ... [40]

Formatted ... [41]

Formatted ... [42]

Formatted ... [43]

Formatted ... [44]

Table 1. Evaluation of insect growth regulators and insecticides against podfly, *Melanagromyza obtusa* in pigeonpea

Treatments	Dose/L Dose (g a.i./ha)	No. of maggots & pupae (No./25 pods) Pod fly maggots + Pupae per 25 pods, pod damage (%) and grain damage (%)								Grain Yield (Kg/ha) DA
		First application spray				Second application spray				
		Precount (No./25 pods)	14 DAS (No./25 pods)	Pod damage (%)	Seed damage (%)	Precount (No./25 pods)	14 DAS (No./25 pods)	Pod damage (%)	Seed damage (%)	
T <sub>1</sub> :Buprofezin 25 % SC	200	11.67	6.33 (2.52) <sup>c</sup>	29.33 <sup>d</sup>	27.46 <sup>d</sup>	9.33 (3.06) <sup>b</sup>	4.76 (2.18) <sup>ab</sup>	10.67 <sup>d</sup>	24.33 <sup>c</sup>	478.3
T <sub>2</sub> :Difenthiuran on 50 % WP	350	12.67	5.33 (2.31) <sup>ab</sup>	16.67 <sup>c</sup>	20.88 <sup>c</sup>	8.00 (2.83) <sup>ab</sup>	5.21 (2.28) <sup>b</sup>	8.70 <sup>c</sup>	21.63 <sup>c</sup>	584.7
T <sub>3</sub> :Dinotofuran 20 % SG	40	11.67	4.67 (2.16) <sup>ab</sup>	15.33 <sup>bc</sup>	16.38 <sup>bc</sup>	8.00 (2.83) <sup>ab</sup>	4.46 (2.11) <sup>ab</sup>	6.49 <sup>b</sup>	15.88 <sup>b</sup>	576.7
T <sub>4</sub> :Flubendiamide 480 SC	30	11.67	3.67 (1.91) <sup>a</sup>	10.00 <sup>a</sup>	12.42 <sup>a</sup>	6.67 (2.58) <sup>a</sup>	3.15 (1.77) <sup>a</sup>	3.72 <sup>a</sup>	11.88 <sup>a</sup>	757.7
T <sub>5</sub> :Lufenuron 5.4 % EC	30	12.00	5.33 (2.31) <sup>ab</sup>	11.33 <sup>ab</sup>	16.77 <sup>ab</sup>	8.00 (2.83) <sup>ab</sup>	4.71 (2.17) <sup>ab</sup>	3.82 <sup>bc</sup>	12.85 <sup>a</sup>	680.0
T <sub>6</sub> :Thiamethoxam 25 % WG	50	12.33	4.33 (2.08) <sup>ab</sup>	14.67 <sup>bc</sup>	14.92 <sup>bc</sup>	7.33 (2.71) <sup>ab</sup>	4.18 (2.04) <sup>ab</sup>	4.28 <sup>a</sup>	14.96 <sup>b</sup>	528.3
T <sub>7</sub> :Untreated Check	-	12.67	14.67 (3.83) <sup>d</sup>	45.33 <sup>e</sup>	41.58 <sup>e</sup>	16.00 (4.00) <sup>c</sup>	16.00 (4.00) <sup>c</sup>	23.78 <sup>e</sup>	39.21 <sup>d</sup>	434.6
	SEd		0.20	1.69	2.08	0.19	0.28	1.75	1.50	
	CD<0.5 %	NS	0.44	3.68	4.54	0.42	0.61	3.83	3.28	

DAS = Days after spray

**Table 2. Cost Economics studies for insecticides on podfly, *M. obtusa* damage in pigeonpea**

Treatments	Dose (g a.i/ha)	Mean damage after two applications (%)		Reduction over control untreated check (%)		Grain Yield (Kg/ha)	Yield Increase over untreated check (%)	B: C ratio
		Pod damage	Seed damage	Pod damage	Seed damage			
T <sub>1</sub> :Buprofezin 25 % SC	200	20.0	25.9	42.2	35.9	478.3	9.8 10.82	1:1.1
T <sub>2</sub> :Difenthiuran 50 WP Difenthiuron 50 % WP	350	12.7	21.3	63.3	47.3	581.7	25.8 34.78	1:1.3
T <sub>3</sub> :Dinotefuran 20 SG Dinotefuran 20 % SG	40	10.9	16.1	68.5	60.1	576.7	25.2 33.62	1:1.3
T <sub>4</sub> :Flubendiamide 480 SC	30	6.9	12.2	80.1	69.8	757.7	43.0 75.56	1:1.6
T <sub>5</sub> :Lufenuron 5.4 % EC	30	7.6	14.8	78.0	63.4	680.0	36.5 57.55	1:1.4
T <sub>7</sub> :Thiamethoxam 25 % WG	50	9.5	14.9	72.5	63.1	523.3	17.5 21.25	1:1.2
T <sub>9</sub> :Untreated Check	-	34.6	40.4	-	-	431.6	-	1:0.9
<b>SEd</b>						12.2		
<b>CD&lt;0.5 %</b>						24.6		

**Comment [MB25]:** \*Data in the table should be given for two digits after the decimals

**Formatted:** Strikethrough

**Formatted Table**

**Comment [MB27]:** \*BC ratio should be given in separate table which should include cost of the chemical per two spray per hectare, labour cost for sprays, other cost of cultivation, gross return, net return and BC ratios

**Comment [MB26]:** Cross check the calculations of yield increase over untreated control treatment

**Formatted:** Left, None, Space Before: 0 pt, Line spacing: single, Don't keep with next, Don't keep lines together

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Formatted:** Strikethrough

**Comment [MB28]:** Instead SEd values S.Em.± values should be added

**Comment [MB29]:** C.D. @ 5 %

**Page 7: [1] Comment [MB21] MANJUNATH BADIGER 11/6/2024 11:12:00 PM**

\*For pre count data also transformation values should be added

\*For pod and seed damages also transformation values should be added (arcsine transformation values)

\*There is sudden rise in the population of pod fly (maggots + pupa) between 14 DAS of 1<sup>st</sup> spray and Pre count of 2<sup>nd</sup> spray with a gap of only 1 to 2 days - clarify

\*Maggot and pupa count data can be given as separate parameters instead of combined

**Page 7: [2] Formatted MANJUNATH BADIGER 11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [3] Formatted MANJUNATH BADIGER 11/6/2024 4:05:00 PM**

Strikethrough

**Page 7: [4] Formatted MANJUNATH BADIGER 11/6/2024 4:05:00 PM**

Not Strikethrough

**Page 7: [5] Formatted MANJUNATH BADIGER 11/6/2024 4:14:00 PM**

Not Strikethrough

**Page 7: [6] Comment [MB22] MANJUNATH BADIGER 11/6/2024 11:19:00 PM**

Data should be given for the two digits after the decimals

**Page 7: [7] Formatted MANJUNATH BADIGER 11/6/2024 4:42:00 PM**

Strikethrough

**Page 7: [8] Formatted MANJUNATH BADIGER 11/6/2024 4:14:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [9] Formatted MANJUNATH BADIGER 11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [10] Formatted MANJUNATH BADIGER 11/6/2024 4:42:00 PM**

Strikethrough

**Page 7: [11] Formatted MANJUNATH BADIGER 11/6/2024 4:14:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [12] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [13] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [14] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [15] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [16] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [17] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [18] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [19] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [20] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single, Don't keep with next, Don't keep lines together

**Page 7: [21] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [22] Change** **MANJUNATH BADIGER** **11/6/2024 4:43:00 PM**

Formatted Table

**Page 7: [23] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [24] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [25] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [26] Formatted** **MANJUNATH BADIGER** **11/6/2024 3:58:00 PM**

Strikethrough

**Page 7: [27] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [28] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [29] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [30] Formatted** **MANJUNATH BADIGER** **11/6/2024 3:59:00 PM**

Strikethrough

**Page 7: [31] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [32] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [33] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [34] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [35] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [36] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [37] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [38] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [39] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [40] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [41] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [42] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [43] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Centered, Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [44] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [45] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [46] Comment [MB23]** **MANJUNATH BADIGER** **11/6/2024 11:12:00 PM**

Instead SEd values S.Em.± values should be added

**Page 7: [47] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:14:00 PM**

Space After: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single

**Page 7: [48] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:41:00 PM**

None, Space Before: 0 pt, Don't add space between paragraphs of the same style, Line spacing: single,  
Don't keep with next, Don't keep lines together

**Page 7: [49] Formatted** **MANJUNATH BADIGER** **11/6/2024 4:42:00 PM**

Indent: Left: 0.06"