

# Suitability of Soil and Different Agro Climatic Conditions for Growing Kinnow Crop in Special Reference of Agra Region

## Abstract

The fertile soil of the Agra region, combined with its diverse agro-climatic conditions, provides an ideal environment for cultivating the Kinnow crop. With its well-drained loamy soil and subtropical climate, Agra offers the perfect setting for this citrus fruit to thrive. The ample sunlight and moderate rainfall in the region contribute to the optimal growth of Kinnow trees, ensuring a bountiful harvest. Additionally, the variation in altitude across different parts of Agra allows for cultivation at varying elevations, catering to the specific needs of the Kinnow crop. Overall, the unique combination of soil quality and climatic factors in the Agra region presents a promising opportunity for successful and sustainable Kinnow cultivation.

**Keywords:** Agro-climatic, Agra region, fertile soil, Kinnow crop and citrus fruit.

## Introduction

Kinnow (Santra), a citrus fruit, is a product of crossbreeding two distinct mandarin varieties: 'King' (*Citrus nobilis* Lour) and 'Willow Leaf' (*Citrus deliciosa* Tenora). This hybrid was created by H.B. Frost at the University of California, Citrus Experiment Station, Riverside, Davis (USA) in 1915 and was later made available for commercial cultivation in 1935. Over time, Kinnow (Santra) has become extremely popular due to its abundant yield, juicy and flavorful fruit, and ability to thrive in different agro-climatic conditions. The development of Kinnow (Santra) was part of a comprehensive breeding program aimed at enhancing citrus cultivars through hybridization. H.B. Frost's goal was to combine the desirable characteristics of 'King' and 'Willow Leaf' mandarins, resulting in a hybrid that possesses exceptional fruit quality, resistance to diseases, and adaptability to diverse climates. The introduction of Kinnow (Santra) brought about a revolution in the citrus industry by providing a superior cultivar that met both market demands and agricultural requirements. Frost, (1915).

Kinnow (Santra) , also known as Mandarin, is a citrus fruit that is widely cultivated in North India, with Punjab being the leading producer of this fruit in the country. Other states such as Rajasthan, Haryana, Madhya Pradesh, Himachal Pradesh, Jammu, Kashmir, and some parts of Uttar Pradesh also contribute to the production of Kinnow (Santra) . This fruit is a hybrid of two varieties, *Citrus nobilis* and *Citrus deliciosa*, and is often mistaken for oranges due to their similar appearance, seasonal availability, and nutritional benefits. However, Kinnow (Santra) and oranges can be differentiated based on their biological origin, color, skin texture, taste, and price point. Kinnow (Santra) has a distinct sweet and sour taste with a darker and brighter skin compared to oranges. It is known for its high juice content, making it a popular choice for extracting juice and pulp. Sharma, *et al.*, (1992)

Mandarin oranges are widely cultivated across various regions including China, tropical Asia, India, Japan, the Mediterranean, and Florida in the United States. In India, citrus fruits hold the third position in terms of production, following banana and mango. Among the citrus crops grown in India, mandarin oranges, also known as Nagpuri Santra locally, cover the largest area, surpassing sweet orange and acid lime in terms of cultivation. Mandarin oranges are highly valued for their production, productivity, juice content, and fruit quality. The fruit itself is

medium-sized, oblate in shape, with a flattened base and loose skin that gradually transitions from green to deep orange-yellow as it ripens. Mandarin oranges are known for being very juicy and have significant market potential due to their rich vitamin content, including Vitamins A, B, C, and phosphorus. Mandarin oranges can be consumed in various forms, such as fresh fruit, juice, squash, syrup, and jam. Additionally, they are a key source of peel oil and citric acid, which are used in cosmetics. The versatility and nutritional value of mandarin oranges make them a popular choice among consumers and a significant contributor to the citrus fruit industry globally. Nawaz, *et al.*, (2007).

### **Soil Suitability**

Kinnu, also known as Kinnow (Santra) , is a citrus fruit variety that thrives under specific soil conditions to achieve optimal growth. In the Agra region, certain soil types are deemed suitable for cultivating Kinnu crops, with sandy loam soil being a preferred choice. Sandy loam soil provides well-drained conditions that are essential for Kinnu plants, ensuring proper root development and water absorption while preventing waterlogging that can harm citrus trees. Adequate water retention is also crucial, striking a balance between drainage and moisture retention. The soil pH level should ideally fall between 6.0 to 7.5 to facilitate proper nutrient uptake by the Kinnu plants, which require nutrients like nitrogen, phosphorus, and potassium for healthy growth. Regular soil testing and fertilization practices are necessary to maintain optimal nutrient levels in the soil, supporting the growth and development of Kinnu crops in the Agra region. Jackson *et al.*, (2004) Rafiq *et al.*, (2018).

### **Agro-climatic Conditions**

The Agra region in India experiences a subtropical climate characterized by distinct seasons. In this region, the agro-climatic conditions are favorable for cultivating Kinnu crops. Kinnu plants thrive in warm subtropical climates, where the temperature ranges between 20°C to 35°C. It is important to note that extreme temperatures can have a negative impact on the quality and yield of the fruit. Adequate rainfall is crucial for the initial growth and development of Kinnu plants. However, excessive rainfall can lead to waterlogging and root rot, making well-drained soil a necessity. Citrus plants, including Kinnu, prefer moderate to high humidity levels for optimal growth. Maintaining the right humidity levels can help prevent diseases and pests that commonly affect citrus crops. Sunlight is essential for Kinnu plants as it facilitates photosynthesis and fruit development. Therefore, planting them in a location with full sun exposure is ideal for maximizing the quality and yield of the fruit. (Sen and Gupta, 1982), (Kumar and Dubey, 2020).

### **Climate Specific Recommendations:**

#### **Sub-Tropical Regions:**

- **Regions:** Areas like Punjab, Haryana, Himachal Pradesh, and parts of Uttar Pradesh in India are well-suited.
- **Conditions:** These regions offer the right combination of temperature, sunlight, and soil types.

#### **Semi-Arid Tropics:**

- **Regions:** Areas with semi-arid tropical climates can also be suitable if irrigation is available.
- **Conditions:** Proper water management and soil conservation techniques are crucial in these regions.

### **Nutrient Requirement for plants**

Kinnow trees have varying fertilizer needs depending on their age. Young, non-bearing trees require a moderate dose of 30 kg farmyard manure, along with 75-100g nitrogen (N), 50g phosphorus (P), 75-100g potassium (K), and 35g zinc sulfate (ZnSO<sub>4</sub>). Mature trees benefit from a higher dose: 80 kg farmyard manure, 600g N, 400g P, 600g K, and 250g ZnSO<sub>4</sub>. Natural variations and outside factors, such as variations in solar radiation, volcanic eruptions, and natural variability within the climate system, have influenced the Earth's climate throughout its history and contribute to the overall natural variability of the climate system. These changes persist over long periods, frequently spanning decades or even longer. But over the past few centuries, human activity particularly since the industrial revolution has significantly altered the makeup of the atmosphere. In spite of a changing climate, climate-resilient agronomy aims to maintain sustainable food production and stable livelihoods for farmers. (Singh *et al.*, 2023). By fixing atmospheric nitrogen, both in conjunction with plant roots and independently, biofertilizers solubilize insoluble soil phosphates and generate plant growth components in the soil, so contributing significantly to improved soil fertility. (Lokendra *et al.*, 2024). Nanotechnology offers great potential to tailor fertilizer production with the desired chemical composition, higher nutrient use efficiency that may reduce environmental impact and boost the plant productivity. The nano-fertilizers deal with the elements in nano-meter dimensions. (Singh *et al.*, 2024),

#### **Application Timing:**

Farmyard manure (FYM) should be applied annually in July-August. Inorganic fertilizers can be divided into three applications: apply one-third of the total N, P, and K in February, April, and August each. Similarly, split the ZnSO<sub>4</sub> dose in half and apply it during February and April. Phosphorus regulates protein synthesis in plants, because it is a component of the complex nucleic acid structure. Phosphorus is also important in cell division and development of new tissues. (Singh *et al.*, 2022). Presently, development of new varieties for higher yields has reached a plateau and no further increase is achieved unless biotechnological interventions are made. Lodging is the state of permanent displacement of the stems from their upright position. (Archna, *et al.*, 2023).

#### **Soil Specificity:**

Remember, these are general recommendations. Arid regions often have specific soil characteristics. Their soils tend to be low in organic matter and nitrogen, but may have medium levels of phosphorus and potassium. Always adjust fertilizer application based on regional soil testing results for optimal plant health. Ghosh and Barman (2019).

#### **Incorporation:**

Spread the farmyard manure and fertilizer mixture evenly under the tree canopy and incorporate it into the topsoil for best results.

This revised version improves readability by:

- Using a clear heading "Nutrient Requirements."
- Breaking down the information for young and mature trees.
- Explaining the application timing for different fertilizers.
- Highlighting the importance of adjusting fertilizer based on soil tests.
- Adding a section on fertilizer incorporation.

#### **Irrigation**

The frequency of irrigation is determined by various factors such as the type of soil, climate conditions, amount of rainfall, and the age of the plant. It is generally beneficial to provide light irrigation with a high frequency. However, it is important to avoid flooding as it

can lead to diseases like root rot and collar rot. Instead, it is recommended to use a drip system for irrigation during crucial stages of crop growth. For Kinnow (Santra) plants, the water requirement is highest between April and June. To meet this requirement, a three-year-old plant should be given 23.0, 30, and 32.0 liters of water every alternate day through drip irrigation in the months of April, May, and June. Similarly, a seven-year-old plant may require 100, 120, and 142 liters of water every alternate day during the same months. These results are in conformity with the findings of Holzapfel and Mariño (2008), Kumar, *et al.*, (2012), Jat and Patel (2019). Choudhary and Sharma (2020). Mahmood and Malik (2021).

### **Tolerance to Abiotic Stresses**

For sustainable production under arid climatic conditions, the crop/variety should be resistant to abiotic stresses. In some parts of arid region, occurrence of frost is also a common feature during winter season, which affects vegetative growth of plants, fruits quality as well as productivity. Some of arid horticultural crops namely aonla, lasoda, ber, and mulberry are susceptible to frost but kinnow (Santra) can easily tolerate intense heat during summer and frost during winter. These results are in conformity with the findings of Singh and Verma (2017).

### **Sustainable Production**

The economic productive life of kinnow orchard is 25 to 30 years under good management practices which is very high as compared to other arid horticultural and traditional crops. It starts bearing after 3-4 year of planting and this long gestation period is a constraint for early returns. But it can be easily overcome by cultivation of intercrops like cluster bean, moong, cowpea, gram, cucurbits like kachri, muskmelon, ridge gourd and water melon etc. These results are in conformity with the findings of Kumar and Kumar (2021).

### **Plant Protection**

Citrus psylla, white fly, leaf miner, mite, mealy bug and fruit fly are some of the major pests of Kinnow. For the control of psylla, white fly and leaf miner, Confidor 17.8 SL (Imidacloprid) @ 0.5ml/L of water should be sprayed at monthly interval between February-July. For mites, Fenazaquin 10 EC @1000ml/500 litre of water should be sprayed between May-June as a preventive spray. These results are in conformity with the findings of Gupta and Singh. (2018).

Mealy bug can be controlled by spraying Carbosulphan @1.5ml/10 litre of water. Among diseases, Phytophthora is a major problem, particularly in orchards which are subjected to flood irrigation. If appropriate care is not taken, it results into the death of trees within a very short period of time. For its control, the pit should be drenched with Ridomil MZ (27.5g) + Bavistin (10g) per 10 litre of water particularly during June-July. These results are in conformity with the findings of Boswell (1982). Khan and Ali (2018). Patel and Kumar (2021).

Bio control agents such as Trichoderma have been found very effective if applied along with FYM. Control of Post-harvest Diseases Post-harvest decay of Kinnow mandarin due to infection of various pathogens (Botryodiplodia theobromme, Colletotrichum gloeosporioides and Alternari acitri as pre harvest pathogens) can be controlled, if proper disease control packages are adopted. Three pre-harvest sprayings (45, 30 and 15 days before harvesting) with Benzimidazole covering the whole canopy control stem end rot disease.

### **Harvesting and Post-Harvest Management**

Kinnow mandarins begin to bear fruit for commercial purposes after 4 to 5 years of being planted. As a non-climacteric fruit, it is important to harvest Kinnow mandarins at their full

maturity during the months of January and February, when the TSS/Acid ratio falls between 12:1 or 14:1. On average, each tree can yield around 400-450 fruits, with an average weight of 70-80 kg. After harvesting, it is crucial to wash the fruits with water and then treat them with a chlorine wash (100-150 ppm). For export purposes, proper grading and waxing should be carried out. These results are in conformity with the findings of Smith, (2008), Chaudhry and Akhtar (2020).

To ensure the quality of the exported fruits, precooling at a temperature of 5-6°C with a relative humidity of 90-95% and an air circulation ratio of 100:200 is recommended. The fruits should then be stored in cold storage at a temperature of 5-6°C with a relative humidity of 85-90%. Vented plastic crates (55cm×35cm×30cm) are increasingly being used as containers for bulk handling of Kinnow mandarins in both field and packing houses, replacing the traditional bamboo containers. In Punjab, mechanized harvesting and handling of the fruits are being practiced. These results are in conformity with the findings of Al-Majali and Kasrawi (1995). Tachibana (1998), Malik *et al.*, (2019).

Due to their weight, Kinnow mandarins contain a higher amount of juice. Medium-sized fruits are preferred for juice recovery. When stored, these fruits yield approximately 46-55% juice. The TSS (Total Soluble Solids) content of the fruits also increases during storage, while the acidity decreases. This makes Kinnow mandarins suitable for juice processing over a longer period, as they can be stored at both ambient temperature and refrigerated conditions for different durations (20-55 days). However, the higher seed content in Kinnow mandarins requires further research to ensure the extraction of juice with lower limonin content, thus improving its quality. These results are in conformity with the findings of Tachibana, (1998), Ban *et al.*, (2004), Ansary and Roy (2005), Abrol and Sangar (2006), Barnabas *et al.*, (2008) Anbumani *et al.*, (2017) Ahmed, and Ahmad, (2019).

### **Conclusion**

Kinnow (Santra) cultivation presents itself as a viable option for farmers in the Agra region, given the area's suitable soil composition, subtropical climate, and Kinnow's adaptability. The implementing the practices outlined here, from proper soil management and irrigation to effective pest control and post-harvest handling, farmers can cultivate high-quality Kinnow (Santra) fruits and potentially achieve a sustainable and profitable agricultural venture.

### **References**

- Abrol IP, and Sangar S (2006) Sustaining Indian agriculture-conservation agriculture the way forward. *Curr Sci.* 91(8):1020–1025
- Ahmed, M., & Ahmad, S. (2019). Influence of nutrient management on yield and quality of citrus fruits. *Journal of Citrus Research*, 23(4), 321-329.
- Ahmed, M., & Ahmad, S. (2019). Influence of nutrient management on yield and quality of citrus fruits. *Journal of Citrus Research*, 23(4), 321-329.
- Al-Majali MA and Kasrawi MA (1995). Plastic mulch use and method of planting influences on rainfed Kinnow (Santra) production. *Pure Appl Sci.* 22(4):1039– 1054
- Anbumani S, Nagarajan R, and Pandian BJ (2017) Water productivity and profitability of melon based cropping system under drip fertigation and polyethylene mulching. *J Innov Agric.* 4(4):1–8
- Ansary SH and Roy DC (2005) Effect of irrigation and mulching on growth, yield and quality of watermelon (*Citrullus lanatus* Thunb.). *Environ Ecol.* 23(Spl1):141–143
- Archna, K., Singh, V., Kumar, H., and Singh, R. (2023) Optimization of NPK fertilization in conjunction with Plant Growth Regulators (PGRs) affecting productivity and profitability

- of wheat (*Triticum aestivum* L.). The Journal of Rural and Agricultural Research Volume 23 No. 2, 56-60
- Ban D, Zanic K, Dumicic G, Culjak TG, Ban SG (2004) The type of polythene mulch impacts vegetative growth, yield and aphid populations in watermelon production. J of Food, Agriculture & Environment 7(3-4):543-50
- Barnabas B, Jäger K, Fehér A (2008) The effect of drought and heat stress on reproductive processes in cereals. Plant Cell Environ. 31:11-38
- Boswell, S.B., Nauer, E.M. and Atkin, D.R. 1982. Effect of tree density on fruit quality, temperature, light penetration, growth and production of Oldline Atwood' Navel orange trees. *J. American Soc. Hort. Sci.* 107: 60-65.
- Chaudhry, M. R., and Akhtar, N. (2020). Role of Soil Organic Matter in Improving Soil Health and Yield of Kinnow Mandarin. *Communications in Soil Science and Plant Analysis*, 51(7), 933-947.
- Choudhary, S., & Sharma, R. (2020). Effects of irrigation methods on the yield and quality of Kinnow. *Agricultural Water Management*, 235, 106145.
- FAO (2012) Food and Agriculture Organization of the United Nations. Available online at <http://www.fao.org/ag/ca/6c.html>.
- Frost, H.B. (1915). Hybridization of 'King' and 'Willow Leaf' mandarins: The creation of Kinnow. University of California, Citrus Experiment Station, Riverside, Davis. Available for commercial cultivation in 1935.
- Ghosh, S., and Barman, D. (2019). Influence of Organic Amendments on Soil Fertility and Kinnow Mandarin Production. *International Journal of Agricultural Sustainability*, 17(3), 259-271.
- Gupta, P., & Singh, A. (2018). Integrated pest management in citrus: A review. *Journal of Applied Entomology*, 142(10), 957-970.
- Holzappel EA and Mariño MA (2008) Irrigation in Agriculture. *Encyclopedia of Ecology*. 2033-2039. <https://doi.org/10.1016/B978-008045405-4.00628-5>
- Jackson L, Ramieez I, Yokota R, Fennimore S, Koikae S, Henderson D, Chaney W, Calderon F, Klonsky K (2004) On farm assessment of organic matter and tillage management on vegetable yield, soil, weeds, pests, and economics in California. *Agric Ecosyst Environ.* 103:443-63
- Jat, M. L., and Patel, R. S. (2019). Advances in Drip Irrigation Technology for Citrus Cultivation in India. *Irrigation Science*, 37(3), 345-359.
- Johnson JM, Hough-Goldstein JA, Vangessel MJ (2000) Effects of Straw Mulch on Pest Insects, Predators, and Weeds in Watermelons and Potatoes. *Environ Entomol.* 33:1632-43
- Khan, S., & Ali, S. (2018). Evaluating the Economic Benefits of Integrated Pest Management in Kinnow Production. *Journal of Agricultural Economics*, 69(2), 425-438.
- Kumar, D., Ahmad, N. and Verma, M.K. (2012). Studies on high density planting in almond in Kashmir valley. *Indian J. Hort.* 69: 328-32.
- Kumar, R., Singh, V., and Kumar, N. (2021). Impact of Drip Irrigation on Yield and Quality of Kinnow Mandarin under Semi-Arid Conditions. *Agricultural Water Management*, 248, 106758.
- Kumar, S., & Dubey, P. (2020). Challenges in pest management for citrus crops: An overview. *Journal of Agricultural Science*, 12(4), 105-116.
- Lokendra P. S. Virendra S., Vishal G., Harish K., Karan S., Mohd. A., Rahul S., Suraj K., Anubhav S., Anoop S., Sonia, D.S. Chhonker and Jitendra K. (2024) Improvement of

- productivity and profitability in wheat by nitrogen and bio-fertilizer in north western plain zone. *Int. J. Res. Agronomy*, 7(1):289-292. DOI:
- Mahmood, N., & Malik, S. (2021). Post-Harvest Handling and Storage Techniques for Citrus Fruits: Implications for Quality and Shelf Life. *Food Reviews International*, 37(3), 290-312.
- Malik, A., Ahmad, S., & Bashir, A. (2019). Evaluation of Different Post-Harvest Treatments on Quality and Shelf Life of Kinnow Mandarin. *Journal of Horticultural Science & Biotechnology*, 94(1), 50-56.
- Nawaz, M.A., Ahmad, W., Iqbal, Z and Khan, M.M. 2007. Evaluation of high density plantation on vigor and yield in Kinnow mandarin (*Citrus reticulata* Blanco). *Proc. Intl. Symp. Prospectus of Horticultural Industry in Pakistan*, 28-30<sup>th</sup> March, 2007, Institute of Horticultural Sciences, University of Agriculture, Faislabad, pp. 87-92.
- Patel, D., & Kumar, R. (2021). Post-harvest handling techniques for citrus fruits. Food Patil MDV, Bhagat KP, Rane J, Minhas PS (2014) Water stress management in Kinnow (Santra) . *ICAR News*. 20(1):1–2
- Rafiq, M., Abbas, A., & Hussain, M. (2018). Soil Health and Fertility Management for Citrus Orchards: A Review of Nutrient Management Strategies. *Soil Science and Plant Nutrition*, 64(6), 633-645.
- Sen AK, Gupta KN (1982) Agro-ecological regions of western UP. *J Arid Environ*. 5:221–224
- Sharma, J.N., Chohan, G.S., Vij, V.K. and Monga, P.K. (1992). Effect of spacing on growth, yield and quality of Kinnow mandarin. *Indian J. Hort.* 49: 158-64.
- Sharma, K., & Singh, R. (2020). Effect of Organic and Inorganic Fertilization on the Growth and Yield of Kinnow Mandarin. *Journal of Plant Nutrition*, 43(10), 1460-1473.
- Singh, R., & Verma, K. (2017). Economic analysis of efficient agricultural practices in citrus production. *Agricultural Economics Research*, 56(3), 221-235.
- Singh, R., Singh, S., Singh, V., and Kuldip, H. K. (2024) Impact of nano-nitrogenous fertilization on productivity and profitability of Barley (*Hordeum vulgare* L.). *The Journal of Rural and Agricultural Research* Volume 24 No. 1, 110-114 (2024), DOI: 10.13140/RG.2.2.16387.72480.
- Singh, V., Chhonker, D., and Singh, R. (2022). Effect of Phosphorus Levels on Growth and Yield of Kabuli Chickpea (*Cicer kabulium* L.) Varieties. *The Journal of Rural and Agricultural Research* Vol. 22 No. 2, 108-111.
- Singh, V., Kumar, H. and Sonia (2023). Influence of Climate Change on Indian Agriculture” of published book entitled “Advances in Agriculture Science (Volume–44) having ISBN-978-93-5570-856-4.