

Effect of particle Size and Concentration of ZnO Nanoparticles on Growth, Yield and Seedling parameters in Buckwheat (*Fagopyrum esculentum* L.)

Abstract

The experiment was conducted in both field and lab conditions to study the Effect of Particle Size and Concentration of ZnO Nanoparticles on Growth, Yield and Seedling Parameters in Buckwheat (*Fagopyrum esculentum* L.) Under Late Sowing Conditions in Eastern Uttar Pradesh Region the lab experiment was performed in CRD with four different size ranges of zinc oxide Nanoparticles (20-30nm, 40-50nm, 60-80nm, 80-100nm) each of three concentrations (30PPM, 60PPM, 80PPM). The field experiment was performed during *Rabi* season of 2022-2023 in Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj Uttar Pradesh with a control was laid out in factorial RBD with 3 Replications. The results revealed that T1(20-30nm, 30PPM for 8hrs) shows greater Performance among parameters viz Plant height 30 DAS (23.45), Plant height 80 DAS (91.36), Days to first flowering (26.11), Number of Branches (4.43), Seeds per Plant (142.39), Seed yield per plot (11.87), Seed yield per hectare (1.19), Economic Yield (11.87), Biological Yield (32.76). and Harvest Index (36.23) and T3(20-30nm, 80PPM for 8hrs) exhibit greater in Leaf surface area (22.37), Test Weight (27.34). The lab results revealed the treatment T1(20-30nm, 30PPM for 8hrs) showing greater performance in Germination Percentage (95.63), Root Length (14.57), Seedling Length (24.76), Vigor Index-1 (2369.25), Electric conductivity (0.327), Protein content (13.59), Seed metabolic efficiency and T3 (20-30nm, 80PPM for 8hrs) shows greater Performance in Shoot length (11.57), Fresh Weight (2.19), Dry Weight (0.80), Vigor Index-2 (72.67), Seed Density (0.8), Chlorophyll Content (2.84), Dehydrogenize Activity (0.996).

Key words: Buckwheat, Zinc Oxide, Yield, Chlorophyll content, Protein content.

1. Introduction

Buckwheat (*Fagopyrum esculentum* L.), Chromosome Number $2n = 16$ is a Cross pollinated annual crop that belongs to the family Polygonaceae. It is grown as a cover crop and for its seeds, which resemble grains. Buckwheat, despite its name, is not related to wheat and is not a grain at all. The names boc from bechnut and wheat from wheat were combined to form the term "buckwheat" (Robinson, 1980). Because the plant's fruit resembled that of the bechnut, the word "beech" was

invented to describe it. Due to its function and chemical makeup resembling that of traditional cereals, it is a member of a dietary category known as pseudo cereals (**Campbell, 1997**). Buckwheat, a pseudo-cereal that grows in easy conditions, has the potential to become an important source of nutrition for the "**Starving World**" (**Leder et al., 2009**). Buckwheat is quite sensitive to low temperatures, while it tolerates hot temperatures relatively well (**Germ and Gabreski 2016**). Buckwheat grows best at temperatures ranging from 18 to 23 degrees Celsius. During the spring and autumn cultivation seasons, the average temperature in Korea is 7.3 °C and 25 °C, respectively (**Jung et al. 2015**). Reduced temperatures affect fertilization processes, resulting in reduced grain output in the spring season (**Farooq et al. 2016**).

Germplasm was morphologically characterized based on stem color, seed coat color, and seed morphology. Green, light Green, Pale Red, Crimson, and Dark Red Stem colors were noted. White, Grey, Pale Brown, Dark Brown, and Black Seed Coat color were noted. Documented seed shape for oval, egg-shaped (ovate), Triangular-type, and winged-form seeds. The grains are rich in polyphenols e.g., rutin and catechins which are reported to have antioxidant activity (**Morishita et al., 2007**). Buckwheat flowers are perfect but incomplete. They have no petals, but the calyx has the appearance of petals. Flowers occur in compact racemes, either terminally on the main stem or on branches from the axil of leaves.

Zinc oxide Nanoparticles have lately been employed and investigated for their impact on plant development in Zn-Nano formulations. Seed, foliar, and soil treatments are used to apply ZnO Nanoparticles. Among these applications, seed invigoration procedures like as seed priming and coating with ZnO Nanoparticles are regarded as relatively inexpensive and ecologically safe. Seed treatments that comprise the addition of ZnO Nanoparticles at the required concentrations result in ZnO Nanoparticles absorption by the seed coat, followed by their penetration and migration through seed tissue layers, which enhances germination and crop biomass. As a result, seed priming and seed coating are a targeted delivery approach for increasing the availability of micronutrients and other elements during the early growth of plants. Cultivation practices, as well as revolutionary seed invigoration techniques and precision farming, can boost seed germination potential, seedling vigor, photosynthesis, and reproductive development, resulting in increased production. Recent Nano-priming, a revolutionary approach of seed priming using designed Nanoparticles, has earned recognition for its unique physiochemical properties in boosting crop output and protection.

Globally, it has been used as Nano-fertilizer especially in Zn-deficit areas in order to enhance plant growth and develop (Sabir *et al.*, 2020). The addition of Nanoparticles may ameliorate nutrient deficiency and improve crop production (Dimkpa *et al.*, 2017). Recent studies have shown a positive impact of ZnO Nanoparticles on plant growth and physiology due to their wide spread applications in agriculture sector.

Objectives :

1. To evaluate the Effect of ZnO Nanoparticles on Growth and Yield parameters of Buckwheat.
2. To find out the Effect of ZnO Nanoparticles on Seedling parameters and Chlorophyll Content, Protein Content and Dehydrogenase Activity of Buckwheat

2. Methods and materials

The experiment was conducted during the Rabi Season of 2022 at field experimentation centre of Department of Genetics and Plant Breeding, Naini Agriculture Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh. The experimentation site is situated 98m above sea level at 25.570N latitude and 81.560N longitude. Buckwheat variety “IC-107575” sown at 25 X 15 cms which were treated with treatments of different concentrations of ZnO nanoparticles of different sizes.. The 13 treatment combinations along with untreated control were employed in the present study. The details of the treatment combinations, their dosage and the duration of the treatment were, T₀- Control, T₁- ZnO @20-30 nm of 30 PPM, T₂- @20-30 nm of 60 PPM, T₃-@20-30 nm of 80 PPM, T₄- @40-50 nm of 30 PPM, T₅-@40-50 nm of 60 PPM, T₆-@40-50 nm of 80 PPM, T₇-@60-80 nm of 30 PPM, T₈- @60-80nm of 60 PPM, T₉-@60-80 nm of 80 PPM, T₁₀-@80-100 nm of 30 PPM, T₁₁-@80-100 nm of 60 PPM, T₁₂-@80-100 nm of 80 PPM For 8 hours.

2.2 Methodology:

• ZnO Nanoparticles suspension preparation

ZnO Nanoparticles used in this study were purchased from Nano research Lab PVT Limited. Sizes of Nanoparticles are 20-30nm, 40-50nm, 60-80nm, 80-100nm Nanoparticles were directly

suspended in de-ionized water and dispersed ultrasonic vibration (100W, 40 KHz) for 10 minutes at lab of Department of Molecular and cellular Engineering, SHUATS. Small magnetic bars were placed in the suspension for stirring through Ultra sonicator to avoid aggregation of the particles and several suspensions of concentration range up to maximum possible limit were tried for uniform particle dispersion, stability and clear suspension (trial and error method) 3 concentrations of each size Nanoparticles (30, 60, 80 PPM) were selected for their evaluation in Buck wheat seed germination and growth study.

3. Result and Discussion

- **Analysis of Variance in Field conditions**

Analysis of Variance for Effect of particle size and concentration of Zinc oxide Nanoparticles on Growth, Yield parameters in Buckwheat (*Fagopyrum esculentum* L.) under late sowing conditions in Eastern Uttar Pradesh region. In Table 1.

- **Analysis of Variance in Seedling Parameters**

The Analysis of Variance for Effect of particle size and concentration of Zinc oxide Nanoparticles on Seedling parameters, Chlorophyll Content, protein Content, Dehydrogenase Activity in Buckwheat (*Fagopyrum esculentum* L.) under late sowing conditions in Eastern Uttar Pradesh region. In Table 2.

Table: 1 Analysis of variance for different characters in Buckwheat under field conditions

S.No	Characters	Mean sum of squares		
		Treatments (d.f = 12)	Replications (d.f = 2)	Error (d.f = 24)
01	Plant Height at 30 DAS	3.53*	0.21	0.35
02	Plant Height at 85 DAS	18.52*	5.21	5.59
03	Days to 50% Flowering	4.11*	0.45	0.62
04	Number of Branches	0.79*	0.00	0.01
05	Leaf Surface Area	8.46*	0.02	0.19
06	Seeds per Plant	622.29*	21.23	10.43
07	Seed Yield per Plot	14.75*	0.14	0.05
08	Seed Yield per ha	0.15*	0.00	0.00
09	Test Weight	9.15*	0.02	0.50
10	Economic Yield	14.75*	0.14	0.05
11	Biological Yield	65.72*	1.22	0.49
12	Harvest Index	31.76*	1.21	1.28

Table: 2 Analysis of variance for different characters in Buckwheat under laboratory conditions

S.No	Characters	Mean Sum Of Squares (MSS)	
		Treatments (d.f = 12)	Error (d.f = 24)
01	Germination percent	34.5*	3.07
02	Root Length	7.63*	0.067
03	Shoot length	4.79*	2.20
04	Seedling length	9.5*	2.25
05	Fresh Weight of seedling	0.3*	0.07
06	Dry Weight of seedling	0.06*	0.01
07	Seedling Vigor Index 1	156025.15*	16460.61
08	Seedling Vigor Index 2	2657.13*	60.73
09	Chlorophyll Content	1.192*	0.003
10	Protein Content	7.16*	0.036
11	Dehydrogenize activity	0.077*	0.00

*indicates significance at 5%

3.1 Physiological Observations:

At 30 DAS significantly higher Plant Height was observed in T₁ seeds treated with ZnO Nanoparticles of 20-30nm (30ppm for 8hrs) with 23.45cm and the lowest was recorded in T₀ (20.16 cm). At 85 DAS Plant Height was significantly higher in (T₁) seeds treated with ZnO Nanoparticles of 20-30nm (30ppm for 8hrs) with 91.36cm, Followed by T₄ is 90.89 cm. Lowest Plant Height was observed in T₀ with 84.14cm. Days to First Flowering that was recorded during the experimental crop growth, showing a significant better performance in the seeds that are primed with ZnO Nanoparticles of size 20-30 nm(30 PPM for 8hrs) T₁ with 26.11 DAS Showig least DAS for 50% flowering and followed by T₆ with 26.47 DAS. A lowest day to 50% Flowering was observed in T₀ with 29.71 DAS. From The present data on Leaf surface area that was recorded during the experimental crop growth, showing a significant higher leaf surface area in the seeds that are primed with ZnO Nanoparticles of size 20-30 nm(80 PPM for 8hrs) T₃ with 22.37nm² that is followed by T₆ with 22.11nm². Lowest Leaf Surface area was observed in T₀ with 17.72nm². Number of Primary Branches that was recorded during the experimental crop growth, showing a significant high number of branches in the seeds that are primed with ZnO Nanoparticles of size 20-30 nm(30 PPM for 8hrs) T₁ with 4.43 branches and followed by T₄ with 4.28 branches . Lowest number of branches was observed in T₀ with 2.84 branches. Number of seed per plant that was recorded during the experimental crop growth, shows a significant high number of seeds per plant in T₁ with ZnO Nanoparticles of size 20-30 nm(30 PPM for 8hrs) with 142 seeds, followed by T₄ with 139 seeds. Lowest number of seeds per plant was observed in T₀ with 101 seeds. Seed yield per plot that was recorded shows a significant high Performance in T₁ with ZnO Nanoparticles of size 20-30 nm(30 PPM for 8hrs) with 11.87g that is followed by T₄ with 11.28g. Lowest seed yield per plot was observed in T₀ with 5.54g. Seed Yield per Hectare that was recorded showed a significant high performance in T₁ with ZnO Nanoparticles of size 20-30 nm(30 PPM for 8hrs) with 1.19 t/ha that is followed by T₄ with 1.13 t/ha. Lowest seed yield per hectare was observed in T₀ with 0.55 t/ha. Test Weight that was recorded showing a significant high Performance in the seeds that are primed with ZnO Nanoparticles of size 20-30 nm(80 PPM for 8hrs) T₃ with 27.34g that is followed by T₆ with 27.19 g . Lowest Test Weight was observed in T₀ with 22.49g. ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) T₁ with 11.87g That is followed by T₄ with 11.28 g. The lowest Economic Yield was obtained in T₀ with 5.54g. Biological Yield was obtained in T₁ with ZnO Nanoparticles of size 20-

30nm (30PPM for 8hrs) with 32.76 g and that is followed by T₄ with 32.14 g and T₀ with 19.33 g was the lowest among all other treatments. Harvest Index was obtained in T₁ with ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) with 36.23% and that is followed by T₄ with 35.09 % and T₀ with 28.41% was the lowest among all other treatments.

Table 3: Mean performance of Effect of particle size and concentration of Zinc oxide Nanoparticles on growth, yield parameters in buckwheat

Treatments	PH 85 DAS	DDF	LSA	NPB	NSP	SYP	SYH	TW	EY	BY	HI
T ₀	84.14	29.71	17.72	2.84	101.28	5.54	0.55	22.49	5.54	19.33	28.41
T ₁	91.36	26.11	21.79	4.43	142.39	11.87	1.19	25.31	11.87	32.76	36.23
T ₂	89.49	26.68	20.83	3.84	127.59	10.45	1.05	26.83	10.45	30.77	33.98
T ₃	87.16	27.83	22.37	3.31	109.35	7.14	0.71	27.34	7.14	25.17	28.51
T ₄	90.89	28.14	21.38	4.28	139.54	11.28	1.13	24.46	11.28	32.14	35.09
T ₅	88.61	26.87	20.64	3.67	121.73	9.67	0.97	26.58	9.67	28.75	33.64
T ₆	85.73	26.47	22.11	3.19	106.68	6.84	0.68	27.19	6.84	23.75	28.80
T ₇	90.46	29.12	18.76	4.11	135.38	11.09	1.11	23.15	11.09	31.64	35.05
T ₈	88.24	28.46	20.37	3.51	117.46	9.25	0.93	23.87	9.25	27.66	33.44
T ₉	85.26	27.19	18.12	3.06	105.73	6.39	0.64	26.17	6.39	21.24	30.08
T ₁₀	90.16	29.38	18.57	3.97	130.24	10.73	1.07	22.64	10.73	31.12	34.50
T ₁₁	87.73	28.74	19.43	3.43	111.29	8.43	0.84	23.69	8.43	26.32	32.06
T ₁₂	84.43	27.58	17.81	2.98	103.21	6.04	0.60	25.67	6.04	20.63	29.32
F Test	S	S	S	S	S	S	S	S	S	S	S
S.E.M (±)	1.37	0.45	0.05	0.05	1.86	0.12	0.01	0.41	0.12	0.4	0.56
CD 5%	3.99	1.32	0.14	0.14	5.44	0.36	0.04	1.19	0.36	1.17	1.63

DAS: Days after sowing; PH: Plant height; LSA: Leaf Surface Area; NPB: Number of Primary branches; DDF: Days to First flowering; NSP: Number of seeds per plant; SYP: Seed Yield per plot; SYH: Seed yield Per Hectare; TW: Test Weight; BY: Biological yield; HI: Harvest Index

3.2 Seedling Observations:

Germination percent shows variation significantly among all treatments. T₁ seeds treated with ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) Showed higher germination percent than other treatments (95.63%) and Followed by T₄ with (94.57%). The lowest germination percent was observed in (T₀) with 85.74%. Root Length shows variation significantly among all treatments. T₁ seeds treated with ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) Showed higher Root Length than other treatments (14.57cm) and Followed by T₄ with (14.25cm). The lowest Root Length was observed in T₀ with (10.69cm). Shoot length shows variation significantly among all treatments. T₃ seeds treated with ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) Showed higher shoot length than other treatments (11.57cm) and Followed by T₆ with (11.41cm). The lowest shoot length was observed in T₀ with (8.53cm). Seedling length shows variation significantly among all treatments. T₁ seeds treated with ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) Showed higher seedling length than other treatments (24.26cm) and Followed by T₄ with (24.18cm). The lowest seedling length was observed in T₀ with (19.22cm). Higher Seedling Fresh Weight recorded in the seeds treated with ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) of T₃ with 2.39g than other treatments that is followed by T₆ with 2.11g. The lowest seedling fresh Weight was observed in T₀ with (1.25g). Higher Seedling Dry Weight recorded in the seeds treated with ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) of T₃ with 0.80g than other treatments that is Followed by T₆ with 0.79g. The lowest seedling Dry Weight was observed in T₀ with (0.43g). Seedling Vigor Index -I was observed in seeds treated with (T₁) ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) with 2369.25. and that is followed by T₄ with 2286.31. The lowest Vigor Index -I was recorded in T₀ with 1648.02. Seedling Vigor Index-II was observed in seeds treated with (T₃) ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) with 72.67 and that is followed by T₆ with 71.07. The lowest Vigor Index -II was recorded in T₀ with 37.23. Seed Density is shown that ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) (T₃) is significantly higher (0.8) over all other treatments and that is followed by T₁, T₂ and T₄ with Seed Density (0.7). The lowest Seed Density was observed in T₀ (0.2). From the data that was recorded, Seeds primed with treatment T₁ of ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) shows significantly higher electrical conductivity with (0.327) that is followed by T₆ with (0.259). The lowest electrical conductivity was recorded in T₀ with (0.153). From the data

that was recorded, Seeds primed with treatment T₁ of ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) had shown higher Seed metabolic efficiency with (1.01) and that is followed by T₃ with (0.97). The lowest seed metabolic efficiency was recorded in T₀ with (0.84).

Table 4: Mean performance of Effect of particle size and concentration of Zinc oxide Nanoparticles on Seedling parameters, Chlorophyll Content, Protein Content, Dehydrogenize activity in buckwheat

Treatments	GP	RL	SL	Se.L	FW	DW	VI-I	VI-II	SD	EC	SME	CC	PC	DE
T ₀	85.74	10.69	8.53	19.22	1.25	0.43	1648.02	37.23	0.2	0.153	0.84	1.21	9.61	0.593
T ₁	95.63	14.57	10.19	24.76	1.85	0.60	2369.25	58.16	0.7	0.327	1.01	2.65	13.59	0.795
T ₂	92.68	13.41	11.35	24.76	2.09	0.76	2295.35	70.58	0.7	0.199	0.97	2.31	12.74	0.935
T ₃	90.29	11.63	11.57	23.20	2.19	0.80	2096.00	72.67	0.8	0.227	0.974	2.84	11.21	0.996
T ₄	94.57	14.25	9.92	24.18	1.73	0.59	2286.31	56.05	0.7	0.164	0.931	2.43	13.35	0.744
T ₅	91.87	12.89	11.16	24.05	2.05	0.71	2208.19	65.20	0.6	0.208	0.963	2.26	12.48	0.887
T ₆	89.62	11.35	11.41	22.76	2.11	0.79	2039.39	71.07	0.8	0.259	0.88	2.70	10.93	0.968
T ₇	93.68	14.09	8.75	22.84	1.56	0.49	2143.52	45.89	0.5	0.186	0.94	1.81	13.11	0.634
T ₈	91.38	12.60	9.75	22.35	1.67	0.56	2043.62	51.23	0.6	0.214	0.942	2.18	11.89	0.705
T ₉	87.74	10.97	10.86	21.835	1.96	0.67	1915.51	59.26	0.4	0.273	0.92	1.42	10.57	0.846
T ₁₀	93.27	13.76	8.61	22.37	1.39	0.44	2080.03	41.80	0.4	0.193	0.91	1.5	12.98	0.611
T ₁₁	90.76	12.35	9.41	21.76	1.63	0.50	1975.94	46.01	0.5	0.22	0.85	2.02	11.46	0.657
T ₁₂	87.48	10.83	10.48	21.31	1.92	0.63	1863.54	55.27	0.3	0.301	0.85	1.35	9.82	0.823
F Test	S	S	S	S	S	S	S	S	S	S	S	S	S	S
S.E.M (±)	0.87	0.13	0.74	0.75	0.13	0.05	64.14	4.95	0.05	0.015	0.014	0.028	0.095	0.007
CD (p=0.05)	2.5	0.37	2.13	2.15	0.38	0.15	184.19	14.23	0.16	0.12	0.11	0.08	0.273	0.021

GP-Germination Percentage; RL-Root length; SL-Shoot length; SeL-Seedling length; FW-Fresh weight ; DW-Dryweight; VI-I-Vigour index-I; VI-II-Vigour index-II; SD-Seed density; SME- Seed metabolic efficiency ;EC-Electrical conductivity

3.3 Biochemical analysis:

Chlorophyll Content which was recorded as significant higher in seeds Primed with (T₃) ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) with 2.84 %. However that is followed by T₆ with (2.64%).The lowest chlorophyll Content was observed in T₀ with (1.21%).Protein Content which was recorded as significant higher in seeds treated with (T₁) ZnO Nanoparticles of size 20-30nm (30PPM for 8hrs) with 13.59. However that is followed by T₄ with protein Content (13.35). The Lowest protein Content was observed in T₀with (9.61).Dehydrogenize Activity. It is clear that

seeds treated with ZnO Nanoparticles of size 20-30nm (80PPM for 8hrs) (T₃) performed better and is significantly higher over all other treatments with 0.996 and that is followed by T₆ with 0.968 respectively and lowest dehydrogenize activity was recorded in T₀ with 0.593.

Table 5: Mean performance of Effect of particle size and concentration of Zinc oxide Nanoparticles on Chlorophyll Content, Protein Content, Dehydrogenize activity in buckwheat

Treatments	CC	PC	DE
T ₀	1.21	9.61	0.593
T ₁	2.65	13.59	0.795
T ₂	2.31	12.74	0.935
T ₃	2.84	11.21	0.996
T ₄	2.43	13.35	0.744
T ₅	2.26	12.48	0.887
T ₆	2.70	10.93	0.968
T ₇	1.81	13.11	0.634
T ₈	2.18	11.89	0.705
T ₉	1.42	10.57	0.846
T ₁₀	1.5	12.98	0.611
T ₁₁	2.02	11.46	0.657
T ₁₂	1.35	9.82	0.823
F Test	S	S	S
S.EM (±)	0.028	0.095	0.007
CD (p=0.05)	0.08	0.273	0.021

CC-Chlorophyll Content ; DE-Dehydrogenize Activity ; PC-Protein Content

Discussion

In present investigation was conducted to know the Effect of particle Size and Concentration of ZnO Nanoparticles on Growth, Yield and Seedling parameters in Buckwheat crop as experimental results proved the beneficial effect of Nanoparticles, some logical explanation has been discussed in detail in favor to particle size and concentration . Buckwheat seeds treated with ZnO Nanoparticles Buckwheat seeds to germination test under laboratory conditions along with untreated seeds (control). The results pertaining to seed germination and early seedling growth of seeds treated with ZnO Nanoparticles are given in table 4.The seeds treated with ZnO Nanoparticles treated Buckwheat

seeds showed significantly more seed germination and seedling performance with respect radical and shoot growth as compared to control (untreated) seed

Tables 3 and 4 show that ZnO Nanoparticles sized 20 to 30, 40 to 50, 60 to 80, and 80 to 100 nm significantly improved vegetative growth, triggered early flowering, and seed output in treated Buckwheat plants.

Conclusion

- Among various treatments T₁ in which seeds are treated with 30PPM of size 20nm-30nm performed better in terms of Plant Height-85DAS, Days to first flowering, Number of branches/Plant, Leaf surface area, Number of Seeds/plant, Seed yield per Plot, Seed yield Per Hectare, Economic Yield, Biological Yield, Harvest Index and seeds treated with 80PPM of size 20nm-30nm (T₃) performed better in terms of Leaf surface area and Test Weight. When compared with the Control (T₀) which has shown lowest performance in all parameters. Among all the treatments T₁ in which seeds treated with 30PPM of Zinc oxide (ZnO) of 20-30nm size for 8hrs has shown higher level of Germination Percentage, Root Length, Seedling length, Vigor Index-1, Electric conductivity, seed metabolic efficiency, Protein Content, and Seeds Treated With 80PPM of Size 20nm-30nm (T₃) Shown high level of Shoot Length, Fresh Weight, Dry Weight, Vigor Index-II Seed Density, Chlorophyll Content and Dehydrogenase Activity. when compared with the Control(T₀) which has shown lowest performance in all parameters

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

References

Abd El-Rahman, K.A., E.A. Teama., F.M. Fathy and Abdelaziz (2020). Effect of Foliar Spray by Different Zinc Oxide Nanoparticles Concentrations at Various Growth Stages on Growth Yield and its Components of Maize. *Assiut J.Agric.Sci.* **51**(2);43-53.

Afshar, i., Akbar, r. and Minoo, S., (2014), Comparison of the effect of spraying different amounts of Nano Zinc Oxide and Zinc Oxide .*Int. Jour. Adv. Biol. Biom. Res.*, 2(4);318-325.

Ahmed, R., Uddin, M. K., Quddus, M. A., Samad, M. Y. A., Hossain, M. M., & Haque, A. N. A. (2023). Impact of foliar application of zinc and zinc oxide nanoparticles on growth, yield, nutrient uptake and quality of tomato. *Horticulturae*, 9(2), 162.

Amita Hajra and Naba Kumar Mondal (2017). Effect of ZnO and TiO₂ Nanoparticles on germination, Biochemical and morphoanatomical attributes of *Cicer arietinum* L. *Energ.Ecol. Environ.* **2**(4);277-288. DOI: 10.1007/s4097-017-0059-6.

Alloway BJ (2008). Zinc in soils and crop nutrition. Brussels, Belgium: Online book published by *the International Zinc Association*; 2008.

Boonyanitipong P., Kositup, B., Kumar, P., Barauh, B. and Dutta, J. (2011). Toxicity of ZnO and TiO₂ Nanoparticles on germination Rice seed (*Oryza sativa* L.) *International Journal of Bioscience, Biochemistry and Bioinformatics.* **1**(4)282-285.

Campbell, C.G. (1997). *Buckwheat: Fagopyrum esculentum Moench Biodiversity International.* **19**(6): 22-29.

Datir, R.B., Apparao B.J. and Laware, S.L. 2012. Application of amino acid chelated micronutrients for enhancing growth and productivity in chilli (*Capsicum annum*L.). *Plant Sciences Feed.* **2** (7): 100-105.

Datir, R.B., Laware, S.L., and Apparao, B.J. 2010. Effect of Organically Chelated Micronutrients on Growth and Productivity in Okra. *Asian J. Exp. Biol. Sci. Spl.* 115-117.

Dimpak.O., Singh, U., Bindraban, P.S., ADISA, I.O., Elmer, W.H., Gardea-Torresdet,J.L., (2019). Addition- Omission of Zinc, Copper, and Boron Nano and Bulk particles demonstrate element and size specific response of soyabean to micronutrients expose. *Science Total Environment.* 665,606-616.

Farooq, S., Rehman, R.U., Pirzadah, T.B., Malik, B., Dar, F.A. and Tahir, I. (2016). Cultivation, agronomic practices, and growth performance of buckwheat. In *Molecular breeding and nutritional aspects of buckwheat*. 22(2): 299-319.

Harish Kumar, K., Aditya, S., and Savalgi V.P, (2021). Evaluation of foliar application of zinc Nanoparticles on growth and yield parameters of maize (*Zea mays* L.) grown under greenhouse conditions. *International Journal of Chemical Studies*.9(1);1464-1467.

Iziy, E., Majd, A., Vaezi-Kakhki, M.R., Nejadstarrari, T., Kazemi Noureini, S. (2019). Effects of zinc oxide Nanoparticles on enzymatic and non enzymatic antioxidant content, germination, and biochemical and ultra structural cell characteristics of *Portulaca oleracea* L. *Acta Soc Bot Pol.* 88(4):3639. 17.

Kisan, B.,Shruthi, H., Sharanagouda, H., Revanappa, S.B. and Pramod, N.K.(2015). Effect of Nano Zinc Oxide on the Leaf Physical and Nutritional quality of soinach. *Agrotechnology*, 5:135-134.

Kolenčík, M., Ernst, D., Komár, M., Urík, M., Šebesta, M., Ďurišová, L., ... & Aydın, E. (2022). Effects of foliar application of ZnO nanoparticles on lentil production, stress level and nutritional seed quality under field conditions. *Nanomaterials*, 12(3), 310.

Kumar, B., & Dhaliwal, S. S. (2021). Zinc bio fortification of dual-purpose cowpea (*vigna unguiculata* L.) walp.] for enhancing the productivity and nutritional quality in a semi-arid region of India. *Archives of Agronomy and Soil Science*, 68(8), 1034–1048.

Laware, S.L. and Raskar, S. (2014). Influence of Zinc Oxide Nanoparticles on Growth, Flowering, Seed productivity in Onion, *International Journal of Current Microbiology and Applied Science*. 3(7):874-881.

Leder, I. (2009). Buckwheat, amaranth, and other pseudocereal plants. *Encyclopedia of Life Support Systems. 1st ed. Ramsey: EOLSS Publishers Co Ltd*, 6(2): 1-17.

Morishita, T., Yamaguchi, H. and Degi, K. (2007). The contribution of polyphenols to antioxidative activity in common buckwheat and tartary buckwheat grain. *Plant production science*. 10(1): 99-104.

Nadi, E., Ayneband and Mojaddam, M. (2013). Effect of Nano Iron chelate fertilizer on Grain yield, Protein percent and Chlorophyll content of Fababean (*Vicia faba* L.) *Int. J. Biosci.*, 3(9):267-272.

Prasad T.N.V.K.V., P.Sudhakar, Y.,Sreenivasulu, P., Latha,V. Munaswamy, K.RajaReddy, T.S.Sreeprasad, P.R.Sajanlal, and T.Pradeep. 2012. Effect of nano scale zinc oxide particles on the germination, growth and yield of peanut. *Journal of Plant Nutrition*. 35,905927.

Ramesh raddy., Mahesh Salimath., K.N. Geetha and Shankar, A.G. (2018). ZnO Nanoparticle Improves Maize Growth, Yield and Seed Zinc under High Soil pH Condition. *Int.J.Curr.Microbiol.App.Sci*. 7(12): 1593- 1601.

Poornima, R., and Koti, R.V. (2019). Effect of nano zinc oxide on growth, yield and grain zinc content of sorghum (*Sorghum bicolor* L). *Journal of Pharmacognosy and Phytochemistry*. 8(4): 727-731.

Racuciu, M., and Creanga, D.2007. TMA-OH coated magnetic Nanoparticles internalized in vegetal tissues. *Romanian Journal of Physics*. 52,395402.

Raskar, S.V. and Laware, S.L.2014. Effect of Zinc oxide Nanoparticles on cytology and seed germination in onion. *Int. J. Curr. Microbiol. App. Sci*. 3(2): 467-473.

Rengel,Z., and Graham, R.D.1995. Importance of seed zinc content for wheat growth on zinc-deficient soil. *I. Vegetative growth*. *Plant and Soil*. 173,259-266.

Robinson RG (1980) The buckwheat crop in Minnesota. In: *Agricultural Experiment station Bulletin* 539, University of Minnesota St.Paul

Sadak, M.S., Bakry, B.A. (2020) Zinc Oxide and Nano ZnO oxide effect on growth, some biochemical aspects, yield quantity, ad quality of flax (*Linum uitatissimum* L.) in absence and presence of Compost under sandy soil. *Bull Nat ResCent* 44, 98.

Satdev, V.J., Zinzala, Bharat N Chavda and Lokesh Kumar Saini. (2019). Effect of nano ZnO on growth and yield of sweet corn under South Gujarat condition. *International Journal of Chemical Studies*. **8**(1); 2020-2023.

Sabir, S., Zahoor, M.A., Waseem, M., Siddique, M.H., Shafique, M., Imran, M., Hayat, S., Malik, I.R., Muzammil, S., (2020). Biosynthesis of ZnO nanoparticles using bacillus subtilis: characterization and nutritive significance for promoting plant growth in *Zea mays* L. *Journal of nanotechnology*. **18**(3): 1-9.

Salah M. H. Gowayed (2017). Impact of Zinc Oxide Nanoparticles on Germination and Antioxidant system of Maize (*Zea mays* L.) Seedling under Cadmium Stress. *Journal of Plant Production Science*, **6**(1):1-11.

Šebesta, M., Kolenčík, M., Sunil, B. R., Illa, R., Mosnáček, J., Ingle, A. P., & Urík, M. (2021). Field application of ZnO and TiO₂ nanoparticles on agricultural plants. *Agronomy*, **11**(11), 2281.

Singh. M. D., Jayadeva, H. M., Chirag, G and Mohan, M.H. (2017). Effect of Nano ZnO Particles on seedling growth of Maize (*Zea mays* L.) in Germination Paper test *Interational Journal of Microbiology Research*. **9**(5);897-898.

Srivastav, A., Ganjewala, D., Singhal, R. K., Rajput, V. D., Minkina, T., Voloshina, M., ... & Shrivastava, M. (2021). Effect of ZnO nanoparticles on growth and biochemical responses of wheat and maize. *Plants*, **10**(12), 2556.

Tarafdar, J. & Sharma, Shikha & Raliya, Ramesh. (2013). Nanotechnology: Inter disciplinary science of applications. *African Journal of Biotechnology*. **12**. 219-226. **10.5897/AJB12**.

Thounaojam, T. C., Meetei, T. T., Devi, Y. B., Panda, S. K., & Upadhyaya, H. (2021). Zinc oxide nanoparticles (ZnO-NPs): a promising nanoparticle in renovating plant science. *Acta Physiologiae Plantarum*, **43**, 1-21.

Uma.V., Jayadeva, H., Rehamn,H. nad Kadalli, G.G. Influence of Nano Zinc Oxide on Yield and Economics of Maize (*Zea mays* L.). *Mysore.J.Agric.Sci.*, **53**(4); **44-48**.