

Effect of nitrogen and boron fertilization on yield and quality of Mustard (*Brassica juncea*) in Chitrakoot area, Madhya Pradesh, India

Abstract

A field experiment was conducted in year 2022 in Rabi season at Rajaula Agriculture farm, Mahatma Gandhi Chitrakoot Gramoday Vishwavidyalaya Chitrakoot, Satna (M.P.). Experiment was laid out in randomized block design with thirteen treatment combinations, varying levels of nano-urea and boron application, alongside a standard 100 % NPK as per RDF treatment as the control. Results showed significant differences in seed and oil yield across treatments. The highest seed yield of 1523.81 kg/ha was achieved with T₁₂ (½ RDN + two nano-urea sprays + 1.25 kg B), a notable improvement over the control (T₀) at 958.73 kg/ha. Oil content ranged from 13.1% to 35.6%, with the highest in T₁₂, which also produced the maximum oil yield of 542.48 kg/ha. The lowest oil content (13.1%) and oil yield (190.47 kg/ha) were observed in T₁₁ (½ RDN + one nano-urea spray + one water spray + 1.25 kg B). Treatments incorporating nano-urea and higher levels of boron showed the most pronounced increases in both oil content and yield, suggesting a synergistic effect of these nutrients on mustard productivity. Statistical analysis confirmed significant improvements in yield and oil quality with combined boron and nano-urea applications, indicating that T₁₂ is the optimal treatment for maximizing mustard production under similar agro-climatic conditions.

Key words: Nitrogen, Boron, Yield, Oil content, Oil yield, Mustard crop.

Introduction

Indian mustard is one of the most important edible oilseed *rabi* crop of North India commonly known as *Sarson*, *Rai* or *Laha*. It belongs to the family Brassicaceae and genus *Brassica*. The oil content in mustard seeds varies from 37-49 percent (Singh *et. al.*, 2023), the seeds are highly nutritive containing 38-57 % eruric acid, and 27 % oleic acid. The seed is used as a condiment in the preparation of pickles and for flavouring curries and vegetables. The oil is utilized for human consumption throughout the northern India for cooking purpose. This is a potential crop in winter (Rabi) season due to its wider adaptability and suitability to exploit residual moisture (Mukherjee, 2010). Oil and fats comprise a vital component of human diet as these are good source of energy and act as carriers of fat-soluble vitamins. Oil cake or meal has high nutritional values in animal diet. Seed owing to its high content of good quality protein. In general, 55g edible oil per day head is essential for human diet.

India ranks third in terms of area and production of rapeseed-mustard after Canada and China. Globally, the area and production of rapeseed-mustard is 36.81 million hectares and 72.61 million tonnes, respectively (**USDA, 2020**). Rapeseed mustard is the second most consumed edible oilseed crop in India, after soybean. India has 6.23 million hectares area under rapeseed mustard and 9.34 million tonnes production with average productivity of 1499 kg ha⁻¹, which is about three-fourth of the world's average productivity (1960 kg ha⁻¹) (**DAC & FW, 2020**). In the Madhya Pradesh, it is grown on an area 1038.15 thousand hectares with a production of 1.69 million tonnes (**Anonymous 2021-22**).

Urea is a rich source of nitrogen, an essential nutrient for plant growth. Nitrogen is a crucial component of chlorophyll, the green pigment in plants responsible for photosynthesis. Adequate nitrogen supply helps mustard plants produce more chlorophyll, leading to improved photosynthesis and overall plant growth. Mustard plants respond well to nitrogen fertilizers like urea during their vegetative stage (**Shorna et al., 2020**). This can lead to better light interception and more efficient utilization of sunlight. The application of urea at the right time and in the right amount can significantly increase mustard crop yield. One of the most common symptoms of nitrogen deficiency is the yellowing of older leaves, starting from the tips and progressing towards the base of the plant. Urea deficiency can lead to reduced plant height, fewer branches, and a generally stunted appearance. Mustard plants may fail to reach their full growth potential (**Iqbal et al., 2011**).

Nano fertilizers possess unique feature which enhance plant performance in terms of ultrahigh absorption, increase in production, rise in the leaves surface area. Beside the controlled released of nutrients contributes in preventing eutrophication and pollution in water resources. Replacement of traditional fertilizers by nano fertilizer is beneficial as upon application, it releases nutrients into the soil steadily and in a controlled way, thus, preventing the water pollution (**Naderi and Danesh Shahraki 2013, Moaveni and Kheiri, 2011**).

The use of nano fertilizers not only causes increased use efficiency through ultra-high absorption of the nutrients, increase in photosynthesis caused by expansion in surface area of the leaves (INIC 2009) but also reduce the toxicity generated due to over application in the soil as well as reduces the split application of fertilizer (**Naderi and Danesh Shahraki, 2013**). Nano fertilizers and nanocomposites can be used to control the release of nutrients from the fertilizer granules so as to improve the nutrient use efficiency while preventing the nutrient ions from either getting fixed or lost to the environment (**Subramanian et al. 2008**).

Boron is essential for the formation and stability of plant cell walls. It is involved in the cross-linking of pectin molecules, which helps maintain the structural integrity of plant cells.

In mustard plants, this is especially important for maintaining stem and seed pod strength. Adequate boron levels promote efficient pollination and higher seed set, ultimately contributing to increased yield. Boron aids in the uptake and translocation of other essential nutrients within the plant. It helps in the movement of calcium, which is important for cell division and overall plant growth (Sharma *et al.*, 2020).

Boron is unique among the essential mineral nutrients because it is the only element that is normally present in soil solution as a non- ionised molecule over the pH range suitable for plant growth. In soil solution, boron mainly exists as undissociated boric acid (H_3BO_3). Boric acid is the major form of boron in soils with H_2BO_3 being predominant only above pH 9.2. Boron occurs in aqueous solution as boric acid $B(OH)_3$, which is a weak monobasic acid that acts as an electron acceptor or as a Lewis acid (Rathore, 2015).

Materials And Method

The experiment was carried out at Rajaula Agriculture farm, Mahatma Gandhi Chitrakoot Gramoday Vishwavidyalaya Chitrakoot, Satna (M.P.) which lies in the semi- arid and sub-tropical region of Madhya Pradesh between 25.148° North latitude and 80.855° East longitude. The altitude of town is about 190-210 meter above mean sea level. Soil Characteristics; pH is soil 7.4, EC 0.34; OC 0.31; Total N 97.68; Available P 16.25; Available K 292.90 and available boron is 0.38. The field was prepared by ploughing with a tractor drawn disc plough by cross harrowing and planking. After preparation of land, the experiment was laid out as per treatment combinations, there were 39 plots and the gross size of each plot was 5.0 m x 4.0 m and the net plot size was 4.5 m x 3.5 m. FYM was applied @ 10 q ha^{-1} as basal dose. After the layout of experimental plot, the fertilizers were weighed and applied in the plots and thoroughly mixed with soil. As per the experimental recommended doses of Nitrogen, Phosphorus, Potassium were applied to assigned plots. Recommended dose of Nitrogen, Phosphorus and Potassium were applied through Urea, DAP and MOP (60:40:40 $kg\ ha^{-1}$) whereas boron was applied through borax (0, 0.5, 1.0, 1.25 $kg\ B\ ha^{-1}$). The seed was sown in line after making a narrow furrow with the help of pointed wooden stick at different row spacing. No irrigation was given to entire experimental field. The crop was harvested on 14th Feb., 2023 when it reached to its physiological maturity i.e. when the leaves were turned yellow and more than 70 % capsules were full matured to avoid shattering of the crop.

The 13 treatments combination of nutrient management practices. Experiment was laid out in Randomized Block Design with three replications. i.e. T0; control, T1; $\frac{1}{2}$ of RDN + (2 water spray + 0.0 $kg\ B$), T2; $\frac{1}{2}$ of RDN + (2 water spray + 0.0 $kg\ B$), T3; $\frac{1}{2}$ of RDN + (1st nano-urea

spray + 2nd nano-urea spray + 0.0 kg B), T4; ½ of RDN + (2 water spray + 0.5 kg B), T5; ½ of RDN + (1st nano-urea spray + 2nd water spray + 0.5 kg B), T6; ½ of RDN + (1st nano-urea spray + 2nd nano-urea spray + 0.5 kg B), T7; ½ of RDN + (2 water spray + 1.0 kg B), T8; ½ of RDN + (1st nano-urea spray + 2nd water spray + 1.0 kg B), T9; ½ of RDN + (1st nano-urea spray + 2nd nano-urea spray + 1.0 kg B) T10; ½ of RDN + (2 water spray + 1.25 kg B), T11; ½ of RDN + (1st nano-urea spray + 2nd water spray + 1.25 kg B), T12; ½ of RDN + (1st nano-urea spray + 2nd nano-urea spray + 1.25 kg B). Collected data and observations i.e. Oil content in seed (%), Oil Yield (kg ha⁻¹),

Data collection

Oil content in seed (%)

Oil content estimated by Soxhelt extraction method.

$$\text{Oil content in seed (\%)} = \frac{W_0}{W_s} \times 100$$

W_0 = weight of oil extracted in grams

W_s = weight of seed in grams

Oil Yield (kg ha⁻¹)

$$\text{Oil yield (kg ha}^{-1}\text{)} = \frac{\text{Oil content (\%)} \times \text{Seed yield (kg ha}^{-1}\text{)}}{100}$$

Statistical analysis: The growth parameters and yields were recorded and analyzed as per Gomez (1984) the tested at 5% level of significance to interpret the significant differences.

Result and discussion

Seed yield

The results revealed that the total seed yield (kg ha⁻¹) of mustard varied in between 958.73 to 1523.81 kg ha⁻¹ all the treatments were significantly superior to T₀ [100 5 NPK as per RDF]. The treatment combination T₁₂ [½ of RDN + (1st nano-urea spray + 2nd nano-urea spray + 1.25 kg B)] gave the maximum total seed yield (1523.81 kg ha⁻¹) followed by the treatment T₉ [½ of RDN + (1st nano-urea spray + 2nd nano-urea spray + 1.0 kg B)] with the value 1485.71 kg ha⁻¹. Minimum total seed yield (958.73 kg ha⁻¹) was found under the treatment T₀ [100 % NPK as per RDF].

Total seed yield increased significantly with the combined use of boron and 50 % nitrogen / urea spray. The minimum total seed yield was noted 1066.66 kg ha⁻¹ in the treatment 0 kg B ha⁻¹ + 50 % RDN + N₀ spray and maximum total seed yield 1523.80 kg ha⁻¹ was noted in the treatment 1.25 kg B ha⁻¹ + 50 % RDN + N₂ spray. Similar findings were reported by **Kumar *et al.* (2016), Sinha *et al.* (2022) and Kumar *et al.* (2022).**

Table-1: Effect of different treatment combination on number of total seed yield (kg ha⁻¹)

Treatment	Treatment Combination	Total seed yield (kg ha ⁻¹)
T ₀	100 % NPK as per RDF	958.73
T ₁	½ of RDN + (2 water spray + 0.0 kg B)	1009.52
T ₂	½ of RDN + (I st nano-urea spray + 2 nd water spray + 0.0 kg B)	1066.67
T ₃	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 0.0 kg B)	1155.55
T ₄	½ of RDN + (2 water spray + 0.5 kg B)	1117.46
T ₅	½ of RDN + (I st nano-urea spray + 2 nd water spray + 0.5 kg B)	1212.70
T ₆	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 0.5 kg B)	1346.03
T ₇	½ of RDN + (2 water spray + 1.0 kg B)	1295.24
T ₈	½ of RDN + (I st nano-urea spray + 2 nd water spray + 1.0 kg B)	1422.22
T ₉	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 1.0 kg B)	1485.71
T ₁₀	½ of RDN + (2 water spray + 1.25 kg B)	1384.13
T ₁₁	½ of RDN + (I st nano-urea spray + 2 nd water spray + 1.25 kg B)	1453.97
T ₁₂	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 1.25 kg B)	1523.81
SEm ±		3.64
C.D. (P=0.05)		10.75

Quality parameters

Oil Content (%)

The oil content ranged from 13.1 % to 35.6 % across the treatments. The lowest oil content (13.1 %) was recorded in treatment T₁₁, which combined half of the recommended dose of nitrogen (RDN) with one nano-urea spray, one water spray, and 1.25 kg of boron (B). The highest oil content (35.6 %) was observed in treatment T₁₂, which used half of the RDN with two nano-urea sprays and 1.25 kg B. Treatments with nano-urea sprays generally led to higher oil content, with T₉ and T₁₂ yielding particularly high results (35.0 % and 35.6 %, respectively).

Similar findings were reported by **Kumar *et al.*, (2014)**, **Yadav *et al.*, (2016)** and **Pandey *et al.*, (2022)**

Oil yield (kg ha⁻¹)

Oil yield ranged from 190.47 kg/ha to 542.48 kg/ha. T₁₂ achieved the highest oil yield at 542.48 kg/ha, indicating that two nano-urea sprays combined with 1.25 kg B and half of the RDN is the most effective combination for maximizing oil yield. The lowest yield was observed in T₁₁ (190.47 kg/ha), where one nano-urea spray, one water spray, and 1.25 kg B were applied. This treatment yielded lower both in oil content and oil yield. As with oil content, treatments incorporating nano-urea sprays and higher boron levels tended to produce higher oil yields. Similar findings were reported by **Sachan *et al.*, (2022)**, **Dhaliwal *et al.*, (2022)** and **Halim *et al.*, (2023)**

Table-2: Effect of different treatment combination on oil content and oil yield

Treatment	Treatment Combination	Oil content (%)	Oil yield (kg ha ⁻¹)
T ₀	100 % NPK as per RDF	31.1	298.17
T ₁	½ of RDN + (2 water spray + 0.0 kg B)	32.2	325.07
T ₂	½ of RDN + (I st nano-urea spray + 2 nd water spray + 0.0 kg B)	32.6	347.73
T ₃	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 0.0 kg B)	33.2	383.64
T ₄	½ of RDN + (2 water spray + 0.5 kg B)	32.9	367.64
T ₅	½ of RDN + (I st nano-urea spray + 2 nd water spray + 0.5 kg B)	33.5	406.25
T ₆	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 0.5 kg B)	34.1	459.00
T ₇	½ of RDN + (2 water spray + 1.0 kg B)	33.8	437.79
T ₈	½ of RDN + (I st nano-urea spray + 2 nd water spray + 1.0 kg B)	34.8	494.93
T ₉	½ of RDN + (I st nano-urea spray + 2 nd nano-urea spray + 1.0 kg B)	35.0	520.00
T ₁₀	½ of RDN + (2 water spray + 1.25 kg B)	34.5	477.52

T₁₁	½ of RDN + (1 st nano-urea spray + 2 nd water spray + 1.25 kg B)	13.1	190.47
T₁₂	½ of RDN + (1 st nano-urea spray + 2 nd nano-urea spray + 1.25 kg B)	35.6	542.48
SEm ±		0.12	2.06
C.D. (P=0.05)		0.35	6.06

Conclusion

The study highlights the beneficial effects of combining nano-urea and boron applications on mustard crop yield and oil quality in the Chitrakoot region. The treatment combination T₁₂ (½ RDN + two nano-urea sprays + 1.25 kg B) proved to be the most effective, significantly enhancing seed yield (1523.81 kg/ha) and oil yield (542.48 kg/ha) compared to the control treatment with 100 % NPK as per RDF. Results indicate that the combined application of boron and nano-urea not only boosts seed yield but also improves oil content, with T₁₂ achieving the highest oil percentage at 35.6 %. This study suggests that using nano-urea and boron synergistically can optimize nutrient efficiency, enhance mustard crop productivity, and improve oil quality, making it a promising nutrient management strategy for similar semi-arid regions.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1.

2.

3.

References

1. **Berger, K. C., & Truog, E. (1939).** Boron determination in soils and plants. *Industrial & Engineering Chemistry Analytical Edition*, 11(10), 540-545.
2. **Dhaliwal, S. S., Sharma, V., Shukla, A. K., Kaur, M., Verma, V., Sandhu, P. S., ... & Hossain, A. (2022).** Biofortification of oil quality, yield, and nutrient uptake in Indian mustard (*Brassica juncea* L.) by foliar application of boron and nitrogen. *Frontiers in Plant Science*, 13, 976391.
3. **Gomez, K. A. (1984).** Statistical procedures for agricultural research. *John NewYork: Wiley and Sons*.

4.

- **Halim, A., Paul, S. K., Sarkar, M. A. R., Rashid, M. H., Perveen, S., Mia, M. L., ... & Islam, A. M. (2023).** Field assessment of two micronutrients (zinc and boron) on the seed yield and oil content of mustard. *Seeds*, 2(1), 127-137.
5. **Hanway, J.J; and Heidel, H. (1952).** Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture* **54**: 1-31.
6. **Iqbal, N., Nazar, R., Syeed, S., Masood, A., & Khan, N. A. (2011).** Exogenously-sourced ethylene increases stomatal conductance, photosynthesis, and growth under optimal and deficient nitrogen fertilization in mustard. *Journal of experimental botany*, 62(14), 4955-4963.
7. **Jackson, M.L. (1973).** Soil chemical analysis. Prentice Hall of India Pvt. Ltd, New Delhi.
8. **Kumar, A., Singh, K., Verma, P., Singh, O., Panwar, A., Singh, T., Kumar, Y. and Raliya, R. (2022).** Effect of nitrogen and zinc nano fertilizer with the organic farming practices on cereal and oil seed crops. *Scientific reports*. Apr 28;12(1):1-7.
9. **Kumar, V., Kandpal, B.K., Dwivedi, A., Kumar, S.V., Kumar, V. and Sharma, D.K. (2016).** Effect of nitrogen and zinc fertilizer rates on growth, quality and yield of Indian Mustard (*Brassica juncea* L.). *International Journal of Agril. Sci.* 8(06) 1031-1035.

10. **Kumar, V., Singh, S. K., & Suman, S. N. (2014).** Zinc-Boron Interaction Effects on Yield, Nutrient Uptake and Quality of Mustard (*Brassica juncea* L.) in Ustifluvents. *RAU Journal of Research*, 24(1-2), 59-63.
11. **Moaveni, P. and Kheiri, T. (2011).** In 2nd International Conference on Agricultural and Animal Science; November 25–27, 2011 in Maldives. Singapore: IACSIT Press 22:160–163.
12. **Mukherjee, D. (2010).** Productivity, profitability and apparent nutrient balance under different crop sequence in mid- hill condition. *Indian Journal of Agricultural Sciences*, 80(5):4 20-22.
13. **Naderi, M.R. and Danesh-Shahraki, A. (2013).** Nano fertilizers and their roles in sustainable agriculture. *International Journal of Agriculture and Crop Sciences* 5, 2229–2232.
14. **Olsen, S.R, Cole, C.V., Watanable, F. S. and Dean, L. A. (1954).** Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA, Cric.* 930:19- 23
15. **Pandey, S. B., Sachan, R., Patel, V., & Bhawan, R. (2022).** Effect of sulphur, potassium and PSB on growth parameters, root architecture and quality of mustard (*Brassica juncea* L.). *International Journal of Plant & Soil Science*, 34(24), 122-127.
16. **Rathore, A. (2015).** Boron: An important element for agricultural productivity. *Asian Journal of Multidisciplinary Studies*, 3(2), 26-28.
17. **Sachan, R., Thomas, T., Pandey, H. P., Tiwari, A., Kumar, A., Yadav, P., ... & Kumar, M. (2022).** Effect of STCR-IPNS based nutrient application on soil health, yield, nutrient content and uptake of mustard (*Brassica juncea* L.) in eastern plain zone of Uttar Pradesh, India. *International Journal of Plant & Soil Science*, 34(20), 248-258.
18. **Sharma, S., Chaudhry, S., & Singh, R. (2020).** Effect of boron and sulphur on growth, yield and nutrient uptake of mustard (*Brassica juncea* L.). *International Journal of Chemical Studies*, 8(4), 1998-2001.
19. **Shorna, S. I., Polash, M. A. S., Sakil, M. A., Mou, M. A., Hakim, M. A., Biswas, A., & Hossain, M. A. (2020).** Effects of nitrogenous fertilizer on growth and yield of Mustard Green. *Tropical Plant Research*, 7(1), 30-36.
20. **Singh, M. K., Singh, S. K., Kumar, A., & Kumari, R. (2023).** Evaluation of varying levels of sulphur and zinc on mustard [*Brassica juncea* (L.) Czern & Coss] yield, nutrient uptake and quality.

21. **Sinha, T., Mishra, A., Mishra, U. S., Sachan, R., & Singh, D. (2022).** Interaction effect of sulphur and boron on growth characteristics, yield components and productivity parameters of mustard (*Brassica juncea* L.) under rainfed condition of Chitrakoot Region. *International Journal of Plant & Soil Science*, 34(22), 1329-1336.
22. **Subbiah, B.V. and Asija, C.L. (1956).** A rapid procedure for the estimation of available N in Soil. *Curr. Sci.* 25:259-260.
23. **Subramanian, K. S., Manikandan, A., Thirunavukkarasu, M. and Rahale, C. S. (2015).** Nano fertilizers for balanced crop nutrition. In: Rai, M, Ribeiro, C., Mattaso, L. and Duran, N.(eds) Nanotechnologies in food and agriculture. Springer, Switzerland, 69- 80.
24. **USDA. (2020).** Oilseeds: world markets and trade. United States of Department of Agriculture, Foreign Agricultural Service. Available at [Usdagov / psdonline / circulars / oilseeds](http://usdagov/psdonline/circulars/oilseeds).
25. **Walkley, A. and Black, C. S.A. (1934).** Old piper, S.S. soil and plant analysis. *Soil Sci.* 37:29- 38.
26. **Yadav, S. N., Singh, S. K., & Kumar, O. (2016).** Effect of boron on yield attributes, seed yield and oil content of mustard (*Brassica juncea* L.) on an Inceptisol. *Journal of the Indian Society of Soil Science*, 64(3), 291-296.